



amidea CHEMISTRY

(Designed according to CBSE Question Paper)

Online Resource Centre www.xamidea.in

Region-wise CBSE

Examination Papers 2016

(Solved)

- Remembering & Understanding Based Questions
- Migher Order Thinking Skills (HOTS)
- Application Based Questions
- Value Based Questions
- Proficiency Exercise

CHEMISTRY

Examination Papers 2008–2015

CONTENT

CBSE Examination Paper–2008 (Delhi)	3
CBSE Examination Paper–2008 (All India)	23
CBSE Examination Paper–2009 (Delhi)	43
CBSE Examination Paper–2009 (All India)	69
CBSE Examination Paper–2009 (Foreign)	92
CBSE Examination Paper–2010 (Delhi)	116
CBSE Examination Paper–2010 (All India)	134
CBSE Examination Paper–2010 (Foreign)	152
CBSE Examination Paper–2011 (Delhi)	171
CBSE Examination Paper–2011 (All India)	191
CBSE Examination Paper–2011 (Foreign)	209
CBSE Examination Paper–2012 (Delhi)	229
CBSE Examination Paper–2012 (All India)	249
CBSE Examination Paper–2012 (Foreign)	267
CBSE Examination Paper–2013 (Delhi)	285
CBSE Examination Paper–2013 (All India)	300
CBSE Examination Paper–2013 (Foreign)	315
CBSE Examination Paper–2014 (Delhi)	331
CBSE Examination Paper–2014 (All India)	349
CBSE Examination Paper–2014 (Foreign)	367
CBSE Examination Paper–2015 (Delhi)	383
CBSE Examination Paper–2015 (Ajmer)	395
CBSE Examination Paper–2015 (Allahabad)	404
CBSE Examination Paper–2015 (Bhubaneshwar)	414
CBSE Examination Paper–2015 (Chennai)	425
CBSE Examination Paper–2015 (Guwahati)	435
CBSE Examination Paper–2015 (Panchkula)	445
CBSE Examination Paper–2015 (Patna)	455
CBSE Examination Paper–2015 (Foreign)	466

CBSE EXAMINATION PAPERS

DELHI-2008

Time allowed: 3 hours Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Delhi) SET-I

- 1. What is the coordination number of each type of ions in a rock-salt type crystal structure?
- 2. Define the term 'order of reaction' for chemical reactions.
- 3. What causes Brownian movement in a colloidal solution?
- **4.** In which one of the two structures, NO_2^+ and NO_2^- , the bond angle has a higher value?
- 5. Write the IUPAC name of the following compound:

6. Arrange the following compounds in an increasing order of their acid strengths:

- 7. Write a chemical reaction in which the iodide ion replaces the diazonium group in a diazonium salt.
- 8. Name a substance that can be used as an antiseptic as well as a disinfectant.
- Explain as to why haloarenes are much less reactive than haloalkanes towards nucleophilic substitution reactions.

OR

Which compound in each of the following pairs will react faster in S_N 2 reaction with —OH? Why?

(i) CH₃Br or CH₃I

- (ii) (CH₃)₃ CCl or CH₃Cl
- **10.** (a) State the IUPAC name of the following compound:

$$H_3C$$
 H
 H
 H
 H
 H
 H

(b) Complete the following chemical equation:

$$CH_3CH_2CH = CH_2 + HBr \xrightarrow{peroxide} \dots$$

- 11. State Henry's law correlating the pressure of a gas and its solubility in a solvent and mention two applications of the law.
- 12. A first order decomposition reaction takes 40 minutes for 30% decomposition. Calculate its $t_{1/2}$ value.

4 | Xam idea Chemistry-XII

- 13. What is meant by the 'rate constant, k' of a reaction? If the concentration is expressed in mol L⁻¹ units and time in seconds, what would be the units for k (i) for a zero order reaction and (ii) for a first order reaction?
- **14.** Define the following terms in relation to proteins:
 - (i) Peptide linkage
- (ii) Denaturation
- 15. List the reactions of glucose which cannot be explained by its open chain structure.
- 16. Assign a reason for each of the following statements:
 - (i) Ammonia is a stronger base than phosphine.
 - (ii) Sulphur in vapour state exhibits a paramagnetic behaviour.
- 17. Draw the structure of the following molecules:
 - (*i*) SF₄

- (ii) XeF₄
- 18. What are biodegradable and non-biodegradable detergents? Give one example of each class.
- **19.** What is a semiconductor? Describe the two main types of semiconductors and explain mechanisms for their conduction.
- **20.** Calculate the temperature at which a solution containing 54 g of glucose, $(C_6H_{12}O_6)$, in 250 g of water will freeze. $(K_f \text{ for water } = 1.86 \text{ K mol}^{-1} \text{ kg})$
- **21.** What are lyophilic and lyophobic sols? Give one example of each type. Which one of these two types of sols is easily coagulated and why?
- 22. State briefly the principles which serve as basis for the following operations in metallurgy:
 - (i) Froth floatation process
 - (ii) Zone refining
 - (iii) Refining by liquation
- 23. Write chemical equations for the following processes:
 - (i) Chlorine reacts with a hot concentrated solution of sodium hydroxide.
 - (ii) Orthophosphorous acid is heated
 - (iii) PtF₆ and xenon are mixed together.

OR

Complete the following chemical equations:

- (i) $\operatorname{Ca}_{3}\operatorname{P}_{2}(s) + \operatorname{H}_{2}\operatorname{O}(l) \longrightarrow \dots$
- (ii) $\operatorname{Cu}^{2+}(aq) + \operatorname{NH}_3(aq) \longrightarrow \dots$ (excess)
- $(iii) F_2(g) + H_2O(l) \longrightarrow ...$
- **24.** (a) What is a ligand? Give an example of a bidentate ligand.
 - (b) Explain as to how the two complexes of nickel, $[Ni(CN)_4]^{2-}$ and $Ni(CO)_4$, have different structures but do not differ in their magnetic behaviour. (Ni = 28)
- 25. Name the reagents which are used in the following conversions:
 - (i) A primary alcohol to an aldehyde
 - (ii) Butan-2-one to butan-2-ol
 - (iii) Phenol to 2, 4, 6-tribromophenol
- **26.** Account for the following observations:
 - (i) pK_b for aniline is more than that for methylamine.
 - (ii) Methylamine solution in water reacts with ferric chloride solution to give a precipitate of ferric hydroxide.
 - (iii) Aniline does not undergo Friedel-Crafts reaction.

- 27. Write the names and structure of the monomers of the following polymers:
 - (i) Buna-S
- (ii) Neoprene

(iii) Nylon-6

28. Conductivity of 0.00241 M acetic acid solution is 7.896×10^{-5} S cm⁻¹. Calculate its molar conductivity in this solution. If Λ_m° for acetic acid is 390.5 S cm² mol⁻¹, what would be its dissociation constant?

OR

Three electrolytic cells A, B and C containing solutions of zinc sulphate, silver nitrate and copper sulphate, respectively are connected in series. A steady current of 1.5 ampere was passed through them until 1.45 g of silver were deposited at the cathode of cell B. How long did the current flow? What mass of copper and what mass of zinc were deposited in the concerned cells? (Atomic masses of Ag = 108, Zn = 65.4, Cu = 63.5)

- **29.** Assign reasons for the following:
 - (i) The enthalpies of atomisation of transition elements are high.
 - (ii) The transition metals and many of their compounds act as good catalysts.
 - (iii) From element to element the actinoid contraction is greater than the lanthanoid contraction.
 - (iv) The E° value for the Mn³⁺ / Mn²⁺ couple is much more positive than that of Cr³⁺ / Cr²⁺.
 - (ν) Scandium (Z = 21) does not exhibit variable oxidation states and yet it is regarded as a transition element.

OR

- (a) What may be the possible oxidation states of the transition metals with the following d electronic configurations in the ground state of their atoms:
 - $3d^3 4s^2$, $3d^5 4s^2$ and $3d^6 4s^2$. Indicate relative stability of oxidation states in each case.
- (b) Write steps involved in the preparation of (i) Na₂CrO₄ from chromite ore and (ii) K₂MnO₄ from pyrolusite ore.
- **30.** (a) Complete the following reaction statements by giving the missing starting material, reagent or product as required:

- (b) Describe the following reactions:
 - (i) Cannizaro reaction
- (ii) Cross aldol condensation

OR

- (a) How would you account for the following:
 - (i) Aldehydes are more reactive than ketones towards nucelophiles.
 - (ii) The boiling points of aldehydes and ketones are lower than of the corresponding acids.
 - (iii) The aldehydes and ketones undergo a number of addition reactions.
- (b) Give chemical tests to distinguish between:
 - (i) Acetaldehyde and benzaldehyde
 - (ii) Propanone and propanol.

CBSE (Delhi) SET-II

Questions Uncommon to Set-I

- 1. What is the total number of atoms per unit cell in a face-centred cubic (fcc) structure?
- 2. What is primary cell? Give an example.
- **6.** Write the IUPAC name of the following compound:

9. State Raoult's law for solutions of volatile liquids. Taking suitable examples explain the meaning of positive and negative deviations from Raoult's law.

OR

Define the term osmotic pressure. Describe how the molecular mass of a substance can be determined by a method based on measurement of osmotic pressure?

- **10.** The conductivity of a 0.20 M solution of KCl at 298 K is 0.0248 S cm⁻¹. Calculate its molar conductivity.
- 11. Formulate the galvanic cell in which the following reaction takes place:

$$Zn(s) + 2Ag^{+}(aq) \longrightarrow Zn^{2+}(aq) + 2Ag(s)$$

State

- (i) Which one of its electrodes is negatively charged?
- (ii) The reaction taking place at each of its electrode.
- (iii) The carriers of current within this cell.
- **14.** Complete the following reaction equations:

(i)
$$C_6H_5N_2Cl + KI \longrightarrow ...$$

(ii)
$$H$$
 $C = C$ H $+ Br_2$ $\xrightarrow{CCl_4}$...

- **15.** (i) Why is it that haloalkanes are more reactive than haloarenes towards nucleophiles.
 - (ii) Which one of the following reacts faster in an S_N 1 reaction and why?

- **28.** (a) Derive the general form of the expression for the half-life of a first order reaction.
 - (b) The decomposition of NH₃ on platinum surface is a zero order reaction. What are the rates of production of N₂ and H₂ if $k = 2.5 \times 10^{-4} \text{ mol}^{-1} \text{ L s}^{-1}$?

OR

- (a) List the factors on which the rate of a chemical reaction depends.
- (b) The half-life for decay of radioactive ¹⁴ C is 5730 years. An archaeological artefact containing wood has only 80% of the ¹⁴ C activity as found in living trees. Calculate the age of the artefact.
- **29.** (a) How will you bring about the following conversions?
 - (i) Ethanol to acetone
- (ii) Benzene to acetophenone
- (iii) Benzoic acid to benzaldehyde
- (b) Describe the following giving a suitable example in each case:
 - (i) Decarboxylation

(ii) Cannizaro's reaction

- (a) An organic compound contains 69.77% carbon, 11.63% hydrogen and the rest is oxygen. The molecular mass of the compound is 86. It does not reduce Tollens' reagent but forms an addition compound with sodium hydrogen sulphite and gives positive iodoform test. On vigorous oxidation it gives ethanoic and propanoic acids. Deduce the possible structure of the organic compound.
- (b) State reasons for the following:
 - (i) Monochloroethanoic acid has a higher pKa value than dichloroethanoic acid.
 - (ii) Ethanoic acid is a weaker acid than benzoic acid.

CBSE (Delhi) SET-III

Questions Uncommon to Set-I and Set-II

- 1. What type of substances exhibit antiferromagnetism?
- 2. Express the relation between conductivity and molar conductivity of a solution.
- 3. Which has a higher enthalpy of adsorption, physisorption or chemisorption?
- 10. The resistance of a conductivity cell containing 0.001 M KCl solution at 298 K is 1500 Ω . What is the cell constant if the conductivity of 0.001 M KCl solution at 298 K is 0.146×10^{-3} S cm⁻¹?
- 19. How would you account for the following?
 - (i) Frenkel defects are not found in alkali metal halides.
 - (ii) Schottky defects lower the density of related solids.
 - (iii) Impurity doped silicon is a semiconductor.

OR

Explain the following properties giving suitable examples:

- (i) Ferromagnetism
- (ii) Paramagnetism
- (iii) Ferrimagnetism
- 21. Explain the basic principles of following metallurgical operations:
 - (i) Zone refining
 - (ii) Vapour phase refining
 - (iii) Electrolytic refining
- 22. Explain what is observed when
 - (i) an electrolyte, KCl, is added to a hydrated ferric oxide sol.
 - (ii) an electric current is passed through a colloidal solution.
 - (iii) a beam of strong light is passed through a colloidal solution.


SOLUTIONS

CBSE (Delhi) SET-I

- 1. Coordination number of Na^+ ion = 6
 - Coordination number of Cl ion = 6
- 2. The sum of powers of the concentrations of the reactants in the rate law expression is called the order of reaction.
- 3. This is due to the unequal bombardment of colloidal particles by the molecules of dispersion medium
- **4.** NO_2^+ has higher bond angle as the central atom nitrogen in NO_2^- has a lone pair of electrons.
- 5. 2, 5-Dimethyl hexane-1, 3-diol.

7.
$$\stackrel{+}{\bigvee_{N_2}Cl}$$
 + KI $\stackrel{\Delta}{\longrightarrow}$ $\stackrel{I}{\bigvee_{Iodobenzene}}$ + KCl + N₂

- 8. Phenol, 0.2% solution of phenol acts as antiseptic where as 1% solution acts as disinfectant.
- **9.** In haloarenes C—X bond acquires a partial double bond character due to resonance. As a result the bond cleavage in haloarenes is difficult than haloalkanes and therefore, they are less reactive towards nucleophilic substitution reaction.



- (i) CH₃-I reacts faster than CH₃-Br as iodine is a better leaving group because of its larger size.
- (ii) CH₃-Cl (1° halide) reacts faster than (CH₃)₃ CCl (3° halide) since in case tertiary butyl chloride three bulky methyl group hinder the approaching nucleophile.
- **10.** (*a*) 1-Bromo but-2-ene.

(b)
$$CH_3 - CH_2 - CH = CH_2 + H - Br \xrightarrow{Peroxide} CH_3 - CH_2 - CH_2 - CH_2 - Br$$

But-1-ene

1-bromo butane

11. It states that at constant temperature the mass of a gas(m) dissolved in a given volume of the liquid is directly proportional to the pressure of the gas(P) present in equilibrium with the liquid.

Mathematically,
$$m \propto P$$

 $m = K_H P$

where K_H is the Henry's law constant.

Applications of Henry's law are

- (i) To increase the solubility of CO₂ in soft drinks and soda water, the bottle is sealed under high
- (ii) To minimize the painful effects accompanying the decompression of deep sea divers, oxygen diluted with less soluble helium gas is used as breathing gas.
- 12. For a first order reaction

$$K = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$
when $t = 40$ minutes, $\frac{[R]_0}{[R]} = \frac{100}{100 - 30} = \frac{10}{7}$

$$K = \frac{2.303}{40} \log \frac{10}{7} = \frac{2.303}{40} \log 1.428 = \frac{2.303}{40} \times 0.1548$$

$$K = 8.91 \times 10^{-3} \text{ min}^{-1}$$

$$t_{1/2} = \frac{0.693}{K} = \frac{0.693}{8.91 \times 10^{-3}}$$

 $t_{1/2} = 77.78$ min.

13. Rate constant is the rate of the reaction when the concentration of each reactant is taken as unity.

$$\therefore$$
 rate = $K[A]^n$

$$\therefore \qquad \text{General unit of } K = \left(\frac{\text{mol.}}{\text{lit.}}\right)^{1-n} \text{ s}^{-1}$$

(i) For a zero order reaction n = 0


$$\therefore$$
 Unit of $K = \text{mol lit}^{-1} \text{ sec}^{-1}$

(iii) For a first order reaction n = 1


$$\therefore$$
 Unit of $K = \sec^{-1}$

14. (i) Peptide linkage: A peptide linkage is an amide linkage (-CONH-) formed between —COOH group of one α-amino acid and NH₂ group of the other amino acid by the elimination of a water molecule.


- (ii) **Denaturation:** When a protein in its native form is subjected to physical change like change in temperature or chemical change like change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and proteins loses its biological activity. During denaturation 2° and 3° structures of proteins are destroyed but 1° structure remains intact, *e.g.*, coagulation of egg white on boiling.
- 15. The following reactions of glucose cannot be explained by its open chain structure.
 - (i) Despite having the aldehyde group glucose does not give 2, 4-DNP test, Schiffs test and it does not form the hydrogen sulphite addition product with NaHSO₃.
 - (ii) The pentacetate of glucose does not react with hydroxylamine indicating the absence of free —CHO group.
 - (iii) When D-glucose is treated with methanol in the presence of dry hydrogen chloride gas, it gives two isomeric mono methyl derivatives known as α -D glucoside and methyl β -D glucoside. These glucosides do not react with hydrogen cyanide or with hydroxylamine.
- **16.** (*i*) As the atomic size of nitrogen is smaller than phosphorus, therefore electron density on nitrogen atom is higher than that on phosphorus atom. Consequently the tendency of N in NH₃ to donate its lone pair of electrons is much higher than that of P in PH₃. Thus, ammonia is stronger base than phosphine.
 - (ii) In vapour state sulphur partly exists as S_2 molecule which has two unpaired electrons in the antibonding π^* orbitals like O_2 and, hence exhibits paramagnetic behaviour.
- 17. (i) sp^3d hybridisation.



(ii) sp^3d^2 hybridisation



18. Biodegradable detergents: Detergents having straight hydrocarbon chains are easily degraded by micro-organism and hence called biodegradable detergents, *e.g.*, sodium–4–(1-dodecyl) benzene sulphonate.



Non-biodegradable detergents: Detergents having branched hydrocarbon chains are not easily degraded by micro-organisms and hence are called non-biodegradable detergents, e.g., sodium-4-(1, 3, 5, 7-tetramethyl octyl) benzenesulphonate.

$$CH_3$$
 CH_3 CH_3 CH_3 CH_3 CH_4 CH_5 CH_5 CH_5 CH_6 CH_7 CH_8 CH_8

Non-biodegradable detergents accumulate in rivers and waterways thereby causing water pollution.

19. Semiconductor: These are the solids with conductivities in the intermediate range from 10^{-6} to $10^4 \text{ ohm}^{-1} \text{ m}^{-1}$.


Semiconductors are of two types

(i) *n*-type of semiconductors

(ii) p-type of semiconductors

n-type semiconductor: When a silicon or germanium crystal is doped with group 15 element like P or As, the dopant atom forms four covalent bonds like a Si or Ge atom but the fifth electron, not used in bonding, becomes delocalised and contribute its share towards electrical conduction. Thus silicon or germanium doped with P or As is called n-type semiconductor, n indicative of negative, since it is the electron that conducts electricity.

p-type semiconductor: When silicon or germanium is doped with group 13 element like B or Al, the dopant has only with three, valence electrons. An electron vacancy or a hole is created at the place of the missing fourth electron. Here, this hole moves through the crystal like a positive charge giving rise to electrical conductivity. Thus Si or Ge doped with B or Al is called p type of semiconductor, (P stands for positive hole) since it is the positive hole that is responsible for conduction.



20. Mass of glucose $(W_R) = 54 \text{ g}$

Molecular mass of glucose $(M_R) = 180$

Mass of water $(W_B) = 250 \text{ g}$

 K_f for water = 1.86 k mol⁻¹ kg

Applying the formula,
$$\Delta T_f = \frac{K_f \times W_B \times 1000}{M_B \times W_A}$$

$$\Delta T_f = \frac{1.86 \times 54 \times 1000}{180 \times 250} = 2.23$$

$$T_f = T^{\circ}_f - \Delta T_f = 0 - (2.23)$$

$$T_f = -2.23^{\circ} \text{ C}$$

- **21. Lyophilic sols:** Lyophilic sols are those sols in which the particles of dispersed phase have great affinity for the dispersion medium, *e.g.*, sols of gum, gelatine, starch, etc.
 - **Lyophobic sols:** In this type of sols the particles of dispersed phase have little or no affinity for the dispersion medium, e.g., gold sol, Fe (OH)₃ sol, As₂O₃ sol., etc.
 - Lyophobic sols easily coagulate on the addition of small amount of electrolyte because these are not stable. The stability of Lyophobic sols is only due to the presence of charge on the colloidal particles, on the other hand stability of lyophilic sol. is due to charge as well as solvation of colloidal particles.
- **22.** (*i*) The principle of froth floatation process is that sulphide ore particle are preferentially wetted by pine oil, whereas the gangue particles are wetted by water.
 - (ii) Zone refining is based on the principle that the impurities are more soluble in (liquid state) than in the solid state of the metal.
 - (iii) The principle of refining by liquation is that the impurities whose melting points are higher than the metal are left behind on melting the impure metal. Hence pure metal separates out.

23. (i)
$$3Cl_2 + 6NaOH \longrightarrow 5NaCl + NaClO_3 + 3H_2O$$

$$\Delta$$

$$(ii) 4H3PO3 \xrightarrow{\Delta} 3H3PO4 + PH3$$

$$(iii) PtF_6 + Xe \longrightarrow Xe^+ [PtF_6]^-$$

(i)
$$\operatorname{Ca}_{3}\operatorname{P}_{2}(s) + 6\operatorname{H}_{2}\operatorname{O}(l) \longrightarrow 2\operatorname{PH}_{3} + 3\operatorname{Ca}\left(\operatorname{OH}\right)_{2}$$

(ii)
$$Cu^{2+}(aq) + 4NH_3(aq) \rightleftharpoons [Cu(NH_3)_4]^{2+}(aq)$$


(blue) (excess) (deep blue)

(iii)
$$2F_2(g) + 2H_2O(1) \longrightarrow 4H^+(aq) + 4F^-(aq) + O_2(g)$$

24. (a) The ion, atom or molecule bound to the central atom/ion in the coordination entity is called ligand. A ligand should have lone pair of electrons in their valence orbital which can be donated to central metal atom/ion.


Bidentate ligand:
$$H_2$$
 N— CH_2 — CH_2 — N H_2 (Ethylene diamine)

(b)



Structure: Square planar.

Magnetic behaviour: Diamagnetic due to the absence of unpaired electrons.



Structure: Tetrahedral

Magnetic behaviour: Diamagnetic due to the absence of unpaired electrons.

- **25.** (i) Pyridinium chlorochromate (C₅H₅ NHCrO₃Cl⁻) or Cu/573 K
 - (ii) LiAlH₄ / ether (iii) Br₂ / H₂O
- **26.** (i) In aniline due to resonance the lone pair of electrons on the nitrogen atom are delocalized over the benzene ring. As a result, the electron density on the nitrogen decreases. On the other hand, in methyl amine +I effect of CH3 increases the electron density on the nitrogen atom. Therefore aniline is a weaker base than methyl amine and hence its pK_h value is higher than that of methyl amine.

(ii)
$$CH_3$$
— NH_2 + H_2O \Longrightarrow CH_3 NH_3 + OH^-

Due to alkaline nature of solution of methylamine precipitation of $Fe(OH)_3$ occurs.

FeCl₃ +3OH⁻
$$\longrightarrow$$
 Fe (OH)₃ \downarrow +3Cl⁻
Ferric hydroxide
(Brown ppt)

(iii) Aniline being a Lewis base, reacts with lewis acid AlCl₃ to form a salt. Due to this N atom of aniline acquires positive charge and hence acts as a strong deactivation group for further reaction.

27.

	Polymer	Monomer	Structure of Monomer
(i)	Buna-S	Buta-1, 3-diene	$CH_2 = CH$ — $CH = CH_2$
			$CH = CH_2$
		and styrene	
(ii)	Neoprene	Chloroprene	$H_2C = C - CH = CH_2$
			 Cl
			Q Cr
(iii)	Nylon-6	Caprolactum	C—NH
			CH ₂ CH ₂
			CH ₂ CH ₂
			CH ₂

28.
$$c = 0.00241 \text{ M}, K = 7.896 \times 10^{-5} \text{ S cm}^{-1}, \lambda_m^{\infty} = 390.5 \text{ S cm}^2 \text{ mol}^{-1}$$
$$\lambda_m = \frac{K \times 1000}{c}$$

Substituting the values

$$\lambda_m = \frac{7.896 \times 10^{-5} \times 1000}{0.002} = 32.76 \text{ S cm}^2 \text{ mol}^{-1}$$
$$\alpha = \frac{\lambda_m^c}{\lambda_m^\infty} = \frac{32.76}{390.5} = 0.084$$

$$\alpha = 8.4\%$$

CH₃ — COOH
$$\Longrightarrow$$
 CH₃ —COO⁻ + H⁺

$$c \qquad 0 \qquad 0$$

$$c(1-\alpha) \qquad c\alpha \qquad c\alpha$$

$$K_a = \frac{[\text{CH}_3 - \text{COO}^-][\text{H}^+]}{[\text{CH}_3 - \text{COOH}]} = \frac{c\alpha \cdot c\alpha}{c(1-\alpha)} = \frac{c\alpha^2}{1-\alpha}$$

$$K_a = \frac{0.0241(0.084)^2}{(1-0.084)} = 1.86 \times 10^{-5}$$
OR

$$Ag^+ + e^- \longrightarrow Ag$$

108 g of Ag are deposited by 96500 C

:. 1.45 g of Ag will be deposited by =
$$\frac{96500}{108} \times 1.45 \text{ C}$$

= 1295.6 C

$$t = \frac{Q}{I} = \frac{1295.6}{1.50} = 863.7 \text{ s.}$$

$$\text{Cu}^{2+} + 2e^{-} \longrightarrow \text{Cu}$$

$$2 \times 96500$$
 C deposit Cu = 63.5 g

:. 1295.6 C deposit Cu =
$$\frac{63.5}{2 \times 96500} \times 1295.6 = 0.426 \text{ g}$$

$$\operatorname{Zn}^{2+} + 2e^{-} \longrightarrow \operatorname{Zn}$$

$$2 \times 96500$$
 C deposits Zn = 65.3 g

$$\therefore 1295.6 \text{ C deposit Zn} = \frac{65.3}{2 \times 96500} \times 1295.6 = 0.438 \text{ g}$$

- 29. (i) This is because transition metals have strong metallic bonds as they have a large number of unpaired electrons.
 - (ii) The catalytic activity of transition metals is attributed to the following reasons:
 - (a) Because of their variable oxidation states transition metals form unstable intermediate compounds and provide a new path with lower activation energy for the reaction.
 - (b) In some cases, the transition metal provides a suitable large surface area with free valencies on which reactants are adsorbed.
 - (iii) This is due to poorer shielding by 5f electrons in actinoids than that by 4f electron in the lanthanoids.

- (iv) This is because half filled d-subshell $(3d^5)$ in Mn²⁺ is more stable.
- (v) This is because scandium has partially filled d orbitals in the ground state $(3d^{1}4s^{2})$.

OR

(*a*)

Electronic Configuration	Element	Possible O.S.	More stable O.S.
$3d^24s^2$	Vanadium	+ 2, + 3, + 4, + 5	+ 5
$3d^5 4s^2$	Manganese	+ 2, + 3, + 4, + 5, + 6, + 7	+ 2, + 7
$3d^{6}4s^{2}$	Iron	+ 2, + 3, + 4, + 6	+ 2, + 3

(b) (i) Chromite ore is fused with sodium carbonate in excess of air.

(ii) Pyrolusite ore (MnO_2) is fused with KOH in the presence of O_2 or oxidising agent such as KNO₃.

$$2MnO_2$$
 + $4KOH$ + O_2 \longrightarrow $2K_2MnO_4$ + $2H_2O$ Pyrolusite ore

(ii) BH_3 / THF , H_2O_2 / OH^- ,

(b) (i) Cannizzaro reaction: Aldehydes which do not have an α-hydrogen, undergo self oxidation and reduction (disproportionation) reaction on treatment with concentrated alkali. In this reaction one molecule of the aldehyde is reduced to alcohol while another is oxidised to carboxylic acid salt.

$$\begin{array}{ccc}
2\text{HCHO} & \xrightarrow{\text{conc. KOH}} \\
\text{Formaldehyde} & \xrightarrow{\text{CH}_3} & \text{OH} + \text{HCOO}^-\text{K}^+ \\
& & \text{Methyl alcohol} & \text{Potasium formate}
\end{array}$$

(ii) Cross aldol condensation: When aldol condensation is carried out between two different aldehydes and/or ketones, it is called cross aldol condensation.

$$\begin{array}{c|c}
C & O & O \\
 & & C \\
\hline
C & CH_3 & OH^- \\
\hline
C & CH_3 & CH = CH \\
\hline
C & CH \\
\hline
C & CH \\
C & CH \\
\hline
C & CH \\
C & CH \\
\hline
C & CH \\
C & CH \\
\hline
C & CH \\
C & CH \\
\hline
C & CH \\
C & CH \\
\hline
C & CH \\
C & CH \\
\hline
C & CH \\
C$$

Benzaldehyde


Acetophenone

Benzal acetophenone

If both of them contain α-hydrogen atoms. It gives a mixture of four products.

OR

- (a) (i) This is due to steric and electronic reasons. Sterically, the presence of two relatively large substituents in ketones hinders the approach of nucleophile to carbonyl carbon than in aldehydes having only one such substituent. Electronically two alkyl groups reduce the positivity of the carbonyl carbon more effectively in ketones than in aldehydes.
 - (ii) This is due to intermolecular hydrogen bonding in carboxylic acids.
 - (iii) Due to greater electronegativity of oxygen than carbon the C atom of the \geq C = O bond acquires a partial positive charge in aldehydes and ketones and hence readily undergo nucleophilic addition reactions.
- (b) (i) Acetaldehyde reacts with NaOI (I2 / NaOH) to form yellow ppt of iodoform while benzaldehyde does not give this test.



(ii) Propanone give orange-red ppt with 2, 4-DNP reagent and yellow ppt of iodoform with sodium hypoiodite, whereas 1-propanol does not give these tests.

- **SE (Delni) SEI-II 1.** 8 (corner atoms) $\times \frac{1}{8}$ atom per unit cell + 6 (face atoms) $\times \frac{1}{2}$ atom per unit cell = 1 + 3 = 4
- A primary cell is one in which the redox reaction occurs only once and the cell becomes dead after some time and cannot be used again, e.g., dry cell.
- Pentane-2, 4-dione.
- Raoult's law: It states that for a solution of volatile liquids the partial pressure of each component is directly proportional to its mole fraction.

Mathematically

$$\begin{array}{ll} P_A \propto x_A & P_B \propto x_B \\ P_A = P_A^{\rm o} x_A & P_B = P_B^{\rm o} x_B \end{array}$$

Positive deviation from Raoult's law: In this type of deviation the partial pressure of each component of solution is greater than that calculated from Raoult's law, i.e., $P_A > P_A^0 x_A$ & $P_B > P_B^{\,o} x_B$. Example: A solution of water and ethanol.

Negative deviation from Raoult's: In this type of deviation the partial pressure of each component of solution is less than that expected from Raoult's law, i.e., $P_A < P_A^0 x_A & P_B < P_B^0 x_B$.

Example: A solution of acetone and chloroform.

Osmotic pressure (π) is defined as the extra pressure that must be applied to the solution side in order to prevent the flow of solvent molecules into it through a semipermeable membrane.

$$\pi = \frac{n_B}{V} RT = CRT$$

where V is the volume of solution in litres containing n_B moles of the solute.

If W_B grams of the solute whose molecular mass M_B is present in the solution then

$$\pi = \frac{W_B RT}{M_B RT}$$

$$M_B = \frac{W_B RT}{\pi V}$$

Thus, knowing W_B , T, π and V molecular mass of the solute, M_B can be calculated. 10. $\kappa = 0.0248 \text{ S cm}^{-1}$, $M = 0.2 \text{ mol L}^{-1}$

Substituting the values

ting the values
$$\Lambda_m = \frac{\kappa \times 1000}{M} = \frac{0.0248 \times 1000}{0.2}$$

$$\Lambda_m = 1248 \text{ cm}^2 \text{ mol}^{-1}$$

$$\Lambda_m = 1248 \text{ cm}^2 \text{ mol}^{-1}$$

11.
$$\operatorname{Zn} \left| \operatorname{Zn}^{2+} (\operatorname{conc.}) \right| \left| \operatorname{Ag}^{+} (\operatorname{conc}) \right| \operatorname{Ag}$$

- (i) Zn electrode is negatively charged.
- $Zn \longrightarrow Zn^{2+} + 2e^{-}$ (ii) At anode

At cathode

$$2Ag^+ + e^- \longrightarrow 2Ag$$


- (iii) Current carriers of cell are
 - electrons in external wire.
 - Zn²⁺ ions in anodic half cell.
 - Ag + ions in cathodic half cell.
 - Ions of salt bridge, *i.e.*, K⁺ and Cl⁻.

14. (i)
$$C_6H_5N_2Cl + KI \longrightarrow C_6H_5I + KCl + N_2$$
Iodobenzene

(ii)
$$\begin{array}{c} H \\ C = C \\ H \end{array} + Br_2 \xrightarrow{CCl_4} \begin{array}{c} H \\ C - C \\ H \\ Br Br Br \end{array}$$

1, 2-Dibromoethane

15. (i) In haloarenes C—X bond acquires a partial double bond character due to resonance. As a result the bond cleavage in haloarenes is difficult than haloalkanes and therefore, they are less reactive towards nucleophilic substitution reaction.



Compound (I) reacts faster in S_N1 reaction as it is a 2° alkyl halide.

28. (a) Consider the first order reaction

$$R \longrightarrow P$$

$$Rate = -\frac{d[R]}{dt} = K[R]$$

$$\frac{d[R]}{[R]} = -K dt$$

or

Integrating both the sides we get

$$ln[R] = -Kt + C \qquad ...(i)$$

when t = 0, $R = [R]_0$ where $[R]_0$ is the initial concentration of the reactant.

Therefore equation (i) can be written as

$$ln [R]_0 = -K \times 0 + C$$

$$ln [R]_0 = C$$

Substituting the value of *C* in equation (i)

$$ln [R] = -Kt + ln [R]_0$$

$$Kt = ln [R]_0 - ln [R]$$

$$Kt = \frac{ln [R]_0}{[R]}$$

$$Kt = 2.303 \log \frac{[R]_0}{[R]}$$
 \Rightarrow $t = \frac{2.303}{K} \log \frac{[R]_0}{[R]}$

when
$$t = t$$

$$t = t_{1/2} [R] = \frac{[R]_0}{2}$$

$$\therefore \quad t_{1/2} = \frac{2.303}{K} \log \frac{[R]_0}{\frac{[R]_0}{2}}$$

$$t_{1/2} = \frac{2.303}{K} \log 2 = \frac{2.303}{K} \times 0.3010$$

$$\therefore \quad t_{1/2} = \frac{0.693}{K}$$

$$(b) \qquad 2NH_3 \longrightarrow N_2 + 3H_2$$

For a zero order reaction

Rate =
$$K [NH_3]^0$$

Rate =
$$K = 2.5 \times 10^{-4}$$

Rate =
$$-\frac{1}{2} \frac{d [NH_3]}{dt} = \frac{d [N_2]}{dt} = \frac{1}{3} \frac{d [H_2]}{dt}$$

$$\therefore \frac{d [N_2]}{dt} = 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ S}^{-1}$$

$$\frac{d [H_2]}{dt} = \frac{3d [N_2]}{dt} = 3 \times 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ S}^{-1}$$

$$\frac{d [H_2]}{dt} = 7.5 \times 10^{-4} \text{ mol L}^{-1} \text{ S}^{-1}$$

OR

- (a) Rate of reaction depends on
 - (i) Concentration
 - (iii) Nature of reactant
 - (v) Surface area

- (ii) Temperature
- (iv) Pressure of the gaseous reactant
- (vi) Catalyst

(b)
$$t_{1/2} = 5730 \text{ years}$$

$$\therefore K = \frac{0.693}{t_{1/2}} = \frac{0.693}{5730} = 1.209 \times 10^{-4} \text{ year}^{-1}$$

$$t = \frac{2.303}{K} \log \frac{[R]_0}{[R]} = \frac{2.303}{1.2 \times 10^{-4}} \log \frac{100}{80}$$


$$t = \frac{2.303 \times 10^4}{1.2} (\log 10 - \log 8) = \frac{2.303 \times 10^4}{1.2} (1 - 3 \log 2)$$

$$t = \frac{2.303}{1.2} \times 10^4 (1 - 3 \times 0.3010) = \frac{2.303 \times 0.097 \times 10^4}{1.209}$$

$$t = 1847.7 \text{ years}$$

t = 1847.7 years

$$t = 1847.7 \text{ years}$$
29. (a) (i) $CH_3 - CH_2 - OH - \underbrace{K_2Cr_2O_7/H_2SO_4}_{(O)} - CH_3 - CHO \xrightarrow{(O)} CH_3 - COOH$
Ethyl alcohol



(b) (i) Decarboxylation: Carboxylic acids lose CO₂ to form hydrocarbons when their sodium salts are heated with soda lime (NaOH and CaO in the ratio 3: 1).

$$R - COONa \xrightarrow{NaOH \text{ and CaO}} R - H + Na_2CO_3$$

$$COONa \xrightarrow{NaOH/CaO} + Na_2CO_3$$
Sodium benzoate
$$R - H + Na_2CO_3$$

$$R - H + Na_2CO_3$$

$$R - H + Na_2CO_3$$

(ii) Cannizzaro Reaction: Aldehydes which do not have an α-hydrogen atom, undergo disproportionation reaction on treatment. In this reaction, one molecule of the aldehyde is reduced to alcohol while the other is oxidised to carboxylic acid.

$$\begin{array}{c} \text{2HCHO} \\ \text{Formaldehyde} \end{array} \xrightarrow{conc. \ NaOH} \begin{array}{c} \text{conc. NaOH} \\ \end{array} \xrightarrow{CH_3 - OH} + \begin{array}{c} -+ \\ \text{HCOONa} \\ \text{Sodium formate} \end{array}$$

(a) %
$$C = 69.77\%$$
, % $H = 11.63\%$

$$%O = 100 - (69.77 + 11.63) = 18.6\%$$

C:H:O=
$$\frac{69.77}{12}$$
: $\frac{11.63}{1}$ = $\frac{18.6}{16}$ =5.81:11.63:1.16

$$\therefore$$
 C:H:O = 5:10:1

The empirical formula of the given compound is
$$C_5H_{10}O$$
.
$$n = \frac{\text{molecular mass}}{\text{Empirical formula mass}}$$

Molecular mass = 86

Empirical formula mass = $5 \times 12 + 10 \times 1 + 1 \times 16 = 86$

$$n = \frac{86}{81} = 1$$

Molecular formula = n (Empirical formula)

$$= C_5 H_{10} O$$

The (forms addition compound with NaHSO₃) given organic compound is methyl ketone as it gives positive iodoform test and also does not reduce Tollen's reagent.

Since on oxidation it gives ethanoic and propanoic acid its possible structural formula is

Reactions involved:

$$\begin{array}{c} O \\ \parallel \\ \text{CH}_3-\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_3 + \text{NaHSO}_3 \longrightarrow \\ \text{Pentan-2-one} \end{array} \\ \begin{array}{c} \text{CH}_3\text{CH}_2-\text{CH}_2\\ \text{CH}_3 \end{array} \\ \begin{array}{c} \text{OH} \\ \text{addition compound} \end{array}$$

$$\begin{array}{c} O \\ \parallel \\ CH_3-C-CH_2-CH_2-CH_3+3I_2+4NaOH \xrightarrow{Iodoform} \\ CH_3-C-CH_2-CH_2-CH_3+3I_2+4NaOH \xrightarrow{Iodoform} \\ +CH_3-CH_2-CH_2-COONa+3NaI+3H_2O \\ O \\ CH_3-C \xrightarrow{\vdots} CH_2-CH_2-CH_3 \xrightarrow{K_2Cr_2O_7/H_2SO_4} \\ CH_3-C \xrightarrow{\vdots} CH_2-CH_2-CH_3 \xrightarrow{K_2Cr_2O_7/H_2SO_4} \\ CH_3-C \xrightarrow{\vdots} CH_2-CH_2-CH_2-CH_3 \xrightarrow{K_2Cr_2O_7/H_2SO_4} \\ CH_3-C \xrightarrow{\vdots} CH_2-CH_2-CH_2-CH_3 \xrightarrow{K_2Cr_2O_7/H_2SO_4} \\ CH_3-COOH + CH_3-CH_2-COOH \\ CH_3-COOH + CH_3-CH_2-COOH \\ CH_3-COOH + CH_3-CH_2-COOH \\ CH_3-COOH + CH_3-COOH \\ CH_3-COOH + CH_3-COOH \\ CH_3-COOH + CH_3-COOH \\ COOH + CH_3-COOH \\ COOH$$

- (b) (i) This is because dichloroethanoic acid is a stronger acid than monochloroethanoic acid.
 - (ii) This is because methyl group due to its positive inductive effect destabilize the acetate anion by intensifying the negative charge.

CBSE (Delhi) SET-III

1. Substance in which domains are oppositely oriented and cancel out each other magnetic moments.

$$\lambda_m = \frac{\kappa \times 1000}{c}$$

where, $\Lambda_m = \text{Molar conductivity}$ $\kappa = \text{Conductivity}$ c = Molar concentration

3. Chemisorption

10.
$$\kappa = \frac{1}{R} \times \text{cell constant}$$

$$\kappa = 0.146 \times 10^{-3} \text{ S cm}^{-1}, \quad R = 1500 \text{ ohm}$$

$$0.146 \times 10^{-3} = \frac{1}{1500} \times \text{cell constant}$$

$$\text{cell constant} = 0.146 \times 10^{-3} \times 1500 = 219 \times 10^{-3} = 0.219 \text{ cm}^{-1}$$

- 19. (i) This is because alkali metal ions have larger size which cannot fit into interstitial sites.
 - (ii) As the number of ions decreases as a result of Schottky defect, the mass decreases whereas the volume remains the same.
 - (*iii*) This is due to additional electron or creation of hole on dopping with impurity. Creation of hole causes *p*-type semiconductor and creation of electron causes *n*-type semiconductor.

OR

(i) **Ferromagnetism:** Ferromagnetic substances are those substance which are strongly attracted by external magnetic fielding, *e.g.*, (iron, cobalt, nickle and CrO₂, etc.] Ferromagnetism arises due to spontaneous alignment of magnetic moments in the same direction.



Alignment of magnetic moments in ferromagnetic substance.

- (ii) **Paramagnetism:** Paramagnetic substances are those substances which are weakly attracted by magnetic field. It is due to the presence of one or more unpaired electrons, *e.g.*, O₂, Fe³⁺, Cr³⁺, etc.
- (iii) **Ferrimagnetism:** Ferrimagnetism is observed when the magnetic moments of the domains in the substance are aligned in parallel and antiparallel directions in unequal number resulting in some net magnetic moment, *e.g.*, Fe₃O₄ (magnetite) and ferrites like MgFe₂O₄, etc.



- **21.** (*i*) Zone refining is based on the principle that the impurities are more soluble in the melt in than solid state of metal.
 - (ii) Vapour phase refining.

In this, metal is converted into its volatile compound and collected elsewhere. It is the decomposed to give pure metal. So, the two requirements are:

- (a) the metal should form a volatile compound with an available reagent.
- (b) the volatile compound should be easily decomposable, so that the recovery is easy.
- (iii) In electrolytic refining impure metal is made to act as anode. A strip of the same metal in pure form is used as cathode. They are put in a suitable electrolytic bath containing soluble salt of the same metal. When electric current is passed, impure metal forms metal ions which are discharged at cathode forming pure metal.

Anode: $M \longrightarrow M^{n+} + ne^-$ Cathode: $M^{n+} + ne^- \longrightarrow M$

- **22.** (*i*) The positively charged colloidal particles of Fe(OH)₃ get coagulated by the oppositely charged Cl⁻ ions provided by KCl.
 - (ii) On passing direct current, colloidal particles move towards the oppositely charged electrode where they lose their charge and get coagulated.
 - (iii) Scattering of light by the colloidal particles takes place and the path of light becomes visible (Tyndall effect).

CBSE EXAMINATION PAPERS

ALL INDIA-2008

Time allowed: 3 hours Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (All India) SET-I

- 1. Define the 'forbidden zone' of an insulator.
- 2. Mention two ways by which lyophilic colloids can be coagulated.
- 3. Mention all the oxidation states exhibited by chlorine in its compounds.
- **4.** Why are lower members of aldehydes easily miscible with water?
- 5. Why do amines behave as nucleophiles?
- **6.** What are disaccharides? Give an example.
- 7. Define the term 'polymerisation'.
- **8.** What is understood by 'chemotherapy'?
- **9.** Define osmotic pressure. How is it that measurement of osmotic pressures is more widely used for determining molar masses of macromolecules than the rise in boiling point or fall in freezing point of their solutions?

OR

Derive an equation to express that relative lowering of vapour pressure for a solution is equal to the mole fraction of the solute in it when the solvent alone is volatile.

- 10. The conductivity of $0.20~\mathrm{M}$ solution of KCl at 298 K is $0.0248~\mathrm{S~cm}^{-1}$. Calculate its molar conductivity in this solution.
- 11. Assign reasons for the following:
 - (i) In liquid state, hydrogen chloride is a stronger acid than hydrogen fluoride.
 - (ii) Phosphorus (P_4) is much more reactive than nitrogen (N_2) .
- **12.** Discuss the relative stability in aqueous solutions of +2 oxidation state among the elements : Cr, Mn, Fe and Co. How would you justify this situation?
 - (At. Nos. Cr = 24, Mn = 25, Fe = 26, Co = 27)
- 13. What happens when bromine reacts with $CH_3 C \equiv CH$? How would you justify this reaction?
- 14. Write the IUPAC names of the following compounds:
 - (i) $(CH_3)_3 CCH_2 Br$

- 15. Alcohols react both as nucleophiles as well as electrophiles. Write one reaction of each type and describe its mechanism.
- **16.** How would you carry out the following conversions?
 - (i) Ethyl magnesium chloride to propan-1-ol
 - (ii) Benzyl chloride to benzyl alcohol
- 17. Write the structures of the monomers of the following polymers:
 - (i) PVC
 - (ii) Polypropene
- 18. What are biodegradable and non-biodegradable detergents? Give one example of each type.
- 19. Niobium (Nb) crystallises in a body-centred cubic (bcc) structure. If its density is 8.55 g cm⁻³, calculate the atomic radius of niobium.

(Atomic mass of Nb = 93 u;
$$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$$
)

Explain with suitable examples the following:

- (a) n-type and p-type semiconductors
- (b) F-centres
- (c) Ferromagnetism
- 20. Calculate the amount of KCl which must be added to 1 kg of water so that its freezing point is depressed by 2 K.

$$(K_f \text{ for water} = 1.86 \text{ K kg mol}^{-1}, \text{ Atomic mass K} = 39, \text{ Cl} = 35.5)$$

21. In the button cells widely used in watches and other devices the following reaction takes place:

$$Zn(s) + Ag_2O + H_2O(l) \longrightarrow Zn^{2+}(aq) + 2Ag(s) + 2OH^{-}(aq)$$

Determine $\Delta_r G^{\circ}$ for the reaction.

(Given:
$$E^{\circ}_{Zn^{2+}/Zn} = -0.76 \text{ V}$$
 and $E^{\circ}_{Ag^{+}/Ag} = +0.34 \text{ V}$)
22. Explain what is observed when

- - (i) a beam of light is passed through a colloidal solution.
 - (ii) an electrolyte NaCl, is added to hydrated ferric oxide sol.
 - (iii) an electric current is passed through a colloidal sol.
- 23. Write the chemical reactions which take place in the following operations:
 - (i) Electrolytic reduction of Al₂O₃.
 - (ii) Isolation of zinc from zinc blende.
 - (iii) Mond's process for refining nickel.
- **24.** Compare actinoids and lanthanoids with reference to their :
 - (i) electronic configurations of atoms
 - (ii) oxidation states of elements
 - (iii) general chemical reactivity of elements.
- 25. Write the IUPAC name and describe the magnetic behaviour (diamagnetic or paramagnetic) of the following coordination entities:
 - (i) $[Cr(H_2O)_2(C_2O_4)_2]^-$


- (*ii*) $[Co(NH_3)_5 Cl]^{2+}$
- (iii) $[NiCl_4]^{2-}$
- (At. Nos. : Cr = 24, Co = 27, Ni = 28)
- **26.** Account for the following:
 - (i) pK_h of methylamine is less than that of aniline.
 - (ii) Aniline does not undergo Friedel-Crafts reaction.
 - (iii) Ethylamine is freely soluble in water whereas aniline is only slightly soluble.
- 27. Define the following in relation to proteins:
 - (i) Primary structure
 - (ii) Denaturation
 - (iii) Peptide linkage
- **28.** (a) A reaction is of first order in A and of second order in B. Write the differential rate equation for this reaction.

How will its initial rate be affected if the concentration of both A and B are together doubled?

(b) The rate constant k of a reaction increases four fold when the temperature changes from 300 K to 320 K. Calculate the activation energy for the reaction. $(R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1})$

OR

- (a) List the factor which affect the rate of a chemical reaction.
- (b) The half-life for radioactive ¹⁴ C is 5730 years. The wooden part of an archaeological artefact has only 80% of the ¹⁴ C activity found in fresh wood. Calculate the age of the artefact.
- **29.** (a) Assign reasons for the following:
 - (i) Bi (V) is a stronger oxidising agent than Sb (V).
 - (ii) Of the noble gases only xenon is known to form established chemical compounds.
 - (b) Draw the structures of the following molecules:
 - (i) $H_2S_2O_7$ (ii) BrF₃ (iii) XeF₂
 - OR
 - (a) Complete the following chemical reaction equations :
 - (i) $Ca_3P_2 + H_2O \longrightarrow$
 - (ii) $XeF_4 + H_2O \longrightarrow$
 - (b) How would you account for the following observations:
 - (i) NH₃ is a stronger base than PH₃.
 - (ii) Sulphur in vapour state exhibits paramagnetism.
 - (iii) Hydrogen fluoride has a higher boiling point than hydrogen chloride.
- **30.** (a) Illustrate the following reactions giving one example for each :
 - (i) Cannizzaro reaction
 - (ii) Decarboxylation
 - (b) Complete the following reaction equations by giving the indicated missing substances:



$$(iii) \qquad \longrightarrow \text{CH}_2 \xrightarrow{\qquad \dots \qquad (?)} \longrightarrow \text{CHO}$$

OR

- (a) State tests to distinguish between the following pairs of compounds:
 - (i) Propanal and propanone
 - (ii) Phenol and benzoic acid
- (b) How will you bring about the following conversions:
 - (i) Propanone to propene
 - (ii) Benzaldehyde to benzophenone
 - (iii) Ethanol to 3-hydroxybutanal

CBSE (All India) SET-II

Questions Uncommon to Set-I

- 1. Which crystal defect lowers the density of a solid?
- **6.** What are reducing sugars? Give one example.
- **8.** What is tincture of iodine?
- 10. Define the following terms giving an example for each:
 - (i) The order of a reaction
 - (ii) The molecularity of a reaction
- **18.** Why do soaps not function in hard water, for washing clothes? How are synthetic detergents better than soaps for this purpose?
- 21. The rate constant for a first order reaction is $60 \, \text{s}^{-1}$. How much time will it take to reduce the initial concentration of the reactant to $1/16^{\, \text{th}}$ of its initial value?
- 23. Describe the principle involved in the following metallurgical operations:
 - (i) Zone refining
 - (ii) Electrolytic refining
 - (iii) Froth-floatation process of concentrating sulphide ores
- 27. What happens when D-glucose is treated with the following reagents?
 - (i) HNO₃
 - (ii) Bromine water
 - (iii) HI

Indicate the products formed.

28. (a) Depict the galvanic cell in which the following reaction takes place:

$$Zn(s) + 2Ag^{+}(aq) \longrightarrow Zn^{2+}(aq) + 2Ag(s)$$

Also indicate that in this cell

- (i) which electrode is negatively charged.
- (ii) what are the carrier of the current in the cell.
- (iii) what is the individual reaction at each electrode.
- (b) Write the Nernst equation and determine the e.m.f. of the following cell at 298 K:

$$Mg\left(s\right)\big|\,Mg^{\,2+}\left(0.001\,M\right)\big|\big|\,Cu^{\,2+}\left(0.0001\,M\right)\big|\,Cu\left(s\right)$$

(Given :
$$E^{\circ}_{\text{Mg}^{2+}/\text{Mg}} = -2.375 \text{ V}, E^{\circ}_{\text{Cu}^{2+}/\text{Cu}} = +0.34 \text{ V})$$
OR

- (a) Define conductivity and molar conductivity for the solution of an electrolyte. How do they vary when the concentration of electrolyte in the solution increases?
- (b) Three conductivity cells A, B and C containing solutions of zinc sulphate, silver nitrate and copper sulphate respectively are connected in series. A steady current of 1.5 amperes is passed through them until 1.45 g of silver is deposited at the cathode of cell B. How long did the current flow? What mass of copper and what mass of zinc got deposited in their respective cells?

(Atomic mass : Zn = 65.4 u, Ag = 108 u, Cu = 63.5 u)

CBSE (All India) SET-III

Questions Uncommon to Set-I and Set-II.

- 1. Name an element with which silicon may be doped to give a p-type semiconductor.
- 10. What is meant by a pseudo first order reaction? Give an example of a pseudo first order reaction and write the rate equation for the same.
- **12.** Assign a reason for each of the following:
 - (i) The third ionization energy of Mn (Z = 25) is higher than that of either Cr (Z = 24) or Fe (Z = 26).
 - (ii) Simple copper (I) salts are not stable in aqueous solutions.
- **18.** What are artificial sweetening agents? Give two examples.
- 21. The rate constant for a first order reaction is $60 \,\mathrm{s}^{-1}$. How much time will it take to reduce the initial concentration of reactant to 1/10th of its value?
- 24. Describe the trends in the following properties of the first series of the transition elements:
 - (i) Oxidation states
 - (ii) Atomic sizes
 - (iii) Magnetic behaviour of dipositive gaseous ions (M²⁺)

SOLUTIONS

CBSE (All India) SET-I

- The difference of energy between conduction band and valence band is called forbidden zone and for insulator its value is averaging between 3-6 eV.
- This can be done
 - (i) by adding an electrolyte.
 - (ii) by adding a suitable solvent.
- 3. Cl₂ exhibits -1, +1, +3, +5, +7 oxidation states in its compounds.
- 4. Lower members aldehydes are able to form intermolecular hydrogen bonds with water molecules. Hence, they are easily miscible with water.
- 5. Due to the presence of a lone pair of electrons on nitrogen atom.
- **6.** Carbohydrates that yield two monosaccharide units, on hydrolysis are called disaccharides, e.g.,
- 7. The process of formation of polymers from respective monomers is called polymerisation.

$$n\text{CH}_2 = \text{CH}_2 \xrightarrow[\text{monomer}]{\text{O}_2, 350\text{K} - 570 \text{ K}} \\ 1000 - 2000 \text{ atm} \xrightarrow{\text{Polymerisation}} -(-\text{CH}_2 - \text{CH}_2)_{\overline{n}} \\ \text{Polymerisation} & \text{Polyethene (LDP)} \\ \text{(polymer)}$$

- **Chemotherapy:** It is the branch of chemistry which deals with the treatment of diseases using suitable chemicals.
- 9. Osmotic pressure (π) may be defined as the extra pressure that must be applied to the solution to prevent the flow of solvent molecules into it through a semipermeable membrane.

$$\pi = \frac{n_B}{V}RT = CRT$$

$$\pi = \frac{W_B \cdot R \cdot T}{M_B \cdot V}$$

where V is the volume of solution in litre containing n_B moles of solute of molecular mass M_B . Thus knowing π , W_R , T and V molecular mass of the solute M_R can be calculated.

The osmotic pressure method has the advantage over rise in boiling point or fall in freezing point for determining molar masses of macromolecules because

- (i) Osmotic pressure is measured at the room temperature and the molarity of solution is used instead of molality.
- (ii) Compared to other colligative properties, its magnitude is large even for very dilute solutions.

For a solution of volatile liquids Raoult's law, is given as

$$P = P_A + P_B$$

If solute (component *B*) is non-volatile then

$$P = P_A = P_A^0 x_A$$

$$P = P_A^0 (1 - x_B)$$

$$P = P_A^0 - P_A^0 x_B$$

$$P_A^0 x_B = P_A^0 - P$$

$$(\because x_A + x_B = 1)$$

$$\frac{P_A^{\, \circ} - P}{P_A^{\, \circ}} = x_B$$

Thus, relative lowering of vapour pressure is equal to the mole fraction of non-volatile solute.

10.
$$K = 0.0248 \text{ S cm}^{-1}$$

$$M = 0.2 \text{ mol L}^{-1}$$

 $\lambda_m = \frac{K \times 1000}{M} = \frac{0.0248 \times 1000}{0.2}$
 $\lambda_m = 124 \text{ S cm}^2 \text{ mol}^{-1}$

- **11.** (*i*) It is due to
 - (a) Intermolecular hydrogen bonding in HF.
 - (b) Higher H—F bond dissociation enthalpy than H—Cl.
 - (ii) As P—P single bond (213 kJ mol⁻¹) in P_4 is much weaker than $N \equiv N$ triple bond $(941.4 \text{ kJ mol}^{-1}) \text{ in N}_2$.
- 12. On the basis of electrochemical series the standard electrode potential shows the following order

$$E^{\circ}_{\text{Mn}^{2+}/\text{Mn}} < E^{\circ}_{\text{Cr}^{2+}/\text{Cr}} < E^{\circ}_{\text{Fe}^{2+}/\text{Fe}} < E^{\circ}_{\text{Co}^{2+}/\text{Co}}$$

Therefore, Co²⁺ gets easily reduced to metallic cobalt while it is difficult to reduce Mn²⁺. Hence Mn^{2+} will be most stable and the increasing stability order will be $Co^{2+} < Fe^{2+} < Cr^{2+} < Mn^{2+}$

$$\text{Co}^{2+} < \text{Fe}^{2+} < \text{Cr}^{2+} < \text{Mn}^{2+}$$

13. When bromine reacts with propyne, the reddish brown colour of bromine is discharged as long as propyne is present in excess.

$$CH_{3} - C \equiv CH + Br_{2} \xrightarrow{CCl_{4}} CH_{3} - C = C - H \xrightarrow{+Br_{2}} CH_{3} - C - C - H$$

$$Br \xrightarrow{Br} Br$$

$$CH_{3} - C = C - H \xrightarrow{+Br_{2}} CH_{3} - C - C - H$$

$$Br \xrightarrow{Br} Br$$

$$1, 1, 2, 2, \text{ Tetra Bromo propane}$$

This is due to the formation of 1, 1, 2, 2-tetrabromo propane which is colourless.


14. (i)
$$CH_3$$
 CH_2 CH_2 CH_2 CH_3 CH_3

1-Bromo-2, 2-dimethyl propane

15. Alcohols as nucleophile:

The bond between O—H is broken when an alcohol reacts as a nucleophile.

Mechanism:



Alcohols as electrophile:

The bond between C—O is broken when alcohols react as electrophiles.

$$\mathbf{R}\mathbf{--}\mathbf{C}\mathbf{H}_{2}\mathbf{--}\mathbf{O}\mathbf{--}\mathbf{H} \ + \ \mathbf{H}^{+} \ \longrightarrow \ \mathbf{R}\mathbf{--}\mathbf{C}\mathbf{H}_{2}\mathbf{--}\mathbf{O}^{+}\mathbf{H}_{2}$$

- 17. Polymer Monomer Structure of the monomer
 - (i) PVC Vinyl chloride $CH_2 = CH$ —Cl (ii) Polypropene Propene CH_3 — $CH = CH_2$
- **18. Biodegradable detergents:** Detergents having straight hydrocarbon chains are easily degraded by micro-organism and hence called biodegradable detergents, *e.g.*, sodium–4–(1-dodecyl) benzene sulphonate.

$$CH_3$$
— $(CH_2)_{11}$ — SO_3^- Na

Non-biodegradable detergents: Detergents having branched hydrocarbon chains are not easily degraded by the micro-organism and hence are called non-biodegradable detergents, *e.g.*, sodium-4-(1, 3, 5, 7-**tetramethyl octyl**) benzenesulphonate.

$$\begin{array}{c|c} \operatorname{CH_3} & \operatorname{CH_3} \\ | & | \\ \operatorname{CH_3} & \leftarrow \operatorname{CH--CH_2} \rightarrow_3 - \operatorname{CH---} \\ \end{array} \begin{array}{c} \operatorname{CH_3} \\ | & | \\ \operatorname{SO_3^-} \\ \operatorname{Na} \end{array}$$

Non-biodegradable detergents accumulate in rivers and waterways thereby causing water pollution.

19. $d = 8.55 \text{ g/cm}^3$, M = 93 g/mol

For bcc,
$$z = 2$$
, $a = ?$ $N_A = 6.02 \times 10^{23}$ $d = \frac{z \times M}{a^3 \times N_A}$

Substituting the values

8.55 =
$$\frac{2 \times 93}{a^3 \times 6.02 \times 10^{23}}$$

 $a^3 = \frac{2 \times 93}{8.55 \times 6.02 \times 10^{23}}$
 $a = \left(\frac{930}{8.55 \times 3.01}\right)^{1/3} 10^{-8}$
Let $x = \left(\frac{930}{8.55 \times 3.01}\right)^{1/3}$
 $\therefore \log x = \log\left(\frac{930}{8.55 \times 3.01}\right)^{1/3}$
 $= \frac{1}{3}(\log 930 - \log 8.55 - \log 3.01)$
 $= \frac{1}{3}(2.9685 - 0.9320 - 0.4786)$
 $\log x = \frac{1}{3}(1.5579) = 0.5193$
 $x = \text{Antilog } (0.5193)$
 $x = 3.306$
 $\therefore a = 3.306 \times 10^{-8} \text{ cm}$

Now
$$r = \frac{\sqrt{3}}{4} a$$

$$\therefore \qquad r = \frac{\sqrt{3}}{4} \times 3.306 \times 10^{-8} = \frac{1.732 \times 3.306 \times 10^{-8}}{4}$$

$$r = 1.4315 \times 10^{-8} \text{ cm} = 143.15 \times 10^{-10} \text{ cm}$$

$$r = 143.15 \text{ pm}$$

(a) n-type semiconductor: When a silicon or germanium crystal is doped with group 15 element like P or As, the dopant atom forms four covalent bonds like a Si or Ge atom but the fifth electron, not used in bonding, becomes delocalised and contribute its share towards electrical conduction. Thus silicon or germanium doped with P or As is called n-type semiconductor, n indicative of negative, since it is the electron that conducts electricity.

p-type semiconductor: When silicon or germanium is doped with group 13 element like B or Al, the dopant has only with three, valence electrons. An electron vacancy or a hole is created at the place of the missing fourth electron. Here, this hole moves through the crystal like a positive charge giving rise to electrical conductivity. Thus Si or Ge doped with B or Al is called p type of semiconductor, (P stands for positive hole) since it is the positive hole that is responsible for conduction.

- (b) **F-centres:** Electrons trapped in anion vacancies are called F-centres. They impart characteristic colour to the compound and increase electrical conductivity.
- (c) Ferromagnetism: Ferromagnetic substances are those substances which are strongly attracted by external magnetic field, e.g., (iron, cobalt, nickle and CrO₂ etc.) Ferromagnetism arises due to spontaneous alignment of magnetic moments in the same direction.



Alignment of magnetic moments in ferromagnetic substance.

20. Assuming 100% dissociation of KCl, i = 2, $\Delta T_f = 2$ K, $K_f = 1.86$ K kg mol⁻¹, $W_R = ?$

$$\begin{split} M_B = 74.5 \text{ g mol}^{-1}, W_A = &1000 \text{ g} \\ \Delta T_f = & \frac{i \times K_f \times W_B \times 1000}{M_B \times W_A} \end{split}$$

Substituting the values
$$2 = \frac{2 \times 1.86 \times W_B \times 1000}{74.5 \times 1000}$$

$$W_B = \frac{74.5}{1.86} = 40.05 \text{ g}$$

21.
$$Zn \longrightarrow Zn^{2+} + 2e^{-}$$

$$Ag_2O + H_2O + 2e^{-} \longrightarrow 2Ag^{+} + 2OH^{-}$$

$$Zn + Ag_2O + H_2O \longrightarrow Zn^{2+} + 2Ag + 2OH^{-}$$

$$E^{\circ}_{cell} = E^{\circ}_{cathode} - E^{\circ}_{anode}$$

$$= E^{\circ}_{Ag^{+}/Ag} - E^{\circ}_{Zn^{2+}/Zn}$$

$$E^{\circ}_{cell} = 0.34 - (-0.76)$$

$$= 1.10 \text{ V}$$

also
$$n = 2$$

$$\Delta G^{\circ} = -nFE^{\circ}_{\text{cell}}$$

$$\Delta G^{\circ} = -2 \times 96500 \times 1.10$$

$$\Delta G^{\circ} = 2.123 \times 10^{5} \text{ J}$$

- 22. (i) Scattering of light by the colloidal particles takes place and the path of light becomes visible (Tyndall effect).
 - (ii) The positively charged colloidal particles of Fe(OH) 3 get coagulated by the oppositely charged Cl ions provided by NaCl.
 - (iii) On passing direct current, colloidal particles move towards the oppositely charged electrode where they lose their charge and get coagulated.
- 23. (i) Electrolytic reduction of Al_2O_3 :

Cathode: Al³⁺ (melt) + 3e⁻
$$\longrightarrow$$
 Al
Anode: C(s) + O²⁻ (melt) \longrightarrow CO(g) + 2e⁻
C(s) + 2O²⁻ (melt) \longrightarrow CO₂(g) + 4e⁻

(ii) Isolation of zinc from zinc blende:

Roasting:
$$2 \operatorname{ZnS} + 3\operatorname{O}_2 \xrightarrow{\Delta} 2 \operatorname{ZnO} + 2 \operatorname{SO}_2$$
Zinc blende

Reduction:
$$ZnO + C \xrightarrow{1673 \text{ K}} Zn + CO$$

(iii) Mond's process for refining nickel:

Ni + 4CO
$$\xrightarrow{330-350 \text{ K}}$$
 Ni(CO)₄
Ni(CO)₄ $\xrightarrow{450-470 \text{ K}}$ Ni + 4CO

24.

	Characteristics	Lanthanoids	Actinoids
(i)	Electronic configuration	[Xe] $4f^{1-14}5d^{0-1}6s^2$	[Rn] $5f^{1-14} 6d^{0-1}7s^2$
(ii)	Oxidation states	Besides + 3 O.S. lanthanoids show +2 and +3 O.S. only in a few cases.	
(iii)	General chemical reactivity of elements	These are less reactive metals	These are highly reactive metals.
		Lesser tendency towards complex formation.	Greater tendency towards complex formation.
		Do not form oxocation	Form oxocation
	Compounds are less basic. Compounds are		Compounds are more basic.

25. (*i*) $[Cr(H_2O)_2 (C_2O_4)_2]^-$

Diaquadioxalatochromate (III) ion

O.S. of
$$Cr = x + 0.2 + (-2).2 = -1$$
, $x = +3$
Electronic configuration of $Cr^{3+} = 3d^3 + 4s^0 = t_{2g}^3 = e_g^0$

Unpaired electrons (n) = 3, Paramagnetic.

 $\left[\operatorname{Co}(\operatorname{NH}_3)_5\operatorname{Cl}\right]^{2+}$ (ii)

Pentaamminechloro cobalt (III) ion

O.S. of Co =
$$x + 0.5 + (-1) \cdot 1 = 2$$
, $x = 3$

Electronic configuration of $\operatorname{Co}^{3+} = 3d^6 4s^0 = t_{2g}^6 e_g^0$

Unpaired electrons (n) = 0, Diamagnetic.

 $[NiCl_4]^{2-}$ (iii)


Tetrachloronickelate (II) ion

O.S. of Ni =
$$x + (-1) 4 = -2$$
, $x = +2$

O.S. of Ni = x + (-1) 4 = -2, x = +2Electronic configuration of Ni²⁺ = $3d^8 4s^0 = t_{2g}^8 e_g^0$

Unpaired electrons (n) = 2, Paramagnetic.

- In aniline, due to resonance, the lone pair of electrons on the nitrogen atom are delocalized over the benzene ring. As a result, the electron density on the nitrogen decreases. On the other hand, in methyl amine +ve I effect of CH₃ increases the electrondensity on the nitrogen atom. Therefore aniline is a weaker base than methyl amine and hence its pK_b value is higher than that of methyl amine.
 - (ii) Aniline being a Lewis base reacts with lewis acid AlCl₃ to form a salt. Due to this, N atom of aniline acquires positive charge and hence acts as a strong deactivation group for further reaction.
 - (iii) Ethyl amine is freely soluble in water because it forms hydrogen bonds with water molecules.



On the other hand in aniline due to large, hydrocarbon part, the extent of hydrogen bonding decreases considerably and hence aniline is slightly soluble.

- **Primary structure of proteins:** The sequence in which various amino acids are arranged in a protein is called its primary structure. Any change in the sequence of amino acids creates different protein which alters biological functions.
 - (ii) Denaturation: When a protein in its native form is subjected to physical change like change in temperature or chemical change like change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and proteins lose its biological activity. During denaturation 2° and 3° structures of proteins are destroyed but 1° structure remains intact, e.g., coagulation of egg white on boiling.
 - (iii) Peptide linkage: A peptide linkage is an amide linkage (—CONH—) formed between —COOH group of one α -amino acid and NH₂ group of the other amino acid by the elimination of a water molecule.

$$\begin{array}{c} O \\ \parallel \\ H_2N-CH_2-C-OH+HNH-CH-COOH \xrightarrow{-H_2O} \\ CH_3 \\ H_2NCH_2-C-NH-CH-COOH \\ CH_3 \\ \end{array}$$

28. (a) Rate =
$$\frac{dx}{dt} = K[A][B]^2$$

If concentration of both A and B are doubled, then

Rate =
$$K[2A][2B]^2$$

= $8K[A][B]^2$

i.e., the rate of reaction becomes 8 times.

(b)
$$K_2 = 4K_1$$
 i.e., $\frac{K_2}{K_1} = 4$

$$T_1 = 300 \text{ K}$$

$$T_2 = 320 \text{ K}$$

$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303 R} \left(\frac{T_2 - T_1}{T_1 T_2}\right)$$

$$\log 4 = \frac{E_a}{2.303 \times 8.314} \left(\frac{320 - 300}{300 \times 320}\right)$$

$$2 \log 2 = \frac{E_a}{19.147} \left(\frac{20}{300 \times 320}\right)$$

$$E_a = \frac{2 \times 0.3010 \times 19.147 \times 300 \times 320}{20} = 55327 \text{ J}$$

$$E_a = 55.327 \text{ kJ}$$
OR

(a) Rate of reaction depends on

- (i) Concentration
- (ii) Temperature
- (iii) Nature of reactant
- (iv) Pressure of the gaseous reactant
- (v) Surface area
- (vi) Catalyst

(b)
$$t_{1/2} = 5730 \text{ years}$$

$$K = \frac{0.693}{t_{1/2}} = \frac{0.693}{5730} = 1.209 \times 10^{-4} \text{ year}^{-1}$$
$$t = \frac{2.303}{K} \log \frac{[R]_0}{[R]} = \frac{2.303}{1.2 \times 10^{-4}} \log \frac{100}{80}$$

$$t = \frac{2.303 \times 10^4}{1.2} (\log 10 - \log 8) = \frac{2.303 \times 10^4}{1.2} (1 - 3 \log 2)$$
$$t = \frac{2.303}{1.2} \times 10^4 (1 - 3 \times 0.3010) = \frac{2.303 \times 0.097 \times 10^4}{1.209}$$
$$t = 1847.7 \text{ years}$$


- **29.** (a) (i) Due to stronger inert pair effect Bi (V) gets readily reduced to Bi (III) therefore, Bi (V) is a stronger oxidising agent than Sb (V).
 - (ii) Xe has least ionization enthalpy among noble gases and hence it readily forms chemical compounds particularly with O_2 and F_2 .

(ii) BrF₃

$$sp^3d$$

hybridisation
$$5e$$


$$1. p. =$$



(iii)
$$XeF_2$$

 sp^3d
hybridisation
$$5e$$

$$OR$$



(a) (i)
$$\operatorname{Ca}_{3}\operatorname{P}_{2} + 6\operatorname{H}_{2}\operatorname{O} \longrightarrow \operatorname{PH}_{3} + 3\operatorname{Ca}(\operatorname{OH})_{2}$$

Phosphine

- (ii) $6XeF_4 + 12H_2O \longrightarrow 2XeO_3 + 24HF + 4Xe + 3O_2$
- (b) (i) As the atomic size of nitrogen is smaller than phosphorus, electron density on nitrogen atom is higher than that on phosphorus atom. Consequently, the tendency of N in NH₃ to denote its lone pair of electrons is much higher than that of P in PH₃. Thus NH₃ is a stronger base than PH₃.
 - (ii) In vapour state sulphur partly exists as S_2 molecule which has two unpaired electrons in the antibonding π^* orbital like O_2 and hence exhibits paramagnetic behaviour.
 - (iii) Because of hydrogen bonding HF exists as associated molecules in a liquid and therefore has high boiling point. Due to large size and low electronegativity of chlorine no hydrogen bonding is present in HCl, only Vander Waal forces are present. The boiling of HCl is therefore low.
- **30.** (a) (i) Cannizzaro reaction:

$$R - COOH + NaOH \longrightarrow R - COONa + H_2O$$

$$R - COONa \xrightarrow{NaOH/CaO} \xrightarrow{A} \xrightarrow{R-H} + Na_2CO_3$$

$$O \qquad O$$

$$CH_3 \qquad \parallel \qquad C = O + H_2NNH - C - NH_2 \longrightarrow H$$

$$Ethanal \qquad Ethanal$$

$$(ii) \qquad O_3 \qquad 2 \longrightarrow O$$

$$CH_2 \xrightarrow{(ii) H_2O_2/\overline{O}H} \longrightarrow CHO$$

$$OR$$

(a) (i) Propanone on treatment with I₂ / NaOH (NaOI) undergoes iodoform reaction to give yellow ppt of iodoform but propanal does not.

Tollen's test: Propanal being an aldehyde reduces Tollen's reagent to silver mirror but propanone being a ketone does not.

(ii) FeCl₃ test: Phenol gives a violet colouration with neutral FeCl₃ solution while benzoic acid gives buff coloured ppt.

$$6C_6H_5OH + FeCl_3 \longrightarrow [Fe(OC_6H_5)_6]^{-3} + 3H^+ + 3HCl$$
Violet complex

NaHCO₃ test: Benzoic acid being a stronger acid than phenol decomposes NaHCO₃ to evolve CO₂ but phenol does not.

(ii)
$$\underbrace{\operatorname{KMnO_4/OH}}_{\text{Benzoyl chloride}}$$
 $\underbrace{\operatorname{COCl}}_{\text{anhyd. AlCl}_3}$ $\underbrace{\operatorname{Benzophenone}}_{\text{Benzophenone}}$

$$(iii) \ \text{CH}_3 - \text{CH}_2 - \text{OH} \xrightarrow{\text{PCC}} \begin{array}{c} \text{OH} \\ \\ | \\ \text{Ethanol} \end{array} \xrightarrow{\text{PCC}} \begin{array}{c} \text{NaOH (dil.)} \\ \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{CHO} \\ \text{Acetaldehyde} \end{array} \xrightarrow{\text{NaOH (dil.)}} \begin{array}{c} \text{OH} \\ \\ \text{S-Hydroxybutanal} \end{array}$$

CBSE (All India) SET-II

- 1. Schottky defect.
- **6.** All carbohydrates which reduce Tollen's reagent and Fehling's solution are referred to as reducing sugars, *e.g.*, glucose.
- **8.** A 2-3 per cent solution of iodine in alcohol water mixture is known as tincture of iodine. It is used as an antiseptic.
- **10.** (*i*) Order of reaction may be defined as the sum of powers of the concentration of the reactants in the rate law expression.

For example consider the reaction

$$NH_4NO_2 \longrightarrow N_2 + 2H_2O$$

Experimentally, it is observed that the rate law for this reaction is

Rate =
$$K$$
 [NH₄NO₂]

Hence, the order of reaction is 1.

(ii) Molecularity of a reaction may be defined as the number of reacting species (atoms, ions or molecules) taking part in an elementary reaction which must collide simultaneously in order to bring about a chemical reaction.

For example molecularity of the reaction

2HI
$$\longrightarrow$$
 H₂ + I₂ is 2 as it involves.

Simultaneous collision between two HI molecules.

18. Hard water contains calcium and magnesium ions. Therefore in hard water soap get precipitated as calcium and magnesium soap which being insoluble stick to the cloth as gummy mass. Hence soap cannot be used with hard water.

$$2C_{17}H_{35}COO^-Na^+ + CaCl_2 \longrightarrow 2NaCl + (C_{17}H_{35}COO)_2Ca$$

Soap insoluble calcium stearate (Soap)

On the other hand calcium and magnesium salts of detergents are soluble in water so they easily form lather with hard water.

21.
$$K = 60 \text{ s}^{-1}$$
, $[R] = \frac{[R]_0}{16}$
 $t = ?$

$$t = \frac{2.303}{K} \log \frac{[R]_0}{[R]}$$

$$t = \frac{2.303}{60} \log \frac{[R]_0}{\frac{[R]_0}{16}}$$

$$t = \frac{2.303}{60} \log 16 = \frac{2.303}{60} \cdot 4 \log 2$$

$$t = \frac{2.303}{15} \times 0.3010 = \frac{0.6932}{15}$$


$$t = 4.62 \times 10^{-2} \text{ s.}$$

- **23.** (*i*) Zone refining is based on the principle that the impurities are more soluble in the liquid state than in the solid state of metal.
 - (ii) In electrolytic refining the impure metal is made to act as anode. A strip of the same metal in pure form is used as cathode. They are put in a suitable electrolytic bath containing soluble salt of the same metal. When electric current is passed, impure metal forms metal ions which are discharged at cathode forming pure metal.

Anode:
$$M \longrightarrow M^{n+} + ne^{-}$$

Cathode: $M^{n+} + ne^{-} \longrightarrow M$

(iii) The principle of refining by froth floatation process is that sulphide particles are preferentially wetted by pine oil, whereas the gangue particles are wetted by water.



- (i) Zn electrode is negatively charged.
 - (ii) Current carriers of cell are
 - electrons in external wire.
 - Zn²⁺ ions in anodic half cell.

- Ag + ions in cathodic half cell.
- Ions of salt bridge, *i.e.*, K⁺ and Cl⁻.

(iii) At anode
$$\operatorname{Zn} \longrightarrow \operatorname{Zn}^{2+} + 2e^{-}$$
At cathode $2\operatorname{Ag}^{+} + e^{-} \longrightarrow 2\operatorname{Ag}$
(b) $\operatorname{Mg} \longrightarrow \operatorname{Mg}^{2+} + 2e^{-}$

$$\operatorname{Cu}^{2+} + 2e^{-} \longrightarrow \operatorname{Cu}$$

$$\operatorname{Mg} + \operatorname{Cu}^{2+} \longrightarrow \operatorname{Cu} + \operatorname{Mg}^{2+}$$

According to Nernst equation,

$$\begin{split} E_{\text{cell}} &= E^{\circ}_{\text{ cell}} - \frac{0.059}{n} \log \frac{[\text{Cu}][\text{Mg}^{2+}]}{[\text{Mg}][\text{Cu}^{2+}]} \\ E_{\text{cell}} &= (E^{\circ}_{\text{Cu}^{2+}/\text{Cu}} - E^{\circ}_{\text{Mg}^{2+}/\text{Mg}}) - \frac{0.059}{2} \log \frac{[\text{Mg}^{2+}]}{[\text{Cu}^{2+}]} \\ &= 0.34 - (-2.375) - \frac{0.059}{2} \log \frac{10^{-3}}{10^{-4}} \\ &= 0.34 + 2.375 - 0.0295 \log 10 \\ E_{\text{cell}} &= 2.6855 \text{ V} \\ E_{\text{cell}} &= 2.685 \text{ V} \\ \mathbf{OR} \end{split}$$

(a) The conductivity of a solution at any given concentration is the conductance of one unit volume of solution kept between two platinum electrodes with unit area of cross section at a distance of unit length.

On increasing the concentration of solution, the number of ions per unit volume of solution increases and thus its conductivity increases.

Molar conductivity (Λ_m) of a solution at a given concentration is the conductance of the volume V of solution containing one mole of electrolyte kept between two electrodes with area of cross section A and distance of unit length. Therefore

$$\Lambda_m = \frac{\kappa A}{l} = \kappa$$

Since l = 1 and A = V (volume containing 1 gram mole of electrolyte)

$$\Lambda_m = \kappa V$$

Molar conductivity increases with decrease in concentration. This is because the total volume, *V* of solution containing one mole of electrolyte also increases. It has been found that decrease in *K* on dilution of solution is more than compensated by increase in its volume.

$$(b) Ag^+ + e^- \longrightarrow Ag$$

108 g of Ag are deposited by 96500 C

∴ 1.45 g of Ag will be deposited by =
$$\frac{96500}{108} \times 1.45 \text{ C}$$

= 1295.6 C
 $t = \frac{Q}{I} = \frac{1295.6}{150} = 863.7 \text{ s}.$

$$Cu^{2+} + 2e^{-} \longrightarrow Cu$$
2 × 96500 C deposit Cu = 63.5 g
∴ 1295.6 C deposit Cu = $\frac{63.5}{2 \times 96500} \times 1295.6 = 0.426$ g
$$Zn^{2+} + 2e^{-} \longrightarrow Zn$$
2 × 96500 C deposit Zn = $\frac{65.3}{2 \times 96500} \times 1295.6 = 0.438$ g
∴ 1295.6 C deposit Zn = $\frac{65.3}{2 \times 96500} \times 1295.6 = 0.438$ g

CBSE (All India) SET-III

- **1.** Boron or Aluminium.
- **10.** A reaction which is of higher order but follows the kinetics of first order under special conditions is called a pseudo first order reaction.

Example, Acid hydrolysis of ethyl acetate.

$$CH_3$$
— $COOC_2H_5$ + H_2O $\xrightarrow{H^+}$ CH_3 — $COOH$ + C_2H_5 — OH

Here, the rate law is given by expression

Rate =
$$K$$
 [CH₃—COOC₂H₅]

The concentration of H_2O is so large that it hardly undergoes any change during the reaction, therefore, it does not appear in the rate law.

- 12. (i) This is because Mn²⁺ is more stable as it has exactly half filled configuration $3d^5 4s^0$.
 - (ii) Cu²⁺(aq) is much more stable than Cu⁺(aq). This is because, although second ionization enthalpy of copper is large but for Cu²⁺(aq) is much more negative than that of Cu⁺(aq) and therefore, it more than compensates for the second ionisation enthalpy of copper. Therefore, Cu⁺ ion aqueous solution undergoes disproportionation.

$$2Cu^{+}(aq) \longrightarrow Cu^{2+}(aq) + Cu(s)$$

18. Artificial sweetening agents: These are the substances which are non-nutritive in nature and used as substitutes for sugar in foods and beverages.

Examples:

Aspartame: Aspartame is 100 times as sweet as cane sugar. Its use is limited to cold foods and soft drinks because it is unstable at cooking temperature.

Saccharin: It is about 550 times as sweet as cane sugar. Its use is of great value to diabetic persons and people who need to control intake of calories.

21.
$$K = 60 \text{ s}^{-1}$$

$$t = \frac{2.303}{K} \log \frac{[R]_0}{[R]}$$

$$t = \frac{2.303}{60} \log \frac{[R]_0}{\frac{[R]_0}{10}} = \frac{2.303}{60} \log 10 = 0.038 \times 1$$

$$t = 0.038 \text{ s}$$

24. (*i*) As there is very little energy difference between 4*s* and 3*d* orbitals, electrons from both energy levels can be used for chemical bond formation. Therefore all elements except Sc and Zn, of the first transition series show a number of oxidation states as shown in table.

Oxidation states of the first series transition elements (the most common ones are in bold letter)

Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
	+2	+2	+2	+2	+2	+2	+2	+1	
+3	+3	+3	+3	+3	+3	+3	+3	+2	+2
	+4	+4	+4	+4	+4	+4	+4		
		+5	+5	+5					
			+6	+6	+6				
				+7					

- (ii) Atomic radii of the first transition series decreases from Sc to Cr, then remains almost constant till Ni and then increases from Cu to Zn.
 - The reason of this variation in atomic radii has been attributed to the increase in nuclear charge in the beginning of the series. But as the electrons continue to be filled in d-orbitals, they screen the outer 4s electrons from the influence of nuclear charge. When the increased nuclear charge and the increased screening effect balance each other in the middle of transition series, the atomic radii becomes almost constant (Mn to Fe). Towards the end of the series, the repulsive interaction between electrons in d orbitals become very dominant. As a result there is an expansion of the electron cloud; consequently, the atomic size increases.
- (iii) Except Zn^{2+} , all other divalent gaseous ions of the first series of the transition elements contain unpaired electrons in their 3d subshell and are therefore paramagnetic in nature.

The magnetic moment (μ) of the elements of the first transition series can be calculated with the unpaired electrons (n) by the spin-only formula

$$\mu = \sqrt{n(n+2)} \text{ B. M.}$$

Ion	Configuration	Unpaired electrons	Magnetic moment (μ) calculated
Mn ²⁺	$3d^5 4s^0$	5	$\sqrt{5(5+2)} = 5.92 \text{ B.M.}$
Cu ²⁺	$3d^{9}4s^{0}$	1	$\sqrt{1(1+2)} = 1.73 \text{ B.M.}$
Zn ²⁺	$3d^{10}4s^0$	0	$\sqrt{0(0+3)}=0$

CBSE EXAMINATION PAPERS

DELHI-2009

Time allowed: 3 hours Maximum marks: 70

General Instructions:


- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Delhi) SET-I

1.	Which point defect in crystals does not alter the density of the relevant solid?	1
2.	Define the term 'Tyndall effect'.	1
3.	Why is the froth floatation method selected for the concentration of sulphide ores?	1
4.	Why is Bi (V) a stronger oxidant than Sb (V)?	1
5.	Give the IUPAC name of the following compound: $ \begin{array}{c c} CH_3-C=C-CH_2OH \\ & CH_3 \end{array} $ CH ₃ Br	1
6.	Write the structure of 3-oxopentanal.	1
7.	Why is an alkylamine more basic than ammonia?	1
8.	Give an example of elastomers.	1
9.	A reaction is of second order with respect to a reactant. How will the rate of reaction be affected if t concentration of this reactant is	he 2
	(i) doubled (ii) reduced to half?	
10.	Explain the role of	2
	(i) Cryolite in the electrolytic reduction of alumina.	
	(ii) Carbon monooxide in the purification of nickel.	
11.	Draw the structures of the following molecules:	2
	(i) XeF ₄ (ii) BrF ₃	
12.	Complete the following chemical reaction equations:	2
	(i) $P_{4(s)}$ NaOH _(aq) $H_2O_{(l)}$	
	(ii) I (aq) $H_2O_{(l)}$ $O_{3(g)}$	

44 | Xam idea Chemistry-XII

- 13. Differentiate between molality and molarity of a solution. What is the effect of change in temperature of a solution on its molality and molarity?
- 14. Which ones in the following pairs of substances undergoes S_N2 substitution reaction faster and why? 2



- (ii) I or C
- **15.** Complete the following reaction equations:

(i)
$$-OH + SOCl_2$$
 $-CH_2OH$ $+ HCl$ $-$

- **16.** Explain what is meant by
 - (i) a peptide linkage
- (ii) a glycosidic linkage

2

2

2

- 17. Name two water soluble vitamins, their sources and the diseases caused due to their deficiency in diet.
- **18.** Draw the structures of the monomers of the following polymers:
 - (i) Teflon

(ii) Polythene

OR

What is the repeating unit in the condensation polymer obtained by combining HO₂CCH₂CH₂CO₂H (succinic acid) and H₂NCH₂CH₂NH₂ (ethylene diamine).

- 19. Iron has a body-centred cubic unit cell with a cell edge of 286.65 pm. The density of iron is 7.87g cm⁻³.
 Use this information to calculate Avogadro's number (At. mass of Fe = 56g mol⁻¹).
 3
- 20. 100 mg of a protein is dissolved in just enough water to make 10.0 mL of solution. If this solution has an osmotic pressure of 13.3 mm Hg at 25°C, what is the molar mass of the protein?
 3 (R = 0.0821 L atm mol⁻¹ K⁻¹ and 760 mm Hg = 1 atm.)
- 21. A first order reaction has a rate constant of 0.0051 min⁻¹. If we begin with 0.10M concentration of the reactant, what concentration of reactant will remain in solution after 3 hours?
- 22. How are the following colloids different from each other in respect of dispersion medium and dispersed phase? Give one example of each type.
 - (i) An aerosol
 - (ii) A hydrosol
 - (iii) An emulsion

23. Account for the following: 3 (i) NH₃ is a stronger base than PH₃. (ii) Sulphur has a greater tendency for catenation than oxygen. (iii) Bond dissociation energy of F2 is less than that of Cl2. OR Explain the following situations: (i) In the structure of HNO₃ molecule, the N—O bond (121 pm) is shorter than N—OH bond (140 pm). (ii) SF₄ is easily hydrolysed whereas SF₆ is not easily hydrolysed. (iii) XeF₂ has a straight linear structure and not a bent angular structure. **24.** For the complex $[Fe(en)_2Cl_2]Cl$, (en = ethylene diamine), identify 3 (i) the oxidation number of iron. (ii) the hybrid orbitals and the shape of the complex. (iii) the magnetic behaviour of the complex. (iv) the number of geometrical isomers. (v) whether there is an optical isomer also, and (vi) name of the complex. (At. no. of Fe = 26) **25.** Explain the mechanism of the following reactions: 3 (i) Addition of Grignard's reagent to the carbonyl group of a compound forming an adduct followed by hydrolysis. (ii) Acid catalysed dehydration of an alcohol forming an alkene. (iii) Acid catalysed hydration of an alkene forming an alcohol. **26.** Giving an example for each, describe the following reactions: 3 (i) Hofmann's bromamide reaction (ii) Gatterman reaction (iii) A coupling reaction **27.** Explain the following types of substance with one suitable example, for each case: 3 (i) Cationic detergents (ii) Food preservatives (iii) Analgesics 28. (a) Define molar conductivity of a substance and describe how for weak and strong electrolytes, molar conductivity changes with concentration of solute. How is such change explained? (b) A voltaic cell is set up at 25 °C with the following half cells: Ag⁺ (0.001 M) | Ag and Cu²⁺ (0.10 M) | Cu What would be the voltage of this cell? $(E_{cell}^{o} = 0.46 \text{ V})$

OR

- (a) State the relationship amongst cell constant of a cell, resistance of the solution in the cell and conductivity of the solution. How is molar conductivity of a solute related to conductivity of its solution?
- (b) A voltaic cell is set up at 25°C with the following half-cells:

Calculate the cell voltage $[E^o_{Ni}^2 + _{/Ni} = -0.25 \text{ V}, E^o_{Al/Al}^3 + = -1.66 \text{ V}]$

- **29.** (a) Complete the following chemical reaction equations:
 - (i) $MnO_{4(aq)}^{-}$ $C_2O_{4(aq)}^{2-}$ H (aq)
 - (ii) $\operatorname{Cr}_2\operatorname{O}_{7(aq)}^{2-}$ F^2 (aq) H (aq)
 - (b) Explain the following observations about the transition/inner transition elements:
 - (i) There is in general an increase in density of element from titanium (Z = 22) to copper (Z = 29).

5

5

- (ii) There occurs much more frequent metal-metal bonding in compounds of heavy transition elements (3rd series).
- (iii) The members in the actinoid series exhibit a large number of oxidation states than the corresponding members in the lanthanoid series.

OR

- (a) Complete the following chemical equations for reactions:
 - (i) MnO_4^- _(aq) + $\text{S}_2\text{O}_3^{2-}$ _(aq) + $\text{H}_2\text{O}_{(l)}$
 - (ii) $\text{CrO}_{7(aq)}^{-} + \text{H}_{2}\text{S}_{(g)} + \text{H}_{(aq)}^{+}$
- (b) Give an explanation for each of the following observations:
 - (i) The gradual decrease in size (actinoid contraction) from element to element is greater among the actinoids than that among the lanthanoids (lanthanoid contraction).
 - (ii) The greatest number of oxidation states are exhibited by the members in the middle of a transition series.
 - (iii) With the same d-orbital configuration (d^4) Cr^{2+} ion is a reducing agent but Mn^{3+} ion is an oxidising agent.
- **30.** (a) Illustrate the following name reactions by giving example:
 - (i) Cannizzaro's reaction
 - (ii) Clemmensen reduction
 - (b) An organic compound A contain 69.77% carbon, 11.63% hydrogen and rest oxygen. The molecular mass of the compound is 86. It does not reduce Tollen's reagent but forms an addition compound with sodium hydrogen sulphite and gives positive iodoform test. On vigorous oxidation it gives ethanoic and propanoic acids. Derive the possible structure of compound 'A'.

- (a) How are the following obtained?
 - (i) Benzoic acid from ethyl benzene.
 - (ii) Benzaldehyde from toluene.
- (b) Complete each synthesis by giving the missing material, reagent or products:

(i)
$$C_6H_5COC1$$
 $\xrightarrow{H_2}$

CBSE (Delhi) SET-II

Questions different from Set-I

- 1. Which point defect in its crystal units alters the density of a solid?
- **4.** Which is a stronger oxidising agent Bi(V) or Sb(V)?
- **21.** What is the difference between multimolecular and macromolecular colloids? Give one example of each. How are associated colloids different from these two types of colloids?
- **24.** Explain the following observations:
 - (i) Fluorine does not exhibit any positive oxidation state.
 - (ii) The majority of known noble gas compounds are those of Xenon.
 - (iii) Phosphorus is much more reactive than nitrogen.
- **27.** How do antisepites differ from disinfectants? Give one example of each type.
- **28.** (a) Complete the following chemical reaction equations:

(i) Fe
$$^{2+}(aq)$$
 MnO $^{-}_{4(aq)}$ H (aq)

(ii)
$$\operatorname{Cr_2O}_{7(aq)}^{2-}$$
 I $_{(aq)}$ H $_{(aq)}$

- (b) Explain the following observations:
 - (i) Transition elements are known to form many interstitial compounds.
 - (ii) With the same d^4 d-orbital configuration Cr^{2+} ion is reducing while Mn^{3+} ion is oxidising.
 - (iii) The enthalpies of atomisation of the transition elements are quite high.

OR

- (a) Complete the following chemical reaction equations:
 - (i) $\text{CrO}_7^{2-}_{(aq)} + \text{H}_2\text{S}_{(g)} + \text{H}^+_{(aq)}$
 - (ii) $MnO_{2(s)} + KOH_{(aq)} + O_{(2)}$
- (b) Explain the following observations:
 - (i) Transition metals form compounds which are usually coloured.
 - (ii) Transition metals exhibit variable oxidation states.
 - (iii) The actinoids exhibit a greater range of oxidation states than the lanthanoids.
- **29.** (a) What type of a cell is the lead storage battery? Write the anode and the cathode reactions and the overall reaction occurring in a lead storage battery while operating.
 - (b) A voltaic cell is set up at 25°C with the half-cells Al | Al³⁺ (0.001 M) and Ni | Ni²⁺ (0.50 M). Write the equation for the reaction that occurs when the cell generates an electric current and determine the cell potential.

(Given:
$$E^{o}_{Ni^2 \mid Ni} = -0.25V$$
, $E^{o}_{Al^3 \mid Al} = -1.66V$).

OR

- (a) Express the relationship amongst cell constant, resistance of the solution in the cell and conductivity of the solution. How is molar conductivity of a solute related to conductivity of its solution.
- (b) Calculate the equilibrium constant for the reaction.

$$Fe_{(s)}$$
 Cd^2 (aq) \rightleftharpoons Fe^2 (aq) $Cd_{(s)}$

(Given:
$$E^{o}_{Cd^2 \mid Cd}$$
 – 0.40 V, $E^{o}_{Fe^2 \mid Fe}$ – 0.44 V).

CBSE (Delhi) SET-III

Ouestions different from Set-I and Set-II

- 1. Which point defect in its crystal units increases the density of a solid?
- **8.** What does the part '6, 6' mean in the name nylon-6,6?
- 19. Calculate the freezing point depression expected for 0.0711m aqueous solution of Na₂SO₄. If this solution actually Freezes at 0.320°C, what would be the value of van't Hoff factor? (K_f for water is 1.86°C mol 1).
- 24. Compare the following complexes with respect to their shape, magnetic behaviour and the hybrid orbitals involved:
 - (*i*) $[CoF_4]^{2-}$

(ii) $[Cr(H_2O)_2(C_2O_4)_2]^{-1}$

(iii) [Ni(CO)₄]

(Atomic number: Co = 27, Cr = 24, Ni = 28)

- 27. What are the following substances? Give one example of each type.
 - (i) Antacid
- (ii) Nonionic detergents
- (iii) Antiseptics
- **28.** (a) Complete the following chemical reaction equations:
 - (i) $Cr_2O_7^2$ (aq) + I (aq) + H (aq)
 - (ii) MnO_{4 (aq)} + Fe²⁺_(aq) + H _(aq)
 - (b) Explain the following observations:
 - (i) In general the atomic radii of transition elements decrease with atomic number in a given series.
 - (ii) The $E^{0}_{M^{2}|M}$ for copper is positive (+ 0.34 V). It is the only metal in the first series of transition elements showing this type of behaviour.
 - (iii) The E^o value for $Mn^{3+} \mid Mn^{2+}$ couple is much more positive than for $Cr^{3+} \mid Cr^{2+}$ or $Fe^{3+} \mid Fe^{2+}$ couple.

OR

- (a) What is meant by the term lanthanoid contraction? What is it due to and what consequences does it have on the chemistry of elements following lanthanoids in the periodic table?
- (b) Explain the following observations:
 - (i) Cu⁺ ion is unstable in aqueous solutions.
 - (ii) Although Co^{2+} ion appears to be stable, it is easily oxidised to Co^{3+} ion in the presence of a strong ligand.
 - (iii) The $E^{o}_{Mn^{2+}|Mn}$ value for manganese is much more than expected from the trend for other elements in the series.
- **29.** (a) Define the term molar conductivity. How is it related to conductivity of the related solution?
 - (b) One half-cell in a voltaic cell is constructed from a silver wire dipped in silver nitrate solution of unknown concentration. Its other half-cell consists of a zinc electrode dipping in 1.0 M solution of Zn(NO₃)₂. A voltage of 1.48 V is measured for this cell. Use this information to calculate the concentration of silver nitrate solution used.

$$(E_{Zn^2/Zn}^o = 0.76V, E_{Ag^2/Ag}^o = 0.80V)$$

OR

- (a) Corrosion is essentially an electrochemical phenomenon. Explain the reactions occurring during corrosion of iron kept in an open atmosphere.
- (b) Calculate the equilibrium constant for the equilibrium reaction.

$$\operatorname{Fe}_{(s)} \operatorname{Cd}^{2}_{(aq)} \Longrightarrow \operatorname{Fe}^{2}_{(aq)} \operatorname{Cd}_{(s)}$$

(Given: $E^{o}_{Cd^2/Cd}$ 0 40V, $E^{o}_{Fe^2/Fe}$ 0 44V)

SOLUTIONS

CBSE (Delhi) SET-I

- 1. Frenkel defect.
- 2. The scattering of light by colloidal particles is known as Tyndall effect.
- 3. As only sulphide ore particles are wetted by oil while gangue particles are wet by water.
- **4.** Because Bi (V) is more stable than Sb (V) due to inert pair effect.
- **5.** 2-Bromo-3-methyl but-2-en-1-ol.
- 7. Alkyl amine is more basic than ammonia because the + I effect or electron donating nature of alkyl group increases electron density on 'N' atom in alkyl amine.
- 8. Buna-S, neoprene
- **9.** Let the rate law be, $r_1 = k [A]^2$
 - (i) If [A] is doubled than rate $r_2 = k(2A)^2 = 4k$ $[A]^2 = 4r_1$, i.e., rate becomes 4 times.
 - (ii) If [A] is reduced to half then rate, r_3 $k = \frac{A}{2} = \frac{1}{4}k[A]^2 = \frac{1}{4}r_1$, i.e., rate becomes $\frac{1}{4}$ times
- **10.** (*i*) Role of cryolite
 - It lowers the melting point of the mixture.
 - It makes alumina a good conductor of electricity.
 - (ii) When nickel is heated with carbon monoxide it forms a volatile complex nickel tetracarbonyl which on further heating at higher temperature decomposes to give pure nickel.

11. (i) No. of electron pairs around central atom (Xe) = $6 \frac{b.p.}{l.p.}$ 2


The shape would be square planar.

(ii) No. of electron pairs around central atom (Br) = $5 \frac{b.p.}{l.p.}$ 2

The shape would be slightly bent T.

12. (*i*) P₄ 3NaOH 3H₂O PH₃ 3NaH₂PO Phosphine





Molality is independent of temperature whereas molarity changes with change in temperature as volume changes with temperature.

14. (i) CH_2Cl : It is primary halide therefore undergoes S_N^2 reaction faster.

(ii) : As iodine is a better leaving group because of its large size, it will be released at a faster rate in the presence of incoming nucleophile

15.
$$(i)$$
 OH SOCl₂ Cl SO₂ + HCl CH₂—OH HCl HO O

- **16.** (*i*) The C—NH— bond formed between two amino acid molecules with loss of water in a polypeptide is called peptide linkage.
 - (ii) The linkage between two monosaccharides molecules through oxygen atom in a disaccharide or polysaccharide is known as glycosidic linkage.
- 17. B group vitamins and vitamins C are soluble in water.

	Name of Vitamins	Sources	Deficiency diseases	
(<i>i</i>)	Vitamin B ₁₂	Meat, fish, egg and curd	Pernicious anaemia	
(ii)	Vitamin C	Citrus fruits and amla	Scurvy	

18.

	Polymer	Monomer	Structure	
(i)	Teflon	Tetrafluoroethene	F F F C C F	
(ii)	Polyethene	Ethene	Н Н Н С С Н	

OR

n

52 | Xam Idea Chemistry-XII

19.
$$d = 7.87 \text{ g cm}^{-3}$$
, For bcc , $z = 2$, $M = 56 \text{ g mol}^{-1}$
 $a = 286.65 \text{pm} = 286.65 \times 10^{-10} \text{cm}$, $N_A = ?$
 $d = \frac{z M}{a^3 N_A}$
7.87g cm⁻³ $\frac{2 56 \text{ g mol}^{-1}}{(286.65 10^3 \text{ cm})^3 N_A}$
 $N_A = \frac{2 56 \text{ g mol}^{-1}}{(2.8665 10^8 \text{ cm})^3 7.87 \text{ g cm}^{-3}}$
 $= \frac{2 56 \text{ g mol}^{-1}}{23.553 10^{-24} \text{ cm}^3 7.87 \text{ g cm}^{-3}}$
 $= \frac{112 \text{ mol}^{-1}}{185.366 10^{-24}} = 0.604 \times 10^{24} \text{ mol}^{-1}$
 $= 6.024 \times 10^{23} \text{ mol}^{-1}$
 $= \frac{W_B R T}{M_B V}$
 $M_B = \frac{W_B R T}{M_B V}$

$$W_{\rm B} = 100 \text{ mg} = 100 \times 100^{-3} \text{g}$$

$$R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

$$T = 25^{\circ}\text{C} = (25 + 273) \text{ K} = 298 \text{K}$$

$$V = 10 \text{ mL} = 10 \times 10^{-3} \text{L}$$

$$= 13.3 \text{ mm Hg} \quad \frac{13.3}{760} \text{ atm}$$

$$M_{\rm B} = \frac{100 \quad 10^{-3} \text{ g}}{13.3 \text{ atm}} \quad \frac{10 \quad 10^{-3} \text{ L}}{10^{-3} \text{ L}}$$

$$M_{\rm B} = \frac{100 \quad 10^{-3} \quad 0.0821 \quad \text{L atm K}^{-1} \text{ mol}^{-1}}{133 \quad 10^{-3} \quad 10^{-3}} = \frac{18594 \text{ g mol}^{-1}}{133}$$

$$M_{\rm B} = \quad 13980.45 \text{ g mol}^{-1}$$

21. For a first order reaction

$$t = \frac{2.303}{k} \log \frac{[R]_o}{[R]}$$

Here

$$t = 3 \text{ h} = 3 \times 60 \text{ min} = 180 \text{ min}$$

$$k = 0.0051 \text{min}^{-1}, [R]_0 = 0.10 \text{ M}, [R] = ?$$

$$180 \text{ min} = \frac{2.303}{0.0051 \text{min}^{-1}} \log \frac{0.10}{[R]}$$

$$\text{Log} \frac{0.1}{[R]} = \frac{180 \text{ min} \quad 0.0051 \text{ min}^{-1}}{2.303} = \frac{918}{2303}$$

$$\log \frac{0.1}{R} = 0.3986$$

$$\frac{0.1}{R} = \text{Anti} \log (0.3986) = 2.503$$

$$R = \frac{0.1}{2.503} = 0.03995 \text{ M}$$

R = 0.04M

22.

Type of Colloid	Dispersed Phase	Dispersion medium	Example	
Aerosol	Solid or liquid	Gas	Smoke, Fog, Cloud	
Hydrosol	Solid	Water	Starch sol, Protein sol	
Emulsion	Liquid	Liquid	Milk, hair cream	

- (i) NH_3 is stronger base than PH_3 . This is because the lone pair of electrons on N atom in NH_3 is 23. directed and not diffused as it is in PH₃ due to larger size of phosphorus and hence more available for donation.
 - (ii) Sulphur has a greater tendency for catenation than oxygen because S-S bond is stronger than O – O bond due to less interelectronic repulsions.
 - (iii) Bond dissociation energy of F₂ is less than Cl₂ this is due to relatively large electron electron repulsion among the lone pairs in F₂ molecule where they are much closer to each other than in case of Cl₂.

(i) H—O—
$$\vec{N}$$
 $\stackrel{\ddot{0}:}{\bar{0}:}$ $\stackrel{\ddot{0}:}{\longleftarrow}$ HO— \vec{N} $\stackrel{\ddot{0}:}{\bar{0}:}$

As a result of resonance, N-O bond length is average of single bond and double bond whereas N-OH bond has purely single bond character. Therefore, N-O bond is shorter than N-OH bond in HNO₃.

(ii) S atom in SF₄ is not sterically protected as it is surrounded by only four F atoms, so attack of H₂O molecules can take place easily and hence hydrolysis takes place easily. In contract, in SF₆, S is sterically protected by six F atoms. Therefore does not allow H₂O molecules to attack S atoms. As a result of this, SF₆ does not undergo hydrolysis.

54 | Xam idea Chemistry-XII

- (iii) In XeF₂, Xe is sp³d hybridised having 2 bond pair and 3 lone pair of electrons. The presence of 3 lone pair of electrons in Xe F₂ at equidistance to have minimum repulsion is responsible for its linear structure.
- 24. Given complex is Fe (en)₂ Cl₂ Cl
 - (i) Let the oxidation number of iron be x.

(ii) Orbitals of Fe (III)



 $d^2 sp^3$ hybridised orbitals of Fe (III)

Thus, hybridisation: $d^2 sp^3$

Shape of the complex: Octahedral

- (iii) Paramagnetic due to presence of three unpaired electrons
- (iv) Two, cis and trans isomers
- (v) Yes, cis isomer will also show optical isomerism
- (vi) Dichlorido bis (ethane-1, 2 diamine) iron (III) chloride or Dichloro bis (ethylenediamine) iron (III) chloride.
- 25. (i) Step I: Nucleophilic addition of Grignard reagent to Carbonyl group.


$$\begin{bmatrix} \searrow_{C} - OMg - X \end{bmatrix} \xrightarrow{H_2O} \searrow_{R} C - OH + Mg(OH)X$$

Step II: Hydrolysis

(ii)
$$CH_3 - CH_2 - OH$$
 H $CH_2 = CH_2$ H_2O

Mechanism

Step I: Formation of protonated alcohols



Step II: Formation of carbocation: It is the slowest step and hence, the rate determining step.

Step III: Formation of ethylene by elimination of a proton

To drive the equilibrium to the right, ethylene is removed as it is formed.

(iii)
$$> C = C < + H_2O \xrightarrow{H^+} > C = C < Alkene$$

Mechanism

Step I: Protonation of alkene to form carbocation by electrophilic attack of H₃O⁺.

Step II: Nucleophilic attack of water on carbocation.

$$-\overset{H}{\overset{}{\subset}} + \overset{H}{\overset{}{\hookrightarrow}} \overset{H}{\overset{}{\hookrightarrow}} -\overset{H}{\overset{}{\hookrightarrow}} -\overset{H}{\overset{}{\hookrightarrow}} -\overset{H}{\overset{}{\hookrightarrow}} +$$

Step III: Deprotonation to form an alchohol

26. (i) Hoffman's bromamide reaction: When a primary acidamide is heated with bromine in an aqueous or ethanolic solution of sodium hydroxide, it gives a primary amine with one carbon atom less. In this degradation reaction, migration of an alkyl or aryl group takes place from carbonyl carbon of the amide to the nitrogen atom.

$$R - C - NH_2 + Br_2 + 4NaOH \longrightarrow R - NH_2 + Na_2CO_3 + 2NaBr + 2H_2O$$

1° acid amide

1° amine

(ii) Gattermann reaction: Chlorine or bromine can be introduced in benzene ring by treating the diazonium salt solution with corresponding halogen acid in the presence of copper powder.

(iii) Coupling reaction: Diazonium salts react with aromatic amines in weakly acidic medium and with phenols in weakly alkaline medium to form coloured compounds called azo dyes by coupling at para position of amines or phenols. The mechanism is basically that of electrophilic aromatic substitution where the diazonium ion is electrophile.

Benzene diazonium chloride
$$NH_2 = NH_2 + N$$

27. (*i*) Cationic Detergents: Cationic detergents are quarternary ammonium salts of amines with acetates, chlorides or bromides as anions. Cationic part possess a long hydrocarbon chain and a positive charge on nitrogen atom. Hence, these are called cationic detergents. Cetyltrimethylammonium bromide is a popular cationic detergent and is used in hair conditioners.

$${\rm CH_3}$$
 ${\rm CH_3}$ ${\rm CH_3}$ ${\rm CH_3}$ ${\rm Br}$ ${\rm CH_3}$

Cetyltrimethyl ammonium bromide

Cationic detergents have germicidal properties and are expensive, therefore, these are of limited use.

(ii) Food Preservatives: These are the chemical substances which are added to the food materials to prevent their spoilage due to microbial growth and to retain their nutritive value for long periods.


Preservatives prevent the rancidity of food and inhibit growth or kill the microganisms.

The most-common preservations used are, sugar table salt, sodium benzoate, Sodium metabisulphite sorbic acid and propanoic acid.

- (iii) Analgesics: Drugs which reduce or abolish pain without causing impairment of consciousness, mental confusion, incoordination or paralysis or some other disturbances of nervous system. These are classified as follow.
 - (a) Non-narcotic (non-addictive) analgesics: Aspirin, paracetamol etc.
 - (b) Narcotic drugs: These are known to be habit forming, e.g., morphine, Codeine, Heroin
- 28. (a) Molar Conductivity (m): It may be defined as the conductance of a solution containing 1 mole of electrolyte such that the entire solution is placed is between two electrodes one centimeter apart.

$$m = k \times v$$
or
$$m = \frac{k \quad 1000}{M}$$

Molar conductivity increases with decrease in concentration or increase in dilution as number of ions as well as mobility of ions increased with dilution.



For strong electrolytes the number of ions do not increase appreciably on dilution and only mobility of ions increases due to decrease in inter-ionic attractions. Therefore m increases a little as shown in graph by a straight line.

For weak electrolyte the number of ions as well as mobility of ions increases on dilution as shown by curve in the figure.

Here
$$E_{\text{cell}}^{0} = 0.46 \text{ V}, n = 2$$

[Ag] 0 001M 1
$$10^{-3}$$
 M, [Cu²] 0.1M

$$E_{cell} = 0 46 \frac{0.0591}{2} log \frac{0.1}{(10^{-3})^2}$$

$$E_{cell} = 0 46 \frac{0.0591}{2} log 10^5 0 46 \frac{0.0591}{2} 5 log 10$$

$$E_{cell} = 0.46 + 0.0591 + 2.5 + 1 + 0.46 + 0.14775 = 0.31225V$$
 $E_{cell} = 0.312 \text{ V}$

OR

$$(a) = \frac{1}{R} \quad \frac{l}{A}$$

Where = Conductivity $\frac{l}{A} = \text{Cell Constant}$

R = Resistance

$$_{m} = \frac{1000}{M}$$

Where m = Molar conductivity

= Conductivity

M = Molarity of Solution

(b) At anode: Al(s) Al
3
 (aq) $3\overline{e}$] 2

At cathode: Ni^2 (aq) $2\bar{e}$ Ni(s)] 3

$$E_{cell} = E_{cell}^{o} = \frac{0.0591}{n} log \frac{[Al^3]^2}{[Ni^2]^3}$$

Here n = 6, $[A1^{3+}] = 0.001M = 1 \times 10^{-3}M$, $[Ni^{2+}] = 0.5M$

$$E_{\text{cell}}^{\text{o}} = E_{Ni^2/Ni}^{\text{o}} E_{Al^3/Al}^{\text{o}} = 0.25V$$
 1.66V

 $E_{\text{cell}}^{\text{o}} = 1.41 \text{V}$

$$E_{\text{cell}}^{0} = 1.41 \quad \frac{0.0591}{6} \log \frac{10^{-3}}{0.5^{-3}} \quad 1.41 \quad \frac{0.0591}{6} \log \frac{10^{-6}}{0.125}$$

$$= 1.41 \quad \frac{0.0591}{6} \log 10^{-6} \quad 8 \quad 1.41 - \frac{0.0591}{6} \quad \log 10^{-6} \quad \log 2^{3}$$

$$= 1.41 \quad \frac{0.0591}{6} \log - 6 \log 10 \quad 3 \log 2 \quad 1.41 - \frac{0.0591}{6} \quad -6 \quad 3 \quad 0.3010$$

$$= 1.41 \quad \frac{0.0591}{6} \quad (5.097) = 1.41 \quad \frac{0.3012}{6}$$

$$= 1.41 + 0.0502 \quad = 1.4602V$$

$$E_{cell} = 1.46 \text{ V}$$

(ii) In acidic medium:

$$Cr_2 O_7^{2-} + 14H^+ + 6\bar{e}$$
 $2Cr^3 7H_2O$
 $Fe^2 Fe^3 \bar{e}_] 6$
 $Cr_2 O_7^{2-} + 6Fe^{2+} + 14H$ $2Cr^3 6Fe^{3+} 7H_2O$

- (b) (i) On moving from titanium to copper in general atomic mass increases where as atomic size decreases, therefore density increases in general.
 - (ii) The frequent metal-metal bonding in compounds of heavy transition elements is due to their high enthalpy of atomization.
 - (iii) This is due to very small energy gap between 5f, 6d and 7s orbitals in the actinoid series.

OR

(a) (i) In neutral or faintly alkaline solutions

$$MnO_4$$
 $2H_2O$ $3e$ MnO_2 $4OH^-$] 8 $S_2O_3^2$ $10OH$ $2SO_4^2$ $5H_2O$ $8e$] 3 $8MnO_4$ $3S_2O_3^2$ $+H_2O$ $8MnO_2$ $6SO_4^2$ $+2OH$

(ii) In acidic solutions

(b)
$$\operatorname{Fe}(s)$$
 $\operatorname{Cd}^2(aq) \Longrightarrow \operatorname{Fe}^2(aq)$ $\operatorname{Cd}(s)$ $\log k_c \quad n \frac{E_{\text{cell}}^0}{0.059}$

Here,
$$n$$
 2
$$E_{\text{cell}}^{0} = E_{\text{cathode}}^{0} \quad E_{\text{anode}}^{0} = E_{\text{Cd}^{2+}/\text{Cd}}^{0} \quad E_{\text{F}e^{2+}/\text{Fe}}^{0}$$

$$= -40 - (-0.44) = 0.04\text{V}$$

$$\log k_{c} \quad \frac{2 \quad 0.04}{0.059} \quad \frac{0.08}{0.059}$$

$$\log k_c = 1.3536$$
 $k_c = \text{Antilog } 1.3536 = 22.57$

30. (a) (i) **Cannizzaro reaction:** Aldehyde which do not have an hydrogen atom, undergo disproportionation reaction on treatment with concentrated alkali, In this reaction, one molecule of aldehyde is reduced to alcohol while other is oxidised to carboxylic acid salt.

(ii) Clemmensen reduction: The carbonyl group of aldehyde and ketones is reduced to CH₂ group on treatment with zinc amalgam and concentrated hydrochloric acid.

$$C = O + 4[H]$$
 $C = O + 4[H]$
 $CH_2 + H_2O$
Aldehyde or Hydrocarbon

(*b*)

Element	Percentage	Atomic mass	No. of moles	Simplest molar ratio
С	69.77	12	$\frac{69\ 77}{12}$ 5 81	$\frac{581}{116}$ 5
Н	11.63	1	$\frac{11 \ 63}{1} \ 11 \ 63$	$\frac{11 \ 63}{1 \ 16} \ 10$
О	(100 - 81.4) = 18.60	16	$\frac{18 \ 60}{16}$ 1 16	$\frac{1}{1}\frac{16}{16}$ 1

Empirical formula of the compound $A = C_5 H_{10} O$

Molecular formula of the compound A = n (Empirical formula)

$$n = \frac{\text{Molecular mass of compound } A}{\text{Empirical formula mass of compound } A}$$

Molecular mass of compound A = 86

Empirical formula mass of compound A = 5 12 1 10 1 16

$$= 60 + 10 + 16$$

$$= 86$$

$$n = \frac{86}{86} \quad 1$$

Molecular formula of the compound $A = 1(C_5 H_{10} O)$

$$= C_5 H_{10} O$$

- As the compound A forms addition compound with NaHSO₃ therefore it must be either an aldehyde or ketone.
- As it does not reduce Tollens reagent and give positive Iodoform test therefore it must be a methyl ketone.
- As on oxidation the compound A gives a mixture of ethanoic acid and propanoic acid, therefore compound A is

$$CH_{3} - C - CH_{2} - CH_{2} - CH_{3}$$
Pentan-2-one

The chemical reactions are:

$$\begin{array}{c} O \\ || \\ CH_3-C-CH_2-CH_2-CH_3 \\ Pentan-2-one \end{array} \quad \begin{array}{c} OH \\ | \\ CH_3-C-CH_2-CH_2-CH_3 \\ | \\ SO_3 \; Na \end{array}$$

Sodium hydrogen sulphite addition product

Benzaldehyde

$$\begin{array}{c} O \\ || \\ CH_3-C-CH_2-CH_2-CH_3-3I_2-4NaOH \\ \hline \\ & \begin{array}{c} Iodoform \\ reaction \end{array} \begin{array}{c} CHI_3 \\ (Iodoform\ yellow\ ppt) \end{array} \begin{array}{c} CH_3-CH_2-CH_2-COONa -3NaI-3H_2O \\ \end{array}$$

CH₃

$$+ \text{ CrO}_2\text{Cl}_2 \xrightarrow{\text{CS}_2} \xrightarrow{\text{Chromium complex}} \xrightarrow{\text{H}_3\text{O}^+} \xrightarrow{\text{Benzaldehyde}}$$

Benzylidene

diacetate

$$(b) (i) \qquad \begin{array}{c} O \\ C - Cl \\ \hline \\ Benzoyl \ chloride \\ \hline \\ (ii) \\ \hline \\ Benzene \\ \hline \\ Benzene \\ \hline \\ Benzene \\ \hline \\ Benzoyl \ chloride \\ \hline \\ Benzophenone \\ \hline \\ Benzophenone \\ \hline \\ Benzophenone \\ \hline \\ C - Cl \\ \hline \\ Acetyl \ chloride \\ \hline \\ Benzophenone \\ \\ Benzophenone \\ \hline \\ Benzophenone \\ \\ Benzophenone \\ \hline \\ Benzophenone \\ \\$$

CBSE (Delhi) SET-II

- 1. Schottky defect
- **21. Multimolecular colloids:** In this type of colloids, colloidal particles are aggregates of atoms or molecules each having size less than 1nm, e.g., sulphur sol, gold sol.

Macromolecular colloids: In this type of colloids, colloidal particles are themselves large molecules of colloidal dimensions, e.g., starch, proteins, polyethene, etc.

Associated colloids: There are certain substances which at low concentrations behave as normal electrolyte, but at higher concentrations exhibit colloidal behaviour due to the formation of aggregates. Such colloids are known as associated colloids, e.g., soaps and detergents.

- **24.** (*i*) Fluorine does not exhibit any positive oxidation state as it is the most electro-negative element and does not have *d* orbitals in its valance shell.
 - (ii) This is due to low ionization enthalpy of xenon.
 - (iii) Phosphorus is much more reactive than nitrogen as the P-P single bond (213 kJ mol ¹) is much weaker than N N triple bond (941.4 kJ mol ¹).
- **27.** Difference between antiseptics and disinfectants.
 - Antiseptics are chemical substances which prevent the growth of micro-organisms and may even kill them but are not harmful to living tissues.
 - Antiseptics are generally applied to living tissues such as wounds, cuts, ulcers and diseased skin surfaces.
 - Dettol, furacine, soframicine are antiseptics.

- Disinfectants are applied to inanimate objects such as floor, drainage system, instrument, etc.
- Chlorine in the concentration of 0.2 to 0.4 ppm in aqueous solution and SO₂ in very low concentration are disinfectants.

28.	(a)	(<i>i</i>)	MnO_4	8H	5e	$Mn^2 4H_2O$
				F	e^2	Fe ³ e] 5
			MnO ₄	5Fe ²	8H	Mn^{2} $5\mathrm{Fe}^{3}$ $4\mathrm{H}_{2}\mathrm{O}$
		(ii)	Cr_2O_7^2	14H	6e	$2Cr^3$ $7H_2O$
					2I	I ₂ 2e] 3
			$Cr_2O_7^2$	6I	14H	2Cr ³ 3I ₂ 7H ₂ O

- (b) (i) Transition elements form many interstitial compounds as they are capable of entrapping small atoms like H, C or N in the interstitial sites in their crystal lattice.
 - (ii) Cr^2 is reducing as its configuration changes from d^4 to d^3 , the latter having a half filled $\operatorname{t_{2g}}$ configuration. On the other hand, the change from Mn^3 to Mn^2 results in the half-filled d^5 configuration which has extra stability therefore Mn^3 is oxidising.
 - (iii) This is because transition metals have strong metallic bonds as they have a large number of unpaired electrons.

(a) (i)
$$Cr_2O_7^2$$
 14H 6e $2Cr^3$ 7H₂O H₂S S 2H 2e] 3
$$Cr_2O_7^2$$
 3H₂S 8H $2Cr^3$ 3S 7H₂O
(ii) 2MnO₂ 4KOH O₂ $2K_2MnO_4$ 2H₂O

- (b) (i) This is due to d d transition as the energy of excitation of d orbital electrons from lower energy to higher energy level lies in the visible region.
 - (ii) The variable oxidation states of transition metals are due to the participation of ns and (n-1)d electrons in bonding as energy difference between ns and (n-1)d orbitals is small.
 - (iii) This is due to comparable energies of 5f, 6d and 7s orbitals of actinoids.
- **29.** (a) The lead storage battery is a secondary cell

The cell reactions when the battery is in use are given below

At anode : Pb(s) SO_4^2 (aq) $PbSO_4(s)$ 2e At cathode : $PbO_2(s)$ SO_4^2 (aq) 4H (aq) 2e $PbSO_4(s)$ 2H₂O(l) Overall cell reaction : Pb(s) $PbO_2(s)$ 2H₂SO₄(aq) 2PbSO₄(s) 2H₂O(l)

64 | Xam idea Chemistry-XII

At anode: Al(s) Al³ (aq)
$$3\bar{e}$$
 2

At cathode: Ni² (aq) $2\bar{e}$ Ni(s)] 3

$$2Al(s) 3Ni2 (aq) 2Al3 (aq) Ni(s)$$

$$E_{cell} = E_{cell}^{0} \frac{0.0591}{n} \log \frac{[Al^{3}]^{2}}{[Ni^{2}]^{3}}$$
Here $n = 6$, $[Al^{3+}] = 0.001M = 1 \times 10^{-3}M$, $[Ni^{2+}] = 0.5M$

$$E_{cell}^{0} = E^{0} \frac{8^{2}}{N^{2}} \frac{1}{N^{1}} E^{0} \frac{10^{-3}}{0.5^{-3}} \frac{1}{1.41} \frac{0.0591}{6} \log \frac{10^{-6}}{0.125}$$

$$= 1.41 \frac{0.0591}{6} \log 10^{-6} 8 1.41 - \frac{0.0591}{6} \log 10^{-6} \log 2^{3}$$

$$= 1.41 \frac{0.0591}{6} \log - 6 \log 10 3 \log 2 1.41 - \frac{0.0591}{6} - 6 3 0.3010$$

$$= 1.41 \frac{0.0591}{6} (5.097) = 1.41 \frac{0.3012}{6}$$

$$= 1.41 + 0.0502 = 1.4602V$$

$$E_{cell} = 1.46 V$$

OR

(a)
$$= \frac{1}{R} \frac{l}{A}$$

Where
$$= \text{Conductivity}$$

$$\frac{l}{A} = \text{Cell Constant}$$

$$R = \text{Resistance}$$

$$m = \frac{1000}{M}$$
Where
$$m = \text{Molar conductivity}$$

$$= C \text{Conductivity}$$

$$= C \text{Solution}$$

(b) $\operatorname{Fe}(s)$ $\operatorname{Cd}^2(aq) \iff \operatorname{Fe}^2(aq)$ $\operatorname{Cd}(s)$

 $\log k_{\rm c}$ $n \frac{E \text{ cell}}{0.059}$

Here, n = 2

E cell = E cathode — E° anode
= E° Cd² /Cd - E° Fe² /Fe = - 0.40 - (- 0.44)

$$E^{\circ}_{cell}$$
 = 0.04 V
 $\log k_{c} = \frac{2 \quad 0.04}{0.059} = \frac{0.08}{0.059}$
 $\log k_{c}$ = 1.3536
 k_{c} = Antilog 1.3536
 k_{c} = 22.57

CBSE (Delhi) SET-III

- 1. Interstitial defect increases the density of a solid.
- It means the two monomers combine to make nylon 6,6, contain six carbon atoms each.

19.
$$T_{f} = [273 \ 15 \ (0 \ 320 \ 273 \ 15)]K = 0 \ 320 \ K$$

$$T_{f} = K_{f} . m$$

$$= 1.86 \ K \ kg \ mol^{-1} \times 0.0711 \ mol \ kg^{-1}$$

$$= 0.132 \ K$$

$$i = \frac{\text{Observed value of}}{\text{Calculated value of}} \frac{T_{f}}{T_{f}} = \frac{0.320 \text{K}}{0.132 \text{K}}$$

$$i = 2.42$$

24.

S. No.	Complex	Central metal ion	Configuration of metal ion	Hybridisation of metal ion	Geometry of the complex	Number of unpaired electrons	Magnetic behaviour
(<i>i</i>)	$[CoF_4]^2$	Co ²	d^7	sp^3	Tetrahedral	3	Paramagnetic
(ii)	$[Cr(H_2O)_2 \\ (C_2O_4)_2]$	Cr ³	$3d^3$	d^2sp^3	Octahedral	3	Paramagnetic
(iii)	Ni(CO) ₄	Ni	$3d^8 4s^2$	sp ³	Tetrahedral	0	Diamagnetic

- 27. (i) Antacids: Chemical substances which removes the excess acid in the stomach & raise the pH to appropriate level, e.g., sodium hydrogen carbonate, a mixture of aluminium and magnesium hydroxide, ranitidine, etc.
 - (ii) Non-ionic Detergents: Non-ionic detergents do not contain any ion in their constitution. One such detergent is formed when stearic acid reacts with polyethylene glycol.
 - (iii) Antiseptics: These are the chemical substances which prevent the growth of micro-organisms and may even kill them but are not harmful to living tissues. Antiseptics are applied to living tissues such as wounds, cuts, ulcers. Dettol, soframicine are antiseptics.

- $2Cr^3$ $7H_2O$ **28.** (*a*) (*i*) $Cr_2 O_7^2$ 14H 6e 21 $Cr_2O_7^2$ 14H 6I $Mn^{\,2}$ MnO_4 H8 5e 4H₂O (ii) Fe^2 Fe^3 5Fe^2 8H 4H₂O MnO_4
 - (b) (i) The atomic radii of transition metals decreases with atomic number in a series as the nuclear charge increases due to poor shielding effect of d orbitals.
 - (ii) This is due to its high enthalpy of atomization and low hydration enthalpy.
 - (iii) This is due to much large third ionisation energy of Mn as Mn^2 is very stable on account of stable d^5 configuration.

OR

(a) Lanthanoid contraction: The steady decrease in the atomic and ionic radii of lanthanoids with increase in atomic number is known as lanthanoid contraction.

Cause of lanthanoid contraction: As we move along the lanthanoid series, for every additional proton in the nucleus, the corresponding electron goes into 4f-subshell, there is poor shielding of one electron by another in this subshell due to the shapes of these f-orbitals. The imperfect shielding is not able to counterbalance the effect of the increased nuclear charge. Thus the net result is decrease in size with increase in atomic number.

Consequences:

- (i) 5d series elements have nearly same radii as that of 4d-series.
- (ii) The basic strength of hydroxides decreases from La(OH)3 to Lu(OH)3.
- (b) (i) Because the high hydration enthalpy of Cu² easily compensates the second ionization enthalpy of Cu.
 - (ii) Because strong ligand cause spin pairing giving rise to diamagnetic octahedral complex which are very stable and have very large crystalfield stabilization energy. This splitting energy overcomes the ionization enthalpy.
 - (iii) This is due to stability of Mn^2 as it has half filled d^5 configuration.
- **29.** (a) Molar conductivity (m): It may be defined as the conductivity of one molar electrolytic solution placed between two electrodes one centimeter apart and have enough area of cross section to hold entire volume.

$$m = \frac{m}{c}$$
Where = Conductivity
$$c = \text{Concentration of solution in mol L}^{-1}$$

(b) At anode :
$$Zn(s)$$
 Zn^2 (aq) 2e

At cathode: Ag
$$(aq)$$
 e Ag (s)] 2

OR

(a) At anode : Oxidation of Fe atoms takes place

Fe Fe² 2e
$$E_{Fe^2/Fe}$$
 0 44V

At cathode: Reduction of oxygen in the presence of H $\,$ ions. The H $\,$ ions are produced by either $\,$ H $_2$ O or $\,$ H $_2$ CO $_3$ (formed by dissolution of $\,$ CO $_2$ in moisture)

2H (aq) 2e 2H
2H
$$\frac{1}{2}$$
 O₂(g) H₂O

Net reaction at cathodic area

2H (aq)
$$\frac{1}{2}$$
O₂ 2e H₂O E _{H /O₂/H₂O} 1.23V

The overall reaction

Fe(s) 2H (aq)
$$\frac{1}{2}$$
O₂(g) Fe² (aq) H₂O(l) E°_{cell} = 1.67V

The ferrous ions are further oxidised by atmospheric oxygen to ferric ions which come out as rust in the form of hydrated ferric oxide (Fe₂O₃.xH₂O).

(b) At anode: Al(s) Al 3 (aq) 3e] 2

At cathode:
$$\operatorname{Ni}^2(aq)$$
 $2\overline{e}$ $\operatorname{Ni}(s)$] 3 $2\operatorname{Al}(s)$ $3\operatorname{Ni}^2(aq)$ $2\operatorname{Al}^3(aq)$ $\operatorname{Ni}(s)$

 $E_{cell} = 1.46 \text{ V}$

$$\begin{split} E_{\text{cell}} &= E_{\text{cell}}^{o} - \frac{0.0591}{n} \log \frac{[\text{AI}^3]^2}{[\text{Ni}^2]^3} \\ \text{Here } &= 6, [\text{AI}^{3+}] = 0.001 \text{M} = 1 \times 10^{-3} \text{M}, [\text{Ni}^{2+}] = 0.5 \text{M} \\ E_{\text{cell}}^{o} &= E_{\text{Ni}^2}^{o} /_{\text{Ni}} - E_{\text{AI}^3}^{o} /_{\text{AI}} - 0.25 V - 1.66 V \\ E_{\text{cell}}^{o} &= 1.41 \text{V} \end{split}$$

$$\begin{split} E_{\text{cell}}^{o} &= 1.41 & \frac{0.0591}{6} \log \frac{10^{-3}}{0.5^{-3}} - 1.41 - \frac{0.0591}{6} \log \frac{10^{-6}}{0.125} \\ &= 1.41 - \frac{0.0591}{6} \log 10^{-6} - 8 - 1.41 - \frac{0.0591}{6} - \log 10^{-6} - \log 2^3 \\ &= 1.41 - \frac{0.0591}{6} \log - 6 \log 10 - 3 \log 2 - 1.41 - \frac{0.0591}{6} - 6 - 3 - 0.3010 \\ &= 1.41 - \frac{0.0591}{6} (-5.097) = 1.41 - \frac{0.3012}{6} \\ &= 1.41 + 0.0502 = 1.4602 \text{V} \end{split}$$

CBSE EXAMINATION PAPERS

ALL INDIA-2009

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (All India) SET-I

- 1. How do metallic and ionic substances differ in conducting electricity?
- 2. What is the 'coagulation' process?
- 3. What is meant by the term 'pyrometallurgy'?
- 4. Why is red phosphorus less reactive than white phosphorus?
- 5. Give the IUPAC name of the following compound:

- **6.** Write the structural formula of 1-phenylpentan-1-one.
- 7. Arrange the following compounds in an increasing order of basic strengths in their aqueous solutions:

1

1

2

- **8.** What does '6, 6' indicate in the name nylon-6,6?
- 9. What type of cell is a lead storage battery? Write the anode and the cathode reactions and the overall cell reaction occurring in the use of a lead storage battery.

OR

Two half cell reactions of an electrochemical cell are given below:

$$MnO_4(aq)$$
 8H (aq) 5e $Mn^2(aq)$ 4H₂O(l), E 1.51V $Sn^2(aq)$ $Sn^4(aq)$ 2e , E 0.15 V

Construct the redox equation from the two half cell reactions and predict if this reaction favours formation of reactants or product shown in the equation.

- **10.** Define the following:
 - (i) Elementary step in a reaction
 - (ii) Rate of a reaction

70 | Xam idea Chemistry-XII

- 11. Describe the underlying principle of each of the following metal refining methods:
 - (i) Electrolytic refining of metals
 - (ii) Vapour phase refining of metals.
- **12.** Complete the following chemical reaction equations:


2

2


2

2

- (i) $XeF_2 + H_2O$
- (ii) PH₃ + HgCl₂
- **13.** Complete the following chemical reaction equations:
 - (i) $MnO_4(aq) + C_2O_4^2(aq) + H^+(aq)$
 - (ii) $Cr_2O_7^2$ (aq) + Fe^2 (aq) + $H^+(aq)$
- **14.** Which one in the following pairs undergoes S_N1 substitution reaction faster and why?



15. Complete the following reaction equations:



- (ii) CH₃CH₂CH CH₂ HBr
- **16.** Name the four bases present in DNA. Which one of these is not present in RNA?
- 17. Name two fat soluble vitamins, their sources and the diseases caused due to their deficiency in diet. 2
- 18. Differentiate between molecular structures and behaviours of thermoplastic and thermosetting polymers. Give one example of each type.
- 19. A first order reaction has a rate constant of 0.0051 min ¹. If we begin with 0.10 M concentration of the reactant, what concentration of the reactant will be left after 3 hours?
- 20. Silver crystallises with face-centred cubic unit cells. Each side of the unit cell has a length of 409 pm. What is the radius of an atom of silver? (Assume that each face atom is touching the four corner atoms.)
- 21. A copper-silver cell is set up. The copper ion concentration in it is 0.10 M. The concentration of silver ion is not known. The cell potential measured 0.422 V. Determine the concentration of silver ion in the cell.3

(Given:
$$E^{o}_{Ag\ /Ag}$$
 0.80V, $E^{o}_{Cu^{2}\ /Cu}$ 0.34 V)

22. What happens in the following activities and why?

3

3

- (i) An electrolyte is added to a hydrated ferric oxide sol in water.
- (ii) A beam of light is passed through a colloidal solution.
- (iii) An electric current is passed through a colloidal solution.
- 23. Giving a suitable example for each, explain the following:
 - (i) Crystal field splitting
 - (ii) Linkage isomerism
 - (iii) Ambidentate ligand.

OR

Compare the following complexes with respect to structural shapes of units, magnetic behaviour and hybrid orbitals involved in units:

$$\left[\text{Co(NH}_3)_6\right]^3$$
, $\left[\text{Cr(NH}_3)_6\right]^3$, Ni(CO)₄

(At No.: Co = 27, Cr = 24, Ni = 28)

24. Explain the following observations:

3

- (i) The boiling point of ethanol is higher than that of methoxymethane.
- (ii) Phenol is more acidic than ethanol.
- (iii) o-and p-nitrophenols are more acidic than phenol.
- **25.** How would you account for the following:

3

- (i) Many of the transition elements and their compounds can act as good catalysts.
- (ii) The metallic radii of the third (5d) series of transition elements are virtually the same as those of the corresponding members of the second series.
- (iii) There is a greater range of oxidation states among the actinoids than among the lanthanoids.
- **26.** Complete the following reaction equations:

3

$$\begin{array}{c} \text{O} \\ \parallel \\ \text{(i) R C NH}_2 \end{array} \quad \begin{array}{c} \text{LiAlH}_4 \\ \text{H}_2 \text{O} \end{array}$$

- (ii) C₆H₅N₂Cl + H₃PO₂+H₂O
- (iii) $C_6H_5NH_2 + Br_2$ (aq)
- **27.** Describe the following substances with one suitable example of each type:

3

- (i) Non-ionic detergents
- (ii) Food preservatives
- (iii) Disinfectants
- **28.** (a) Define the following terms:

5

- (i) Mole fraction
- (ii) Van't Hoff factor

72 | Xam idea Chemistry-XII

(b) 100 mg of a protein is dissolved in enough water to make 100 mL of a solution. If this solution has an osmotic pressure 13.3 mm Hg at 25° C, what is the molar mass of protein?

 $(R = 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1} \text{ and } 760 \text{ mm Hg} = 1 \text{ atm.})$

OR

What is meant by:

- (i) Colligative properties
- (ii) Molality of a solution.
- (b) What concentration of nitrogen should be present in a glass of water at room temperature? Assume a temperature of 25° C, total pressure of 1 atmosphere and mole fraction of nitrogen in air of 0.78. [K_H for nitrogen = 8.42×10^{-7} M/mm Hg]
- **29.** (a) Draw the structures of the following:
 - (i) H₂S₂O₈
 - (ii) HClO₄
 - (b) How would you account for the following:
 - (i) NH₃ is a stronger base than PH₃.
 - (ii) Sulphur has a greater tendency for catenation than oxygen.
 - (iii) F_2 is a stronger oxidising agent than Cl_2 .

OR

5

5

- (a) Draw the structures of the following:
 - (i) H₂S₂O₇
 - (ii) HClO₃
- (b) Give an explanation for each of the following observations:
 - (i) In the structure of HNO₃, the N − O bond(121 pm) is shorter than the N − OH bond (140 pm)
 - (ii) All the P Cl bonds in PCl₅ are not equivalent.
 - (iii) ICI is more reactive than I_2 .
- **30.** (a) Write chemical equations to illustrate the following name bearing reactions:
 - (i) Cannizzaro's reaction
 - (ii) Hell-Volhard-Zelinsky reaction
 - (b) Give chemical tests to distinguish between the following pairs of compounds:
 - (i) Propanal and Propanone
 - (ii) Acetophenone and Benzophenone
 - (iii) Phenol and Benzoic acid

1

3

- (a) How will you bring about the following conversions:
 - (i) Ethanol to 3-hydroxybutanal
 - (ii) Benzaldehyde to Benzophenone.
- (b) An organic compound A has the molecular formula C₈H₁₆O₂. It gets hydrolysed with dilute sulphuric acid and gives a carboxylic acid B and an alcohol C. Oxidation of C with chromic acid also produced B. C on dehydration reaction gives but-1-ene. Write equations for the reactions involved.

CBSE (All India) SET-II

[Questions different from Set-I]

- 1. Which point defect of its crystals decrease the density of a solid?
- 8. What is the primary structural feature necessary for a molecule to make it useful in a condensation polymerisation reaction?
- **20.** For a decomposition reaction the values of rate constant k at two different temperatures are given below:

$$k_1 = 2.15 \times 10^{-8} \text{ L mol}^{-1} \text{ s}^{-1} \text{ at } 650 \text{ K}$$

 $k_2 = 2.39 \times 10^{-7} \text{ L mol}^{-1} \text{ s}^{-1} \text{ at } 700 \text{ K}$

Calculate the value of activation energy for this reaction.

$$(R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1})$$

- **27.** Complete the following reaction equations:
 - (i) C₆H₅N₂Cl + CH₃COCl
 - (ii) C₂H₅NH₂ + C₆H₅SO₂Cl
 - (iii) C₂H₅NH₂ + HNO₂
- **28.** (a) Give chemical tests to distinguish between compounds in the following pairs of substances:
 - (i) Ethanal and Propanal
 - (ii) Benzoic acid and Ethyl benzoate
 - (b) An organic compound contains 69.77% carbon, 11.63% hydrogen and rest oxygen. The molecular mass of the compound is 86. It does not reduce Tollen's reagent but forms an addition compound with sodium hydrogensulphite and gives positive iodoform test. On vigorous addition, it gives ethanoic and propanoic acids. Derive the structure of the compound 'A'. 5

- (a) Arrange the following compounds in an increasing order of their indicated property:
 - (i) Benzoic acid, 4-Nitrobenzoic acid, 3,4-Dinitrobenzoic acid, 4-Methoxybenzoic acid (acid strength)
 - (ii) CH₃CH₂CH(Br)COOH, CH₃CH(Br) CH₂COOH, (CH₃)₂CHCOOH, CH₃CH₂COOH (acid strength)

74 | Xam idea Chemistry–XII

	(b) How would you bring about the following conversions:
	(i) Propanone to Propene
	(ii) Benzoic acid to Benzaldehyde
	(iii) Bromobenzene to 1-phenylethanol
29.	(a) Draw the structure of the following:
	(i) H ₃ PO ₂
	(ii) BrF ₃
	(b) How would you account for the following observations:
	(i) Phosphorus has a greater tendency for catenation than nitrogen.
	(ii) Bond dissociation energy of fluorine is less than that of chlorine.
	(iii) No chemical compound of helium is known. 5
	OR
	(a) Draw the structures of the following:
	$(i) N_2O_5$
	(ii) XeOF ₄
	(b) Explain the following observations:
	(i) The electron gain enthalpy of sulphur atom has a greater negative value than that of
	oxygen atom.
	(ii) Nitrogen does not form pentahalides.
	(iii) In aqueous solutions HI is a stronger acid than HCl.
CBS	E (All India) SET-III
[Duo	stions different from Set–I and Set–II]
ĮŲuG	Stivils unferent from Set-1 and Set-11]
1.	What is the number of atoms in a unit cell of a face-centred cubic crystal?
13.	Describe the role of the following:
	(i) NaCN in the extraction of silver from a silver ore.
	(ii) Cryolite in the extraction of aluminium from pure alumina.
14.	Define the following:
	(i) Order of a reaction
	(ii) Activation energy of a reaction
17.	Differentiate between condensation and addition polymerisations. Give one example each of the
17.	resulting polymers.
21.	A voltaic cell is set up 25° C with the following half-cells:
21.	Al Al ³⁺ (0.0010 M) and Ni Ni ²⁺ (0.50 M).
	Write the equation for the cell reaction that occurs when the cell generates an electric current and determine the cell potential.
	(Given: $E_{Ni^2/Ni}^0 = 0.25V, E_{Al^3/Al}^0 = 1.66V$)

- **23.** Explain the following:
 - (i) Low spin octahedral complexes of nickel are not known.
 - (ii) The complexes are known for transition elements only.
 - (iii) CO is a stronger ligand than NH₃ for many metals.

OR

Compare the following complexes with respect to structural shapes of units, magnetic behaviour and hybrid orbitals involved in units:

- (i) $[Ni(CN)_4]^2$
- $(ii) [NiCl_4]^2$
- (iii) [CoFe₆]³

[At. Nos,: Ni = 28; Co = 27]

- 27. What are the following substances? Give one example of each of them.
 - (i) Cationic detergents
 - (ii) Enzymes
 - (iii) Sweetening agents
- **30.** (a) Draw the structures of the following:
 - (i) XeF₄
 - (ii) H₂S₂O₇
 - (b) Explain the following observations:
 - (i) Phosphorus has a greater tendency for catenation than nitrogen.
 - (ii) The negative value of electron gain enthalpy is less for fluorine than that for chlorine.
 - (iii) Hydrogen fluoride has a much higher boiling point than hydrogen chloride.

OR

- (a) Draw the structures of the following:
 - (*i*) PCl₅(*s*)
 - (ii) SO_3^2
- (b) Explain the following observations:
 - (i) Ammonia has a higher boiling point than phosphine.
 - (ii) Helium does not form any chemical compound.
 - (iii) Bi(V) is a stronger oxidising agent than Sb(V).

3

3

5

SOLUTIONS

CBSE (All India) SET-I

- Metallic substances conduct electricity through electrons while ionic substances conduct electricity in molten state or in solution through ions.
- 2. The process of settling of colloidal particles is called coagulation.
- 3. It is a thermal process of extracting a metal from its ore.
- **4.** This is due to polymeric structure of red phosphorus or angular strain in P_4 molecule of white phosphorus where the angle is only 60° .

5.
$$H_2C = CH - CH_2 - CH_2 - CH_3$$
 Hex-1-en-3-ol. OH

6. 1-Phenyl pentan-1-one
$$CH_3$$
— CH_2 — CH_2 — CH_2 — C

- 7. $NH_3 < (CH_3)_3N < CH_3 NH_2 < (CH_3)_2 NH$.
- 8. The two monomers (adipic acid and hexamethylene diamine) contain 6 carbon atoms each.
- 9. Lead storage battery is a secondary cell.

At anode:
$$Pb(s)$$
 SO_4^2 (aq) $PbSO_4(s)$ 2e
At cathode: $PbO_2(s)$ SO_4^2 (aq) 4H (aq) 2e $PbSO_4(s)$ 2H₂O(l)
Overall reactions: $Pb(s)$ $PbO_2(s)$ 2H₂SO₄ (aq) 2PbSO₄ (s) 2H₂O(l)

OR

At cathode:
$$MnO_4(aq)$$
 8H 5e $Mn^{2+}(aq)$ 4H₂O(l)] 2 E 1.51 V At anode: Sn^2 Sn^4 (aq) 2 e] 5 E° = +0.15 V Overall reaction: $2MnO_4(aq)$ $5Sn^2$ (aq) $16H^+(aq)$ $2Mn^{2+}(aq)$ $5Sn^{4+}(aq)$ $8H_2O(l)$ $E^o_{Sn^4/Sn^2}$ $E^o_{Sn^2/Sn^4}$ 0.15 V E^o_{cell} $E^o_{cathode}$ E^o_{anode} $E^o_{MnO_4/Mn^2}$ $E^o_{Sn^4/Sn^2}$ = 1.51 – (–0.15) E^o_{Cell} 1.66V

As $E_{\text{cell}}^{\text{o}}$ is +ve therefore the reaction will take place in forward direction, *i.e.*, favours the formation of products.

- 10. (i) Elementary step: Each step of a complex reaction is called an elementary step.
 - (ii) Rate of reaction: It is the change in the concentration of any of the reactants or products per unit time.
- 11. (i) Electrolytic refining: In this method, the impure metal is made to act as anode. A strip of same metal in pure form is used as cathode. They are put in a suitable electrolytic bath containing soluble salt of same metal. When electric current is passed, the metal from the anode goes into solution as ions due to oxidation while pure metal gets deposited at the cathode due to reduction of metal ions. The less electropositive impurities settle down below the anode as anode mud.

At anode: M M^n ne At cathode: M^n ne M

(ii) Vapour phase refining of metals: In this method, crude metal is freed from impurities by first converting it into volatile compound and collected elsewhere. It is then decomposed to give pure metal. For example refining of nickel by Mond process.

 $\begin{array}{ccccc} Ni & 4CO & ^{330} & ^{350K} & Ni(CO)_4 \\ impure & & & Tetracarbonyl nickel (volatile) \\ Ni(CO)_4 & ^{450} & ^{470K} & Ni & ^{4CO} \\ & & Pure & \end{array}$

- **12.** (*i*) $2XeF_2 + 2H_2O$ $2Xe + 4HF + O_2$
 - $\begin{array}{ccc} \textit{(ii)} & 2\text{PH}_3 + & 3\text{HgCl}_2 & & \text{Hg}_3\text{P}_2 & 6\text{HCl} \\ & & \text{Mercuric chloride} & & \text{Mercuric phosphide} \end{array}$
- 13. (i) MnO_4 8H 5e Mn^2 4H₂O] 2 $\frac{C_2O_4^2}{2MnO_4^- + 5C_2O_4^2}$ 16H $2Mn^2$ 10CO₂ 8H₂O

- 14. (i) Cl 3° halide reacts faster than 2° halide because of the greater stability of tertiary carbocation.

 tert-Butyl chloride
 - (ii) 2° halide reacts faster than 1° halide because of the greater stability of secondary carbocation than primary.

(ii)
$$\text{CH}_3$$
— CH_2 — $\text{CH}=\text{CH}_2+\text{H}$ — Br CH_3 — CH_2 — CH — CH_3 — Br
2 Bromo butane

16. The four bases present in DNA are adenine (A), guanine (G), cytosine(C) and thymine (T). Thymine(T) is not present in RNA.

17.

Fat soluble vitamins	Sources	Deficiency diseases	
Vitamin A	Fish liver oil, carrots, milk	Xerophthalmia, Night blindness.	
Vitamin D	Exposure to sunlight, fish and egg.	Rickets and osteomalacia.	
Vitamin E	Vegetable oils like wheat germ oil, sunflower oil, etc.	Sterility and muscular atrophy	
Vitamin K	Green leafy vegetables	Haemorrhages and increased blood clotting time.	

18.

	Thermoplastics	Thermosetting plastics		
(<i>i</i>)	These polymer are linear or slightly branched chain molecules		These polymers are cross linked or heavily branched molecules	
(ii)	(ii) Soften on heating and harden on cooling and can be remoulded.		On heating undergo extensive cross linking in moulds and become infusible.	
(iii)	Some common examples are polyethene, PVC, polystyrene, etc.	()	Some common examples are bakelite, urea-formaldehyde resins, terylene, etc.	

19. For a first order reaction

$$t = \frac{2.303}{k} \log \frac{[R]_o}{[R]}$$

Here

t = 3 h = 3 × 60 min = 180 min
k = 0.0051min⁻¹, [R]_o = 0.10 M, [R] = ?
180 min =
$$\frac{2.303}{0.0051 \text{ min}} \log \frac{0.10}{[R]}$$

Log $\frac{0.1}{[R]} = \frac{180 \text{ min}}{2.303} \log \frac{0.051 \text{ min}}{2.303} = \frac{918}{2303}$
log $\frac{0.1}{R} = 0.3986$
 $\frac{0.1}{R} = \text{Anti log } (0.3986) = 2.503$
R = $\frac{0.1}{2.503} = 0.03995 \text{ M}$
R = 0.04M

$$r = \frac{a}{2\sqrt{2}}$$

Given a = 409 pm

$$r = \frac{409}{2\sqrt{2}} \quad \frac{409\sqrt{2}}{4}$$

$$r = 144.58 \text{ pm}$$

 Cu^{2} (aq) $2e^{-}$ **21.** At anode: Cu(s)

> At cathode: 2Ag (aq) e 2Ag(s)

 Cu^2 (aq) 2Ag(s)Overall reaction: Cu(s) 2Ag (aq)

$$n=2$$
,

[Ag] ?

Given, E_{cell} 4.22 V

 $[Cu^2] 0.1 M$

$$E^{o}_{Ag\ /Ag} = 0.80 \text{ V}$$

$$E_{Cu^2}^{o} / Cu = 0.34 \text{ V}$$

$$E_{cell}^{o} = E_{cathode}^{o}$$
 E_{anode}^{o} E_{Ag}^{o} Ag $E_{Cu^{2}}^{o}$ /Cu

$$E_{cell}^{o} = 0.80 - 0.34 = 0.46 \text{ V}$$

$$E_{cell} = E_{cell}^{o} \quad \frac{0.0591}{n} \log \frac{[Cu^2]}{[Ag]}$$

$$0.422 = 0.46 - \frac{0.0591}{2} \log \frac{0.1}{[Ag]^2}$$

$$0.422 - 0.46 = -0.0295 \log \frac{10^{-1}}{[Ag]^2}$$

$$-0.038 = -0.0295 \log \frac{10^{-1}}{[Ag]^2}$$

$$\frac{0.038}{0.0295}$$
 $\log 10^{-1}$ $\log [Ag]^2$

$$1.2881 = -\log 10 - 2\log [Ag^{+}]$$

$$1.2881 = -1 - 2 \log [Ag^{+}]$$

$$2 \log [Ag^+] = 2.2881$$

$$log [Ag^+] = -1.144$$

$$[Ag^+]$$
 = Antilog $\overline{2}.856$

$$[Ag^+] = 7.178 \times 10^{-2} M$$

- 22. (i) The positively charged colloidal particles of Fe(OH)₃ get coagulated by the negatively charged ions provided by electrolyte.
 - (ii) The path of light becomes visible due to scattering of light by colloidal particles (Tyndall effect).
 - (iii) Electrophoresis takes place in which colloidal particles move towards the oppositely charged electrode where they lose their charge and get coagulated.
- **23.** (*i*) **Crystal field spliting:** When the ligands approach the central metal ion, the electrons in the d-orbitals of central metal ion will be repelled by the lone pairs of the ligands. Because of these interactions the degeneracy of d orbitals of the metal ion is lost and these split into two sets of orbitals having different energies. This is known as crystal field splitting, *e.g.*, for d⁴, configuration is $t^3_{2g} e^1_g$ in the presence of weak field ligand.
 - (ii) Linkage isomerism: The isomers which have same molecular formula but differ in the linkage of ligand atom to the central metal atom are called linkage isomers, e.g.,

Pentaamminenitrito-N-Cobalt (III) chloride, pentaamminenitrito-O-Cobalt(III) chloride

(iii) Ambidentate ligand: A ligand which can bind to the central metal atom through any of the two donor atoms present in it is called ambidentate ligand, e.g., NO₂ can bind to metal either through nitrogen, i.e., as nitrito-N (\(\bigcup N = O \)) or through oxygen atom, i.e., as nitrito -O (\(O - N = O \))

Complex/Ion	Central metal ion/atom	Configuration of metal ion	Hybri- disation of metal ion involved	Geometry of the complex	Number of unpaired electrons	Magnetic behaviour
$[Co(NH_3)_6]^{3+}$	Co ³⁺	$d^{6}4s^{0}$	d^2sp^3	octahedral	0	Diamagnetic
$[Cr(NH_3)_6]^{3+}$	Cr ³⁺	$3d^34s^0$	d^2sp^3	octahedral	3	Paramagnetic
[Ni(CO) ₄]	Ni	$3d^34s^2$	sp^3	Tetrahedral	0	Diamagnetic

- 24. (i) Ethanol undergoes intermolecular hydrogen bonding due to the presence of a hydrogen attached to oxygen atom. As a result, ethanol exist as associated molecules and hence it has higher boiling point than methoxy ethane which does not form hydrogen bonds.
 - (ii) Phenol is stronger acid than ethanol because the phenoxide ion left after the release of proton is stabilized by resonance but ethoxide ion is not. Moreover, ethoxide ion is destabilised by +I effect of ethyl group.
 - (iii) Due to –I effect or –R effect of –NO₂ group, the resulting phenolate ion is more stable than phenoxide ion. Therefore o- and p--nitrophenols are more acidic than phenol.
- **25.** (*i*) The catalytic activity of transition metals is attributed to the following reasons:
 - (a) Because of their variable oxidation states transition metals form unstable intermediate compounds and provide a new path with lower activation energy for the reaction.

- (b) In some cases, the transition metal provides a suitable large surface area with free vacancies on which reactants are adsorbed.
- (ii) This due to filling of 4f orbitals which have poor shielding effect or due to lanthanoid contraction.
- (iii) This is due to comparable energies of 5f, 6d and 7s orbital in actinoids.

26. (i)
$$R - C - NH_2$$

$$\begin{array}{c|c}
 & \text{LiAIH}_4 \\
 & \text{H}_2O
\end{array}
R - CH_2 - NH_2$$
(ii) $\begin{array}{c}
 & \stackrel{\uparrow}{N_2}C\overline{l} \\
 & \text{Benzene} \\
 & \text{diazonium chloride}
\end{array}$

$$\begin{array}{c|c}
 & \text{NH}_2 \\
 & \text{Aniline}
\end{array}$$

2, 4, 6 - Tribromo aniline

- (i) Non-ionic detergents: These are the esters of high molecular mass alcohols with fatty acids. 27. These are named so because they do not contain any ion in their constitution, e.g., polyethylene glycol stearate.
 - (ii) Food preservatives: These are the substances which prevent spoilage of food due to microbial growth, e.g., sodium benzoate, potassium metabisulphite, salts of sorbic acid and propanoic acid, etc.
 - (iii) Disinfectants: These are the chemical substances which kill micro-organisms or stop their growth but are harmful to human tissues, e.g., phenol (1%), chlorine in concentration of 0.2 to 0.4 pm in aqueous solution, SO_2 , etc.
- **28.** (a) (i) Mole fraction: It may be defined as the ratio of number of moles of one component to the total number of moles of all the components present in the solution.
 - (ii) Van't Hoff factor (i): It may be defined as the ratio of normal molar mass to the observed molar mass, i.e., Van't Hoff factor
 - Normal molar mass Observed molar mass

OR

Observed colligative property Calculated colligative property

$$(b) = \frac{W_B - R - T}{M_B - V}$$

$$M_B = \frac{W_B - R - T}{V}$$


Here

$$\begin{split} W_B &= \ 100 \ \text{mg} = 100 \times 10^{-3} \ \text{g} \\ R &= \ 0.0821 \ \text{L} \ \text{atm} \ \text{K}^{-1} \ \text{mol}^{-1} \\ T &= \ 25^{\circ}\text{C} = (25 + 273) \ \text{K} = 298 \text{K} \\ V &= \ 100 \ \text{mL} = 10 \times 10^{-3} \ \text{L} \\ &= \ 13.3 \ \text{mm} \ \text{Hg} \ \frac{13.3}{760} \ \text{atm} \\ M_B &= \frac{100 \ 10^{-3} \ \text{g} \quad 0.0821 \ \text{L} \ \text{atm} \ \text{K}^{-1} \ \text{mol}^{-1} \quad 298 \text{K}}{\frac{13.3}{760} \ \text{atm} \quad 100^{-10}^{-3} \ \text{L}} \\ M_B &= \frac{100 \ 10^{-3} \ 0.0821 \ 298 \ 760 \ \text{g} \ \text{mol}^{-1}}{133 \ 10^{-10}^{-3}} = \frac{18594 \ \text{g} \ \text{mol}^{-1}}{133} \end{split}$$


- (i) Colligative properties: Those properties which depend on the number of solute particles irrespective of their nature relative to the total number of particles present in the solution are called colligative properties of solutions.
- (ii) Molality (m): It is the number of moles of the solute per kilogram of the solvent and is expressed as

(b)
$$P_{N_2} = 0.78 \text{ atm} = 0.78 \times 760 \text{ mm Hg}$$

 $= 592.8 \text{ mm Hg}$
 $K_H = 8.42 \times 10^{-7} \text{ M/mmHg}$
 $X_{N_2} = ?$
 $X_{N_2} = K_H \times P_{N_2}$
 $= 8.42 \times 10^{-7} \text{ M/mmHg} \times 592.8 \text{ mmHg} = 4991.376 \times 10^{-7}$
 $X_{N_2} = 4.99 \times 10^{-4}$
 $X_{N_2} = \frac{X_{N_2}}{X_{N_2}} \frac{X_{N_2}}{X_{H_2O}} \frac{X_{N_2}}{X_{H_2O}}$
 $X_{N_2} = X_{H_2O} \times X_{N_2} \frac{1000}{18} 4.99 \cdot 10^{-4} \cdot 0.0277$
 $X_{N_2} = 2.77 \times 10^{-2} = 2.77 \times 10^{-2} \text{M}$




Peroxodisulphuric acid $(H_2S_2O_8)$



Perchloric acid $(HClO_4)$

- (b) (i) This is because the lone pair of electrons on N atom in NH_3 is directed and not delocalised as it in PH_3 due to larger size of P.
 - (ii) This is because S-S single bond is stronger O—O single bond.
 - (iii) Due to small size of fluorine atom there are strong interelectric repulsions in the relatively small size 2p orbitals of fluorine and thus, the incoming electron does not experience much attraction.



- (i) Due to resonance, N—O bond length is the average of single and double bond whereas N—OH bond is purely single bond.
- (ii) PCl₅ has trigonal bipyramidal structure in which the three equatorial P—Cl bonds are equivalent, while the two axial bonds are longer than equational bonds. This is due to the fact that axial bond pairs suffer more repulsion as compared to equatorial bond pairs.
- (iii) This is because I—Cl bond has lower bond dissociation enthalpy than Cl—Cl bond.

30. (a) (i) Cannizzaro Reaction:

(ii) Hell-Volhald-Zelinsky reaction:

(b) (i) Propanal and Propanone:

Tollen's reagent test: Propanal being an aldehyde reduces Tollens reagent to silver mirror but propanone being a ketone does not.

(ii) Benzophenone and Acetophenone:

Iodoform test: Acetophenone being a methyl ketone on treatment with NaOI (I₂/NaOH) gives yellow ppt of iodoform but benzophenone does not.

(iii) Phenol and Benzoic acid:

FeCl₃ test: Phenol gives a violet colouration with neutral FeCl₃ solution while benzoic acid gives buff coloured ppt ferric benzoate.

(ii) O OMgBr
$$H$$
—C—CH₃

Benzal dehyde

OMgBr H —C—CH₃
 H_2O/H^+
 $K_2Cr_2O_7/H_2SO_4$
(O)

O
C—CH₃

O
C—CH₃

(b) A = Butyl butanoate
$$B = Butanoic acid$$

C = Butanol $D = But-1$ -ene

Reactions involved:

$$\begin{array}{c} \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{3} & \text{H}_{2}\text{O} \\ \text{Butylbutanoate(A)} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{C} - \text{OH} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{OH} \\ \text{Butanoic acid(B)} & \text{Butanol(C)} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{OH} \\ \text{Butanol(C)} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CHO} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{COOH} \\ \text{Butanolic acid (B)} \\ \text{CH}_{3} - \text{CH}_{2} \\ \text{Butan-I-ol} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{Butan-I-ene.} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{Butan-I-ene.} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{Butan-I-ene.} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{Butan-I-ene.} & \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2} \\ \text{CH}_{3} - \text{CH}_{2} - \text{CH}_{2}$$

CBSE (All India) SET-II

- 1. Schottky defect.
- **8.** The monomers must be bifunctional.

19. Here,
$$z = 2$$
, $M = 56$ g mol⁻¹, $d = 7.87$ g cm⁻³

$$a = 286.65 \text{ pm} = 286.65 \times 10^{-10} \text{ cm} = 2.8$$
Substituting the values in the expression,

$$N_A = \frac{z}{a^3} \frac{M}{d}, \text{ we get}$$

$$N_A = \frac{2 \cdot 56 \text{ g mol}^{-1}}{(286.65 \cdot 10^{-10} \text{ cm})^3 \cdot 7.87 \text{ g cm}^{-3}}$$

$$= 6.042 \times 10^{23} \text{ mol}^{-1}$$

$$20. \quad \log \frac{k_2}{k_1} \quad \frac{E_a}{2303R} \quad \frac{T_2 \cdot T_1}{T_1 \cdot T_2}$$

20.
$$\log \frac{x_2}{k_1} = \frac{-a}{2303R} = \frac{x_2 - x_1}{T_1 T_2}$$

$$E_a = \frac{2303 R T_1 T_2}{T_2 - T_1} \log \frac{k_2}{k_1}$$

$$E_a = \frac{2303 - 8314 \mathrm{J \ mol}^{-1} \mathrm{K}^{-1} - 650 \, \mathrm{K}}{700 \, \mathrm{K} - 650 \, \mathrm{K}} \log \frac{239 - 10^{-7}}{215 - 10^{-8}}$$

$$E_a = \frac{19.147 - 650 - 700}{50} \log (23.9 - \log 2.15) \text{J mol}^{-1}$$

$$E_a = 174237.7(1.3783 - 0.3324) \text{J mol}^{-1}$$

$$E_a$$
 174237.7 1.0459 J mol⁻¹ = 182235.2 J mol⁻¹

$$E_a$$
 182.24 kJ mol⁻¹.

(ii)
$$C_2H_5NH_2$$
 $C_6H_5SO_2Cl$ $C_6H_5SO_2NHC_2H_5$

(iii)
$$C_2H_5$$
 — NH_2 HNO₂ C_2H_5 — OH H₂O N₂

$$C_2H_5$$
 — OH H_2O N_2

(i) Ethanol, when warmed with NaOI (I2/NaOH) gives yellow ppt of Iodoform while propanal 28. does not

(ii) Benzoic acid being an acid, decomposes NaHCO3 to produce brisk effervescence due to evolution of CO₂ while ethyl benzoate does not.

(*b*)

Element	Percentage	Atomic mass	No. of moles	Simplest molar ratio
С	69.77	12	$\frac{69\ 77}{12}$ 5 81	$\frac{581}{116}$ 5
Н	11.63	1	$\frac{11 \ 63}{1} \ 11 \ 63$	$\frac{11 \ 63}{1 \ 16} \ 10$
О	(100 - 81.4) = 18.60	16	$\frac{18 \ 60}{16} \ 116$	$\frac{1\ 16}{1\ 16}$ 1

Empirical formula of the compound $A = C_5 H_{10} O$

Molecular formula of the compound A = n (Empirical formula)

$$n = \frac{\text{Molecular mass of compound } A}{\text{Empirical formula mass of compound } A}$$

Molecular mass of compound A = 86

Empirical formula mass of compound $A = 5 \ 12 \ 1 \ 10 \ 1 \ 16$

$$= 60 + 10 + 16$$

$$= 86$$

$$n = \frac{86}{86} \quad 1$$

Molecular formula of the compound $A = 1(C_5H_{10}O)$

$$= C_5 H_{10} O$$

As the compound A forms addition compound with NaHSO3 therefore it must be either an aldehyde or ketone.

As it does not reduce Tollens reagent and give positive Iodoform test therefore it must be a methyl ketone.

As on oxidation the compound A gives a mixture of ethanoic acid and propanoic acid, therefore compound A is

The chemical reactions are:

$$\begin{array}{c} O \\ || \\ CH_3 - C - CH_2 - CH_2 - CH_3 - CH_3 \end{array} \quad \text{NaHSO}_3 \qquad \begin{array}{c} OH \\ | \\ CH_3 - C - CH_2 - CH_2 - CH_3 - CH_3 - CH_2 - CH_3 - CH_3$$

Sodium hydrogen sulphite addition product

$$\begin{array}{c} O \\ || \\ CH_3-C-CH_2-CH_2-CH_3-3I_2-4NaOH \\ \hline \\ Pentan 2 one \\ \hline \\ Iodoform \\ reaction \\ \hline \\ (Iodoform yellow ppt) \\ \end{array} CH_3-CH_2-CH_2-COONa 3NaI 3H_2O \\ \\ \end{array}$$

$$\begin{array}{c} O \\ || \\ CH_3-C-CH_2-CH_2-CH_3 \\ \text{Pentan 2 one} \end{array} \\ \begin{array}{c} K_2Cr_2O_7 \\ H_2SO_4 \end{array} \\ \begin{array}{c} CH_3-COOH \\ \text{Ethanoic acid} \end{array} \\ \begin{array}{c} CH_3-CH_2-COOH \\ \text{Propanoic acid} \end{array}$$

- (a) (i) Acid strength: 4-Methoxy benzoic acid <Benzoic acid < 4-Nitrobenzoic acid < 3, 4-Dinitrobenzoic acid.
 - (ii) Acid strength: $(CH_3)_2CHCOOH < CH_3CH_2COOH < CH_3CH(Br)CH_2COOH$ < CH₃CH₂CH(Br)COOH

- (*b*) (i) Due to greater bond strength of P—P single than N—N single bond.
 - (ii) Due to large electron-electron repulsion among the lone pairs in F₂ as fluorine has very small size.

BrF3: Bent "T"

(iii) Due to very high ionization enthalpy of helium.

(a) (i)
$$\overset{\bullet}{\mathbf{O}}$$
 $\overset{\bullet}{\mathbf{O}}$ $\overset{\bullet}{\mathbf{O$

- (b) (i) Due to small size of oxygen atom there will be greater interelectronic repulsions in oxygen.
 - (ii) Due to non-availability of d orbitals in valence shell nitrogen does not form pentahalide.
 - (iii) Due to low bond dissociation enthalpy of H-I as compared to H—Cl.

CBSE (All India) SET-III

- 1. (8 corner atoms) $\frac{1}{8}$ (6 face centre atoms) $\frac{1}{2} = 1 + 3 = 4$
- **13.** (*i*) Dilute NaCN forms a soluble complex with Ag or Ag₂S while the impurities remain unaffected which are filtered off.

- (ii) The role of cryolite is two fold.
 - It lowers the melting point of the mixture to about 1140K.
 - It increase the electrical conductivity of the mixture.
- **14.** (*i*) Order of a reaction: It may be defined as the sum of exponents of the concentration terms in the rate law expression.
 - (ii) Activation energy of a reaction: It may be defined as the extra amount of energy over and above the average energy of reactants which must be supplied to them to undergo a chemical reaction.

17.

	Addition Polymerisation		Condensation Polymerisation
(<i>i</i>)	Monomers are unsaturated molecules	(<i>i</i>)	Monomers have di or polyfunctional groups.
(ii)	Involves chain reaction	(ii)	Does not involve chain reaction.
(iii)	Formed by adding monomers to a growing chain without loss of any molecules.	(iii)	Monomers combine together with the loss of molecules like H_2O , NH_3 , etc.
	Examples: Polyethene, Polystyrene, Teflon etc.		Examples: Terylene, Bakelite, Nylon-66, etc.

 $E_{cell} = 1.46 \text{ V}$

21.

At anode: Al(s) Al³ (aq)
$$3\overline{e}$$
] 2

At cathode: Ni² (aq) $2\overline{e}$ Ni(s)] 3

$$2Al(s) \ 3Ni^2 \ (aq) \ 2Al^3 \ (aq) \ Ni(s)$$

$$E_{cell} = E_{cell}^0 \frac{0.0591}{n} \log \frac{[Al^3]^2}{[Ni^2]^3}$$
Here $n = 6$, $[Al^{3+}] = 0.001M = 1 \times 10^{-3}M$, $[Ni^{2+}] = 0.5M$

$$E_{cell}^0 = E^0_{Ni^2} /_{Ni} E^0_{Al^3/Al} \quad 0.25V \quad 1.66V$$

$$E_{cell}^0 = 1.41V$$

$$E_{cell}^0 = 1.41 \frac{0.0591}{6} \log \frac{10^{-3}}{0.5^{-3}} \quad 1.41 \frac{0.0591}{6} \log \frac{10^{-6}}{0.125}$$

$$= 1.41 \frac{0.0591}{6} \log 10^{-6} \quad 8 \quad 1.41 - \frac{0.0591}{6} \log 10^{-6} \quad \log 2^3$$

$$= 1.41 \frac{0.0591}{6} \log - 6 \log 10 \quad 3 \log 2 \quad 1.41 - \frac{0.0591}{6} - 6 \quad 3 \quad 0.3016$$

$$= 1.41 \frac{0.0591}{6} (5.097) = 1.41 \frac{0.3012}{6}$$

$$= 1.41 + 0.0502 = 1.4602V$$


- 23. (i) Ni in its atomic ionic state can not afford two vacant 3d orbitals hence d^2sp^3 hybridisation is not possible.
 - (ii) Transition metals have vacant d orbitals in their atoms or ions into which the electron pairs can be donated by ligands containing electrons, e.g., C_6H_6 , $CH_2 = CH_2$, etc. Thus d P bonding is possible.
 - (iii) Because in case of CO back bonding takes place in which the central metal uses its filled d orbital with empty anti bonding * molecular orbital of CO.


OR

Complex ion	Central metal ion	Configuration of metal ion	Hybridisation of metal ion involved	Geometry of complex ion	Number of unpaired electrons	Magnetic behaviour
$[Ni(CN)_4]^2$	Ni ²⁺	d^8	dsp^2	Square planar	0	Diamagnetic
[Ni(Cl) ₄] ²⁻	Ni ²⁺	d^8	sp^3	Tetrahedral	2	Paramagnetic
[CoF ₆] ³⁻	Co ³⁺	d ⁶	sp^3d^2	Octahedral	4	Paramagnetic

(ii) Enzymes: Enzymes are globular proteins with high molecular mass ranging from 15,000 to 1,000,000 g mol⁻¹, and form colloidal solution in water. A number of reactions that occur in the body of animals and plants to maintain the life process are catalysed by enzymes therefore enzymes are termed as biochemical catalysts.

(b) Sweetening agents: These are non-nutritive substances which are used as substitute of sugar in food and beverages, e.g., ortho-sulphobenzimide (saccharin), sucrolose, aspartame, etc.






(b) (i) This is because P-P single bond is stronger than the N-N single bond.

(ii) Due to extremely small size of F atom, interelectronic repulsions are more in fluorine.

(iii) There is H-bonding in HF molecules due to high electronegativity and small size of fluorine atom but there is no H-bonding in HCl.



(b) (i) Due to small size and high electronegativity of nitrogen, molecules of ammonia are highly associated through hydrogen bonding.

(ii) This is due to small size and high ionization enthalpy of helium.

(iii) As Bi(V) is more stable than Sb(V) due to inert pair effect.

CBSE EXAMINATION PAPERS

FOREIGN-2009

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Foreign) SET-I

- 1. What is the number of atoms in a body-centred cubic unit cell of a crystal?
- 2. What is an emulsion?
- 3. Which one has higher electron gain enthalpy with negative sign, sulphur or oxygen?
- 4. Give the IUPAC name of the following compound:

- **5.** Write the structural formula of 3-oxopentanal.
- **6.** Name two metals which occur in nature as oxides.
- 7. Arrange the following compounds in an increasing order of their basic strength in aqueous solutions:

$$NH_3$$
, RNH_2 , R_2NH , R_3N

1

1

1

1

2

- **8.** Write the name of an antacid which is often used as a medicine.
- Differentiate between molality and molarity of a solution. What is the effect of rise in temperature on molality and molarity of the solution?
- **10.** Describe the role of the following:
 - (i) NaCN in the extraction of silver
 - (ii) CO in the purification of nickel
- 11. Corrosion is essentially an electrochemical phenomenon. With the help of a diagram explain the reactions occurring during the corrosion of iron kept in open atmosphere.

OR

Define the term molar conductivity and indicate how molar conductivity of a substance changes with change in concentration of a weak electrolyte and a strong electrolyte in its solution.

12. Draw the structures of the following molecules: 2 (*i*) BrF₃ (ii) H₂S₂O₇ **13.** State reasons for the following observations: 2 (i) The enthalpies of atomisation of transition elements are quite high. (ii) There is a greater horizontal similarity in the properties of the transition elements than of the main group elements. **14.** Give a chemical equation for each of the following reactions: 2 (i) Williamson's synthesis (ii) Reimer-Tiemann reaction **15.** Explain the mechanism of each of the following processes: 2 (i) Acid catalysed dehydration of an alcohol (ii) Hydration of ethene to yield ethanol. 16. Name two water soluble vitamins, state their sources and the diseases caused due to their deficiency in diet. 2 **17.** What are the following substances? (i) Invert sugar (ii) Polypeptides **18.** State reasons for the following occurrences: 2 (i) Soaps do not do the cleansing in hard water. (ii) Synthetic detergents are preferred to soaps in washing machines. OR What is the repeating unit in the condensation polymer obtained by combining HO₂CCH₂CH₂CO₂H (succinic acid) and H₂NCH₂CH₂NH₂ (ethylenediamine). 19. Silver crystallises in face-centred cubic unit cells. Each side of the unit cell has a length of 409 pm. What is the radius of silver atom? (Assume that each face atom is touching the four corner atoms in the unit 20. Calculate the equilibrium constant for the reaction equilibrium $Fe(s) + Cd^{2+}(aq) \Longrightarrow Fe^{2+}(aq) + Cd(s)$ $E_{Cd^2/Cd}^{o}$ 0.40V; $E_{Fe^2/Fe}^{o}$ 0.44V Given: 3 21. Calculate the freezing point depression for 0.0711 m aqueous solution of sodium sulphate, if it is completely ionised in solution. If this solution actually freezes at – 0.320 °C, what is the value of Van't Hoff factor for it at the freezing point? (K_f for water is 1.86°C mol⁻¹) 22. Describe what is observed when 3 (i) an electric current is passed through a colloidal solution. (ii) a beam of light is passed through a colloidal solution. (iii) an electrolyte such as NaCl, is added to hydrated ferric oxide sol.

- **23.** Explain the following observations:
 - (i) With the same d-orbital configuration (d^4) Cr^{2+} ion is a reducing agent while Mn^{3+} ion is an oxidising agent.

3

3

3

- (ii) Cu⁺ ion is not stable in aqueous solutions.
- (iii) Among the 3d series of transition elements, the largest number of oxidation states are exhibited by manganese.
- Three geometrical isomers are possible for [Co(en)(H₂O)₂(NH₃)₂]³⁺. Draw molecular structure of these three isomers and indicate which one of them is chiral.
- 25. Complete the equations for the following reactions:

$$(i) \qquad \begin{array}{c} H \\ + HBr \end{array}$$

(ii)
$$CH_3 + HI \longrightarrow$$

- **26.** How are the following conversions carried out:
 - (i) Aniline to nitrobenzene
 - (ii) Ethanamine to N-ethylethanamide
 - (iii) Chloroethane to propan-1-amine

OR

Give one chemical test each to distinguish between the compounds in the following pairs:

- (i) Methylamine and dimethylamine
- (ii) Aniline and benzylamine
- (iii) Ethylamine and aniline
- 27. Differentiate between the modes of formation of an addition polymer and a condensation polymer.Give one example of each of these formations.
- **28.** (a) How are the following obtained:
 - (i) Benzoic acid from ethylbenzene
 - (ii) Benzaldehyde from toluene
 - (b) Complete each of the following reactions by giving the missing reactant, reagent or product:
 - (i) C_6H_5COC1 H_2 Pd-BaSO₄

5

5

OR

- (a) How will you bring about the following conversions:
 - (i) Ethanol to 3-hydroxybutanal
 - (ii) Benzaldehyde to benzophenone
- (b) An organic compound A contains 69.77% carbon, 11.63% hydrogen and the rest oxygen. The molecular mass of the compound is 86. It does not react with Tollen's reagent but forms an addition compound with sodium hydrogen sulphite and gives a positive iodoform test. On vigorous oxidation it gives a mixture of ethanoic and propanoic acids. Derive the structure of compound A.
- **29.** (a) Complete the following reaction equations:
 - (i) XeF₂ PF₅
 - (ii) $Cl_2(g)$ NaOH(aq) (hot and conc.)
 - (b) Explain the following observations:
 - (i) +3 oxidation state becomes more and more stable from As to Bi in the group.
 - (ii) Sulphur in vapour state exhibits paramagnetism.
 - (iii) Fluorine does not exhibit any positive oxidation state.

- (a) Complete the following reaction equations:
 - (i) $PCl_5 + H_2O$ (excess)
 - (ii) $F_2 + H_2O$
- (b) Explain the following observations:
 - (i) No distinct chemical compound of helium is known.
 - (ii) Phosphorus has a greater tendency for catenation than nitrogen.
 - (iii) In solutions of H_2SO_4 in water, the second dissociation constant K_{a_2} , is less than the first dissociation constant K_{a_1} .

96 | Xam idea Chemistry-XII

 $4PH_3(g)$

 $P_4(g) + 6H_2(g)$

It is found that this reaction follows the following rate equation:

30.	(a) A reaction is of second order with respect to a reactant. How is the rate of reaction affected in
	the concentration of this reactant is
	(i) doubled, (ii) reduced to half?
	(b) A first order reaction has a rate constant of 0.0051 min ⁻¹ . If we begin with 0.10 M concentration of the reactant, what will be the concentration of the reactant left after 3 hours?
	OR
	(a) Define the following:
	(i) Rate of reaction. (ii) Elementary step in a reaction
	(b) For a decomposition reaction, the values of rate constant k at two different temperature are given below:
	$k_1 = 2.15 \times 10^{-8} \text{ L mol}^{-1} \text{ s}^{-1} \text{ at } 650 \text{ K}$
	$k_2 = 2.39 \times 10^{-7} \text{ L mol}^{-1} \text{ s}^{-1} \text{ at } 700 \text{ K}$
	Calculate the value of activation energy (E_a) for this reaction. (R = 8.314 J K ⁻¹ mol ⁻¹)
BS	E (Foreign) SET-II
Que	stions different from Set-I]
2.	What is 'reverse osmosis'?
6.	Why is it that sulphide ores are concentrated by the 'froth floatation process'?
8.	What is meant by the 'broad spectrum antibiotics'?
13.	Complete the following chemical equations:
	(i) $C_2O_7^2$ (aq) $C_2O_4^2$ (aq) H (aq)
	(ii) $MnO_4(aq)$ $Fe^2(aq)$ H (aq)
17.	Describe the following types of substances giving one example of each:
	(i) Cationic detergents
	(ii) Food preservatives
19.	Iron has a body-centred cubic unit cell with a cell edge of 286.65 pm. The density of iron is 7.87 g cm^{-3} Use this information to calculate Avogadro's number. (At. mass of Fe = 56 g mol^{-1})
22.	Write three special features of chemisorption which are not found in physisorption.
24.	Compare the following complexes with respect to their molecular shape and magnetic behaviour:
	(i) $[Cr(NH_3)_6]^{3+}$ (ii) $[Fe(CN)_6]^{4-}$ (iii) $[NiCl_4]^{2-}$
	(At. No. $Cr = 24$, $Fe = 26$, $Ni = 28$)
30.	Decomposition of phosphine (PH ₃) at 120°C proceeds according to the equation:

5

Rate = $k[PH_3]$

The half-life of PH_3 is 37.9 s at 120°C.

- (i) How much time will be required for 3/4 of PH₃ to decompose?
- (ii) What fraction of the original amount of PH₃ will remain undecomposed after 1 minute?

OR

Hydrogen peroxide, H_2O_2 (aq) decomposes to $H_2O(l)$ and $O_2(g)$ in a reaction that is of first order in H_2O_2 and has a rate constant, $k = 1.06 \times 10^{-3} \text{ min}^{-1}$.

- (i) How long will it take 15% of a sample of H₂O₂ to decompose?
- (ii) How long will it take 85% of a sample of H_2O_2 to decompose?

CBSE (Foreign) SET-III

[Questions different from Set-I and Set-II]

1.	Which point defect in its crystal unit cells decreases the density of a solid?					
2.	Write the structure of pent-2	-enal.		1		
9.	State the principles on which	h the following operati	ons are based:			
	(i) Zone refining	(ii) Vapour phase refi	ning.	2		
12.	Draw the structures of the fe	ollowing molecules:				
	(i) XeF ₄	$(ii) H_2S_2O_7$		2		
16.	Name two fat soluble vitam diet.	ins. State their sources	and the diseases caused due to their deficiency	in 2		
18.	State what the following are	and how they differ fr	rom each other:			
	(i) a nucleotide, and	(ii) a nucleoside.		2		
24.	Explain the following giving	g an example in each c	ase:			
	(i) Linkage isomerism	(ii) An outer orbital	complex			
	(iii) A bidentate ligand			3		
27.	Draw the structures of the m	onomers of the followi	ng materials:	3		
	(i) PVC	(ii) Teflon	(iii) Neoprene			
30.	(a) Complete the following	ng reaction equations:				
	$(i) SO_2 + MnO_4 +$	H_2O				
	(ii) HgCl ₂ + PH ₃					
	(b) Explain the following	observations:				
	(i) Sulphur has a g	reater tendency for car	tenation than oxygen.			
	(ii) Fluorine is a str	onger oxidising agent	than chlorine.			

(iii) The +5 oxidation state becomes less stable down the group in group 15 of the periodic table.

- (a) Complete the following reaction equations:
 - $(i) P_4 + NaOH + H_2O$
 - (ii) $Cu + HNO_3$ (dilute)
- (b) Explain why
 - (i) H₂O is a liquid while, inspite of a higher molecular mass, H₂S is a gas.
 - (ii) iron dissolves in HCl to form FeCl2 and not FeCl3.
 - (iii) helium is used in diving equipment

SOLUTIONS

CBSE (Foreign) SET-I

- 1. 8 (Corner atoms) $\times \frac{1}{8}$ 1 (body centre atom) $\times 1 = 1 + 1 = 2$
- 2. Emulsion is a colloidal solution in which both the dispersed phase and dispersion medium are liquids e.g. milk, cod liver oil, etc.
- **3.** Sulphur.

- **6.** Aluminium occurs as Al₂O₃.2H₂O Iron occurs as Fe₂O₃
- 7. $NH_3 < R-NH_2 < R_3N < R_2NH$
- **8.** Ranitidine (zantac).
- 9. Molality is the number of moles of solute per thousand grams of solvent whereas molarity is the number of moles of solute dissolved in one litre of solution.

Molality is independent of temperature whereas molarity changes with change in temperature as volume changes with temperature.

(i) Dilute NaCN forms a soluble complex with Ag or Ag₂S while the impurities remain unaffected 10. which are filtered off.

$$4Ag + 8NaCN + O_2 + 2H_2O \qquad 4Na[Ag(CN_2)] + 4NaOH$$
 Or
$$Ag_2S + 4NaCN \qquad 2Na[Ag(CN)_2] \quad Na_2S \\ Sodium \ dicyanoargentate(I) \\ (Soluble \ complex)$$

(ii) Carbon monoxide forms a volatile complex, nickel tetracarbonyl with nickel which on decomposition gives pure nickel.

$$\begin{array}{ccc} \text{Ni} & \text{4CO} & \text{330-350K} & \text{Ni(CO)}_{4} \\ \text{Impure} & & \text{Nickel tetracarbonyl} \\ \text{Ni(CO)}_{4} & \text{450 470K} & \text{Ni} & \text{4CO} \\ & & \text{Pure} & \end{array}$$

11. At anode: Oxidation of Fe atoms takes place

Fe Fe² 2e
$$E_{Fe^2/Fe} = 0.44V$$

At cathode: Reduction of oxygen in the presence of H ions. The H ions are produced by either H_2O or H_2CO_3 (formed by dissolution of CO_2 in moisture)

2H
$$\frac{1}{2}$$
O₂(g) H₂O

Net reaction at cathodic area

2H (aq)
$$\frac{1}{2}$$
O₂ 2e H₂O E _{H /O₂/H₂O 1.23V}

The overall reaction

Fe(s) 2H (aq)
$$\frac{1}{2}$$
O₂(g) Fe² (aq) H₂O(l) E°_{cell} = 1.67V

The ferrous ions are further oxidised by atmospheric oxygen to ferric ions which come out as rust in the form of hydrated ferric oxide (Fe₂O₃.xH₂O).

OR

Molar Conductivity ($_{\mathbf{m}}$): It may be defined as the conductance of a solution containing 1 mole of electrolyte such that the entire solution is placed is between two electrodes one centimeter apart.


$$_{\rm m} = \times v$$
or
 $_{\rm m} = \frac{1000}{M}$

Molar conductivity increases with decrease in concentration or increase in dilution as number of ions as well as mobility of ions increased with dilution.


For strong electrolytes the number of ions do not increase appreciably on dilution and only mobility or ions increases due to decrease in inter-ionic attractions. Therefore $\ _m$ increases a little as shown in graph by a straight line.


For weak electrolyte the number of ions as well

as mobility of ions increases on dilution as shown by curve in the figure.



(F)





Pyrosulphuric acid $(H_2S_2O_7)$

- **13.** (*i*) This is because, transition elements have strong metallic bonds due to presence of large number of unpaired electrons.
 - (ii) This is because in transition elements incoming electron goes into d-orbitals of inner shell whereas in main group elements, the incoming electron goes to outermost shell.
- 14. (i) Williamson's Synthesis:

$$\begin{array}{cccc} CH_3 & CH_3 \\ CH_3 - C - ONa + CH_3 - Br & CH_3 - C - O - CH_3 + NaBr \\ CH_3 & CH_3 & CH_3 \\ Sodium tert-butoxide & Methyl tert-butyl (ether) \\ \end{array}$$

(ii) Reimer-Tiemann reaction:

15. (i)
$$CH_3 - CH_2 - OH$$
 H $CH_2 = CH_2$ H_2O

Mechanism

Step I: Formation of carbocation: It is the slowest step and hence, the rate determining step.

Step II: Formation of protonated alcohols

Step III: Formation of ethylene by elimination of a proton

To drive the equilibrium to the right, ethylene is removed as it is formed.

(ii)
$$> C = C < + H_2O \xrightarrow{H^+} > C - C < Alkene$$

Mechanism

Step I: Protonation of alkene to form carbocation by electrophilic attack of H₃O⁺.

Step II: Nucleophilic attack of water on carbocation.

Step III: Deprotonation to form an alchohol

16. B group vitamin and vitamin C are soluble in water.

	Name of Vitamins	Sources	Deficiency diseases	
(<i>i</i>)	Vitamin B ₁₂	Meat, fish, egg and curd	Pernicious anaemia	
(ii)	Vitamin C	Citrus fruits and amla	Scurvy	

- 17. (i) Invert Sugar: Sucrose is dextrorotatory, on hydrolysis in the presence of HCl or enzyme invertase, it produces a mixture of D–C(+)–glucose and D–(–)–fructose which is laevorotatory called invert sugar.
 - (ii) **Polypeptide:** If more than ten -amino acids are joined together by peptide bond (-CONH-) the polyamide thus formed is called polypeptide.
- **18.** (*i*) This is because the Ca²⁺ and Mg²⁺ ions present in hard water get precipitated as calcium and magnesium soap which being insoluble stick to the clothes as gummy mass.
 - (ii) This is because detergents can be used in hard water as well as in acidic solutions, as sulphuric acid and their calcium and magnesium salts are soluble in water but the fatty acids and their calcium and magnesium salts are insoluble.

19. For fcc unit cell

$$r = \frac{a}{2\sqrt{2}}$$

Given a = 409 pm

$$r = \frac{409}{2\sqrt{2}} \quad \frac{409\sqrt{2}}{4}$$

$$r = 144.58 \text{ pm}$$

20. Fe(s)
$$\operatorname{Cd}^2(aq) \rightleftharpoons \operatorname{Fe}^2(aq) \operatorname{Cd}(s)$$

$$\log k_{\operatorname{c}} \quad n \frac{E \text{ cell}}{0.059}$$

Here,
$$n = 2$$

$$E_{cell} = E_{cathode} - E_{anode}^{\circ}$$

= $E_{Cd^2/Cd}^{\circ} - E_{Fe^2/Fe}^{\circ} = -0.40 - (-0.44)$
 $E_{cell}^{\circ} = 0.04V$

$$\log k_{\rm c} = \frac{2 + 0.04 \text{ V}}{0.059} = \frac{0.08}{0.059}$$

$$\log k_{\rm c} = 1.3536$$

$$k_{\rm c}$$
 = Antilog 1.3536

$$k_c = 22.57$$

21.
$$T_f$$
 [273 15 (0 320 273 15)]K 0 320 K

$$T_f K_f.m$$


=
$$1.86 \text{ K kg mol}^{-1} \times 0.0711 \text{ mol kg}^{-1}$$

$$= 0.132 \text{ K}$$

$$i = \frac{\text{Observed value of}}{\text{Calculated value of}} \quad \frac{\text{T}_f}{\text{T}_f} \quad \frac{0.320 \text{K}}{0.132 \text{K}}$$

104 | Xam idea Chemistry-XII

- 22. (i) Electrophoresis takes place in which colloidal particles move towards the oppositely charged electrode where they lose their charge and get coagulated.
 - (ii) The path of light becomes visible due to scattering of light by colloidal particles (Tyndall
 - (iii) The positively charged colloidal particles of Fe(OH)₃ get coagulated by the negatively charged ions provided by electrolyte.
- (i) ${\rm Cr}^2$ is reducing as its configuration changes from d^4 to d^3 , the latter having a half filled ${\rm t_{2g}}$ configuration. On the other hand, the change from ${\rm Mn}^3$ to ${\rm Mn}^2$ results in the half-filled d^5 23. configuration which has extra stability therefore Mn³ is oxidising.
 - (ii) Because the high hydration enthalpy of Cu² easily compensates the second ionization enthalpy of Cu.
 - (iii) This is because manganese has electronic configuration $[Ar]3d^54s^2$, with five unpaired electrons in 3d orbitals.



(iii)
$$\sim$$
 OH + SOCl₂ \rightarrow Cl + SO₂ + HCl Chlorocylohexane

26. (i)
$$N_2 = \frac{N_2 \overline{Cl}}{273-278K}$$
 $N_2 = \frac{N_2 \overline{BF}_4}{N_2 \overline{BF}_4}$ $NO_2 = \frac{NaNO_2/HCl}{273-278K}$ $NaNO_2 = \frac{NaNO_2}{Cu, \Delta}$ Nitrobenzene

OR

(i) Methylamine on treatment with alcoholic KOH and CHCl₃ gives offensive smell of methyl isocyanide but dimethyl amine does not.

(ii) Aniline on treatment with NaNO₂/HCl (HNO₂) at 0–5°C followed by treatment with an alkaline solution of naphthol gives an orange coloured azodye while benzylamine does not give this test.

$$NH_{2} \xrightarrow{\text{NaNO}_{2}/\text{HCl}} \bigvee N = N \xrightarrow{\text{NCl}} \frac{\text{b-Naphthol}}{\text{dil NaOH}} \bigvee N = N \xrightarrow{\text{Orange dye}} Orange dye$$

(iii) Add Br₂(aq), aniline forms white ppt while ethyl amine does not form such ppt.

$$\begin{array}{c|c} NH_2 & NH_2 \\ \hline \\ +3Br_2(aq) & Br \\ \hline \\ Aniline & Br \\ \end{array}$$

2, 4, 6–Tribromo aniline (white ppt)

77	
41	-

	Addition Polymerisation		Condensation Polymerisation
(<i>i</i>)	Monomers are unsaturated molecules	(<i>i</i>)	Monomers have di or polyfunctional groups.
(ii)	Involves chain reaction	(ii)	Does not involve chain reaction.
(iii)	Formed by adding monomers to a growing chain without loss of any molecules.	(iii)	Monomers combine together with the loss of molecules like H ₂ O, NH ₃ , etc.
	Examples: Polyethene, Polystyrene, Teflon etc.		Examples: Terylene, Bakelite, Nylon-66, etc.

106 | Xam idea Chemistry-XII

28. (*a*) (*i*)

OR

Benzoyl chloride

(ii)
$$\bigcirc$$
 + Ar/R - C - Cl Acetyl chloride Acetyl chloride Acetyl chloride

(iii)
$$\bigcirc$$
 + \bigcirc C - Cl anhyd. AlCl₃ + HCl Benzophenone + HCl

Benzaldehyde

$$(a) \quad (i) \ \operatorname{CH}_3 - \operatorname{CH}_2 - \operatorname{OH} \quad \overset{\operatorname{PCC}}{\underset{\operatorname{Ethanol}}{\operatorname{CH}_3}} - \overset{\operatorname{CHO}}{\underset{\operatorname{Ethanol}}{\operatorname{CH}_0}} \quad \overset{\operatorname{dil} \operatorname{NaOH}}{\underset{\operatorname{Ethanol}}{\operatorname{CH}_3}} \quad \overset{\operatorname{OH}}{\underset{\operatorname{Durphy}}{\operatorname{CH}_1}} - \overset{\operatorname{OH}}{\underset{\operatorname{Durphy}}{\operatorname{CH}_2}} - \overset{\operatorname{CH}}{\underset{\operatorname{Durphy}}{\operatorname{CH}_3}}$$

(ii) O OH OH CH—C₆H₅

$$+ C_6H_5 - MgBr$$
Benzaldehyde
$$(O) K_2Cr_2O_7/H_2SO_4$$

$$C - C_6H_5$$

$$C - C_6H_5$$

$$C - C_6H_5$$

$$C - C_6H_5$$
Benzophenone

(*b*)

Element	Percentage	Atomic mass	No. of moles	Simplest molar ratio
С	69.77	12	$\frac{69\ 77}{12}$ 5 81	$\frac{581}{116}$ 5
Н	11.63	1	$\frac{11 \ 63}{1} \ 11 \ 63$	$\frac{11 \ 63}{1 \ 16} \ 10$
О	(100 - 81.4) = 18.60	16	$\frac{18 \ 60}{16}$ 1 16	$\frac{1}{1}\frac{16}{16}$ 1

Empirical formula of the compound $A = C_5 H_{10} O$

Molecular formula of the compound A = n (Empirical formula)

$$n = \frac{\text{Molecular mass of compound } A}{\text{Empirical formula mass of compound } A}$$

Molecular mass of compound A = 86

Empirical formula mass of compound A = 5 12 1 10 1 16

$$= 60 + 10 + 16$$

$$n = \frac{86}{86}$$
 1

Molecular formula of the compound $A = 1(C_5 H_{10} O)$

$$= C_5 H_{10} O$$

As the compound A forms addition compound with NaHSO₃ therefore it must be either an aldehyde or ketone.

As it does not reduce Tollens reagent and give positive iodoform test therefore it must be a methyl ketone.

As on oxidation the compound A gives a mixture of ethanoic acid and propanoic acid, therefore compound A is

$$CH_{3} - C - CH_{2} - CH_{2} - CH_{3}$$
Pentan-2-one

The chemical reactions are:

$$\begin{array}{c} O \\ || \\ CH_3 - C - CH_2 - CH_2 - CH_3 - CH_3 \\ \text{Pentan-2-one} \end{array} \quad \begin{array}{c} OH \\ | \\ CH_3 - C \\ | \\ \end{array} \quad \begin{array}{c} CH_3 - C \\ | \\ \end{array} \quad \begin{array}{c} CH_2 - CH_2 - CH_2 - CH_3 \\ | \\ \end{array}$$

Sodium hydrogen sulphite addition product

$$\begin{array}{c} O \\ || \\ CH_3 - C - CH_2 - CH_2 - CH_3 \\ Pentan 2 one \end{array} \quad \begin{array}{c} K_2Cr_2O_7 \\ H_2SO_4 \end{array} \quad \begin{array}{c} CH_3 - COOH \\ Ethanoic acid \end{array} \quad \begin{array}{c} CH_3 - CH_2 - COOH \\ Propanoic acid \end{array}$$

- **29.** (a) (i) XeF₂ PF₅ [XeF] [PF₆]
 - (ii) 3 Cl₂(g) 6NaOH(aq) 5NaCl NaClO₃ 3H₂O (Hot and conc.)
 - (b) (i) This is due to inert pair effect.
 - (ii) In vapour state sulphur partly exists as S_2 molecule having two unpaired electrons in the anti bonding orbitals like O_2 and, hence exhibits paramagnetism.
 - (iii) This is because fluorine is the most electronegative element and does not have d orbitals in its valence shell.

OR

(a) (i)
$$PCl_5 + 4H_2O$$
 (excess) $H_3PO_4 + 5HCl$

(ii)
$$2F_2(g) + 2H_2O(l)$$
 $4H^+(aq) + 4F^-(aq) + O_2(g)$

- (b) (i) This is due to small size, high ionisation enthalpy and stable electronic configuration of helium.
 - (ii) This is because P-P single bond is stronger than N-N single bond.
 - (iii) $Ka_2 \ll Ka_1$, because HSO_4 ion has much less tendency to donate a proton to H_2O as compared to H_2SO_4 .
- **30.** (*a*) Let the rate law be, $r_1 = k [A]^2$
 - (i) If [A] is doubled than rate $r_2 = k(2A)^2 = 4k$ [A]² = $4r_1$, i.e., rate becomes 4 times.
 - (ii) If [A] is reduced to half then rate, r_3 $k + \frac{A}{2} = \frac{1}{4}k[A]^2 + \frac{1}{4}r_1$, i.e., rate becomes $\frac{1}{4}$ times
 - (b) For a first order reaction

$$t = \frac{2.303}{k} \log \frac{[R]_o}{[R]}$$

Here,
$$t = 3 h = 3 \times 60 min = 180 min$$

$$k = 0.0051 \text{min}^{-1}$$
, $[R]_0 = 0.10 \text{ M}$, $[R] = ?$

180 min =
$$\frac{2.303}{0.0051 \,\text{min}^{-1}} \log \frac{0.10}{[R]}$$

$$Log \frac{0.1}{[R]} = \frac{180 \min \quad 0.0051 \min^{-1}}{2.303} = \frac{918}{2303}$$

$$\log \frac{0.1}{R} = 0.3986$$

$$\frac{0.1}{R}$$
 = Anti log (0.3986) = 2.503

$$R = \frac{0.1}{2.503} = 0.03995 \text{ M}$$

$$R = 0.04M$$

OR

- (a) (i) Elementary step: Each step of a complex reaction is called an elementary step.
 - (ii) Rate of reaction: It is the change in the concentration of any of the reactants or products per unit time.

(b)
$$\log \frac{k_2}{k_1} = \frac{E_a}{2303R} = \frac{T_2 - T_1}{T_1 - T_2}$$

$$E_a = \frac{2303 R T_1 T_2}{T_2 - T_1} \log \frac{k_2}{k_1}$$

$$E_a = \frac{2303 8.314 \text{J mol}^{-1} \text{K}^{-1} 650 \text{ K}^{-700} \text{K}}{700 \text{K} - 650 \text{ K}} \log \frac{2.39 10^{-7}}{2.15 10^{-8}}$$

$$E_a = \frac{19.147 650 700}{50} \log (23.9 - \log 2.15) \text{J mol}^{-1}$$

$$E_a = 174237.7 (1.3783 - 0.3324) \text{J mol}^{-1}$$

$$E_a = 174237.7 1.0459 \text{ J mol}^{-1} = 182235.2 \text{ J mol}^{-1}$$

$$E_a = 182.24 \text{ kJ mol}^{-1}.$$

CBSE (Foreign) SET-II

- **2. Reverse osmosis:** If the pressure applied on the solution is greater than the osmotic pressure then the solvent molecules start to move from solution into solvent through semipermeable membrane. This process is called reverse osmosis.
- **6.** As only sulphide ore particles are wet by oil while gangue particles are wet by water.
- **8. Broad spectrum antibiotic:** Antibiotics which kill or inhibit a wide range of Gram-positive and Gram-negative bacteria are known as broad spectrum antibiotics, *e.g.*, chloramphenicol, ofloxacin, vancomycin, etc.

13. (i)
$$\operatorname{Cr_2O_7^2}(aq)$$
 14H (aq) 6e⁻ 2Cr³ (aq) 7H₂O
 $\operatorname{C_2O_4^2}(aq)$ 2CO₂ 2e⁻] 3
 $\operatorname{Cr_2O_7^2}(aq)$ 3(C₂O₄² (aq) 14H (aq) 2Cr³ (aq) 6CO₂ 7H₂O
(ii) MnO₄(aq) 8H (aq) 5e⁻ Mn² (aq) 4H₂O
Fe² (aq) Fe³ (aq) e⁻] 5
 $\operatorname{MnO_4}(aq)$ 5Fe² (aq) 8H (aq) Mn² (aq) 5Fe³ (aq) 4H₂O

17. (i) Cationic Detergents: Cationic detergents are quaternary ammonium salts of amines with acetates, chlorides or bromides as anions. Cationic part possess a long hydrocarbon chain and a positive charge on nitrogen atom. Hence, these are called cationic detergents. Cetyltrimethylammonium bromide is a popular cationic detergent and is used in hair conditioners.

$${\rm CH_3}$$
 ${\rm CH_3}$ ${\rm CH_3}$ ${\rm CH_3}$ ${\rm Br}$ ${\rm CH_3}$

Cetyltrimethyl ammonium bromide

Cationic detergents have germicidal properties and are expensive, therefore, these are of limited use.

(ii) Food Preservatives: These are the chemical substances which are added to the food materials to prevent their spoilage due to microbial growth and to retain their nutritive value for long periods.

Preservatives prevent the rancidity of food and inhibit growth or kill the microorganisms.

The most-common preservations used are, sugar table salt, sodium benzoate, Sodium metabisulphite sorbic acid and propanoic acid.

19. $d = 7.87 \text{ g cm}^{-3}$, For bcc, Z = 2, $M = 56 \text{ g mol}^{-1}$ $a = 286.65 \text{pm} = 286.65 \times 10^{-10} \text{cm}$, $N_A = ?$

$$d = \frac{Z - M}{a^3 - N_A}$$

$$7.87g \text{ cm}^{-3} \frac{2 \cdot 56 \text{ g mol}^{-1}}{(286.65 \cdot 10^{-3} \text{ cm})^{3} \cdot \text{N}_{A}}$$

$$N_{A} = \frac{2 \cdot 56 \text{ g mol}^{-1}}{(2.8665 \cdot 10^{-8} \text{ cm})^{3} \cdot 7.87 \text{ g cm}^{-3}}$$

$$= \frac{2 \cdot 56 \text{ g mol}^{-1}}{23.553 \cdot 10^{-24} \text{ cm}^{3} \cdot 7.87 \text{ g cm}^{-3}}$$

$$= \frac{112 \text{ mol}^{-1}}{185.366 \cdot 10^{-24}} = 0.604 \times 10^{24} \text{ mol}^{-1}$$

$$= 6.024 \times 10^{23} \text{ mol}^{-1}.$$

- **22.** (*i*) Chemisorption is caused by chemical bond formation where as physisorption arises due to Van der Waals' forces.
 - (ii) Chemisorption is highly specific in nature whereas physisorption is not specific.
 - (iii) Chemisorption results into unimolecular layer whereas physisorption results into multimolecular layers.
 - (iv) Chemisorption is irreversible in nature whereas physisorption is reversible in nature.
 - (v) Chemisorption first increases with increase in temperature, then decreases while physisorption decreases with increase in temperature.

24.


(i) Orbitals of Cr^{3+} ion


d²sp³ hybridised orbitals of Cr³⁺

$$[\text{Co(NH}_3)_6]^{3+}$$

Shape: Octahedral


Magnetic behaviour: Paramagnetic.





Six pairs of electrons from six NH₃ molecules

112 | Xam idea Chemistry-XII




d²sp³ hybridised orbitals of Fe²⁺

↑

 $[\mathrm{Fe}(\mathrm{CN})_6]^{4-}$





Six pairs of electrons from six CN⁻ions


Shape: Octahedral


Magnetic behaviour: Diamagnetic

(iii) Orbitals of Ni²⁺ ions

sp³ hybridised orbitals of Ni²⁺

[Ni(Cl)₄]²⁻





Four pairs of electrons from four Cl¯ions

Shape: Tetrahedral

Magnetic behaviour: Paramagnetic

30. (*i*) Given
$$t_{1/2} = 37.95$$

$$t_{1/2} = \frac{0.693}{k}$$
, $k = \frac{0.693}{t_{1/2}} = 1.83 \times 10^{-2} \, s^{-1}$

$$k = \frac{0.693}{37.9} s^{-1}, t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$$

$$t_{3/4} = \frac{2303}{1.83 \cdot 10^{-2} \, s^{-1}} \log \frac{[A]_0}{\underline{[A]_0}} = \frac{2303}{1.83} = 100 \log 2^2 \, s$$

$$t_{3/4}$$
 $\frac{2.303 \ 100 \ 2 \ 0.30105}{1.83}$ $\frac{138.64}{1.83}s$

t_{3/4} 75.76s

(ii)
$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{37.9} s^{-1} = 1.83 \cdot 10^{-2} s^{-1}$$

$$t = 1 \text{ minute} = 60 \text{ s}$$

$$t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$$

$$60 \text{ s} = \frac{2.303}{1.83 \cdot 10^{-2} \, \text{s}^{-1}} \log \frac{[A]_0}{[A]}$$

$$\log \frac{[A]_0}{[A]} \quad \frac{60 \quad 1.83 \quad 10^{-2}}{2.303} \quad 0.4768$$

$$\log \frac{[A]}{[A]_0} = 0.4768 - 1.5232$$

$$\frac{[A]}{[A]_0} = \text{Anti log } 1.5232 = 0.3336$$

$$\frac{[A]}{[A]_0} = 0.334$$

OR

$$(i) \quad t \quad \frac{2303}{k} \log \frac{[A]_0}{[A]}$$

Given:
$$k$$
 1.06 10 3 min 1 , $\frac{[A]_{0}}{[A]}$ $\frac{100}{85}$

$$t \frac{2.303}{1.06 \cdot 10^{-3} \text{ min}^{-1}} \log \frac{100}{85} \frac{2303}{1.06} [2 \log 10 - \log 85] \text{ min}$$

$$t \frac{2303}{1.06} [2 \cdot 1 \cdot 1.9294] \frac{2303}{1.06} \cdot 0.0706 = 153.39 \text{ min}$$

(ii) Given:
$$k = 1.06 \times 10^{-3} \text{ mm}^{-1}$$
, $\frac{[A]_0}{[A]} = \frac{100}{15}$
 $t = \frac{2.303}{1.06 \cdot 10^{-3} \text{ min}^{-1}}$, $\log \frac{100}{15} = \frac{2303}{1.06} [2 \log 10 - \log 15] \text{ min}$
 $\frac{2303}{1.06} [2 - 1 - 1.1761] = \frac{2303 - 0.8239}{1.06} \text{ min}$

t 1790 min.

CBSE (Foreign) SET-III


1. Schottky defect.

- Pent-2-enol: CH_3 — CH_2 —CH = CH—C—H
- (i) Zone refining based on the principle that impurities are more soluble in the melt than in the solid state of the metal.
 - (ii) Vapour phase refining of metals: In this method, crude metal is freed from impurities by first converting it into volatile compound and collected elsewhere. It is then decomposed to give pure metal. For example refining of nickel by Mond process.

12. (i) No. of electron pairs around central atom (Xe) =
$$6 \frac{b.p.}{l.p.}$$
 2

The shape would be square planar.

O



H₂S₂O₇ (Pyrosulphuric acid)

O

Fat soluble vitamins	Sources	Deficiency diseases
Vitamin A	Fish liver oil, carrots, milk	Xerophthalmia, Night blindness.
Vitamin D	Exposure to sunlight, fish and egg. Rickets and osteomalacia.	
Vitamin E	Vegetable oils like wheat germ oil, sunflower oil, etc.	Sterility and muscular atrophy
Vitamin K	Green leafy vegetables	Haemorrhages and increased blood clotting time.

- **18.** (*i*) **Nucleotides:** The monomeric unit of nucleic acid is called nucleotide. When a nucleoside is linked to phosphoric acid at 5' position of sugar moiety, we get nucleotide.
 - (ii) Nucleoside: A nucleoside is the condensation product of purine or pyrimidine base with pentose sugar.

24. (*i*) Linkage isomerism: The isomers which have same molecular formula but differ in the linkage of ligand atom to the central metal atom are called linkage isomers, *e.g.*,

$$[Co(NH_3)_5 NO_2]Cl_2$$
 and $[Co(NH_3)_5ONO]Cl_2$

Pentaamminenitrito-N-Cobalt (III) chloride pentaamminenitrito-O-Cobalt (III) chloride

- (ii) Outer orbital complex: When ns, np and nd orbitals are involved in hybridisation, outer orbital complex is formed, e.g., $[CoF_6]^2$ in which cobalt is sp^3d^2 hybridised.
- (iii) Bidentate ligand: When a ligand is bound to a metal ion through two donor atoms it is said to be bidentate ligand, e.g., $H_2\ddot{N}$ — CH_2 — CH_2 — $\ddot{N}H_2$ (ethane–1, 2–diamine, $C_2O_4^2$ (oxalate), etc.

27.	Polymer	Monomer	Structure
	PVC	Vinyl chloride	$CH_2 = CH - Cl$
	Teflon	Tetrafluro ethene	$CF_2 = CF_2$
	Neoprene	Chloroprene	CH ₂ C—CH CH ₂
30.	(a) (i) SO_2 $2H_2O$	SO_4^2 4H 2e	Cl] 5
	MnO_4 8H 5 e	$Mn^2 4H_2O]$	2
	2MnO ₄ 5SO ₂ 2H ₂	$_2$ O $2Mn^2$ $5SO_4^2$	4H
	(ii) 3HgCl ₂ 2PH ₃	Hg ₃ P ₂ 6HCl	

- (b) (i) Sulphur has a greater tendency for catenation than oxygen because S-S bond is stronger than O - O bond due to less interelectronic repulsions.
 - (ii) It is due to
 - low enthalpy of dissociation of F-F bond.
 - high hydration enthalpy of F⁻.
 - (iii) This is due to inert pair effect.

OR

- $PH_3 + 3NaH_2PO_2$ (a) (i) $P_4 + 3NaOH + 3H_2O$
 - $3Cu(NO_3)_2 + 2NO + 4H_2O$ (ii) 3Cu + 8HNO₃ (dilute)
- (b) (i) Because of small size and high electro negativity of oxygen, molecules of water are highly associated through H-bonding resulting in its liquid state.
 - (ii) Its reaction with iron produces H₂

Fe + 2HCl
$$FeCl_2 + H_2$$

Liberation of hydrogen prevents the formation of ferric chloride.

(iii) Because of its very low solubility in blood it prevents 'bends'.

CBSE EXAMINATION PAPERS

DELHI-2010

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Delhi) SET-I

- 1. Write a feature which will distinguish a metallic solid from an ionic solid.
- 2. Define 'order of a reaction'.
- **3.** What is an emulsion?
- **4.** Why does NO₂ dimerise?
- **5.** Give an example of linkage isomerism.
- **6.** A solution of KOH hydrolyses CH₃CHC*l*CH₂CH₃ and CH₃CH₂CH₂CH₂C*l*. Which one of these is more easily hydrolysed?
- 7. Draw the structural formula of 1-phenylpropan-1-one molecule.
- 8. Give the IUPAC name of $H_2N CH_2 CH_2 CH = CH_2$.
- **9.** Non-ideal solutions exhibit either positive or negative deviations from Raoult's law. What are these deviations and why are they caused? Explain with one example for each type.
- **10.** A reaction is of first order in reactant A and of second order in reactant B. How is the rate of this reaction affected when (i) the concentration of B alone is increased to three times (ii) the concentrations of A as well as B are doubled?
- 11. The rate constant for a reaction of zero order in A is 0.0030 mol L⁻¹ s⁻¹. How long will it take for the initial concentration of A to fall from 0.10 M to 0.075M?
- **12.** Draw the structures of white phosphorus and red phosphorus. Which one of these two types of phosphorus is more reactive and why?
- **13.** Explain the following observations:
 - (i) Generally there is an increase in density of elements from titanium (Z = 22) to copper (Z = 29) in the first series of transition elements.
 - (ii) Transition elements and their compounds are generally found to be good catalysts in chemical reactions.

- **14.** Name the following coordination compounds according to IUPAC system of nomenclature:
 - (i) $[Co(NH_3)_4 (H_2O) Cl]Cl_2$
 - (ii) $[CrCl_2(en)_2]Cl$, (en = ethane -1, 2 diamine)
- **15.** Illustrate the following reactions giving a chemical equation for each:
 - (i) Kolbe's reaction,
- (ii) Williamson synthesis.
- **16.** How are the following conversions carried out?
 - (i) Benzyl chloride to benzyl alcohol,
 - (ii) Methyl magnesium bromide to 2-methylpropan-2-ol.
- **17.** Explain the following terms:
 - (i) Invert sugar
- (ii) Polypeptides

OR

Name the products of hydrolysis of sucrose. Why is sucrose not a reducing sugar?

- **18.** What are essential and non-essential amino acids in human food? Give one example of each type.
- 19. The well known mineral fluorite is chemically calcium fluoride. It is known that in one unit cell of this mineral there are 4 Ca^{2+} ions and 8 F^- ions and that Ca^{2+} ions are arranged in a fcc lattice. The F $^-$ ions fill all the tetrahedral holes in the face centred cubic lattice of Ca^{2+} ions. The edge of the unit cell is 5.46×10^{-8} cm in length. The density of the solid is 3.18 g cm^{-3} . Use this information to calculate Avogadro's number (Molar mass of $\text{CaF}_2 = 78.08 \text{ g mol}^{-1}$)
- **20.** A solution prepared by dissolving 1.25 g of oil of winter green (methyl salicylate) in 99.0 g of benzene has a boiling point of 80.31°C. Determine the molar mass of this compound. (B.P. of pure Benzene = 80.10°C and K_b for benzene = 2.53°C kg mol⁻¹)
- 21. What is the difference between multimolecular and macromolecular colloids? Give one example of each type. How are associated colloids different from these two types of colloids?
- **22.** Describe how the following changes are brought about:
 - (i) Pig iron into steel.
 - (ii) Zinc oxide into metallic zinc.
 - (iii) Impure titanium into pure titanium.

OR

Describe the role of

- (i) NaCN in the extraction of gold from gold ore.
- (ii) SiO₂ in the extraction of copper from copper matte
- (iii) Iodine in the refining of Zirconium

Write chemical equations for the involved reactions.

- **23.** How would you account for the following?
 - (i) The atomic radii of the metals of the third (5d) series of transition elements are virtually the same as those of the corresponding members of the second (4d) series.

118 Xam idea Chemistry—XII

- (ii) The E $^{\circ}$ value for the Mn 3 /Mn 2 couple is much more positive than that for Cr 3 /Cr 2 couple or Fe 3 /Fe 2 couple.
- (iii) The highest oxidation state of a metal is exhibited in its oxide or fluoride.
- **24.** (i) State one use each of DDT and iodoform.
 - (ii) Which compound in the following couples will react faster in S_{N^2} displacement and why?
 - (a) 1-Bromopentane or 2-bromopentane
 - (b) 1-Bromo-2-methylbutane or 2-bromo-2-methylbutane.
- **25.** In the following cases rearrange the compounds as directed:
 - (i) In an increasing order of basic strength:

$$C_6H_5NH_2$$
, $C_6H_5N(CH_3)_2$, $(C_2H_5)_2$ NH and CH_3NH_2

(ii) In a decreasing order of basic strength:

Aniline, p-nitroaniline and p-toluidine

(iii) In an increasing order of pK_b values:

- **26.** Give **one** example each of
 - (i) addition polymers,
- (ii) condensation polymers,

- (iii) copolymers.
- 27. What are analgesic medicines? How are they classified and when are they commonly recommended for use?
- **28.** (a) State Kohlrausch law of independent migration of ions. Write an expression for the molar conductivity of acetic acid at infinite dilution according to Kohlrausch law.
 - (b) Calculate m for acetic acid.

Given that
$$_{m}$$
 (HCl) = 426 S cm² mol⁻¹
 $_{m}$ (NaCl) = 126 S cm² mol⁻¹
 $_{m}$ (CH₃COONa) = 91 S cm² mol⁻¹

OR

- (a) Write the anode and cathode reactions and the overall reaction occurring in a lead storage battery.
- (b) A copper-silver cell is set up. The copper ion concentration is 0.10 M. The concentration of silver ion is not known. The cell potential when measured was 0.422 V. Determine the concentration of silver ions in the cell. (Given E $_{\rm Ag}$ /Ag 0.80V, E $_{\rm Cu}^2$ /Cu 0.34 V)
- **29.** (a) Complete the following chemical equations:
 - $\begin{array}{ccc} \textit{(i)} \;\; \text{NaOH}_{(aq)} \quad \text{Cl}_{\,2(g)} \\ & \text{(Hot and conc.)} \end{array}$
 - (ii) $XeF_6(s)$ $H_2O(l)$

- (b) How would you account for the following?
 - (i) The value of electron gain enthalpy with negative sign for sulphur is higher than that for oxygen.
 - (ii) NF₃ is an exothermic compound but NCl₃ is endothermic compound.
 - (iii) CIF₃ molecule has a T-shaped structure and not a trigonal planar one.

OR

- (a) Complete the following chemical reaction equations::
 - $(i) P_4 SO_2Cl_2$
 - (ii) XeF₄ H₂O
- (b) Explain the following observations giving appropriate reasons:
 - (i) The stability of +5 oxidation state decreases down the group in group 15 of the periodic table.
 - (ii) Solid phosphorus pentachloride behaves as an ionic compound.
 - (iii) Halogens are strong oxidizing agents.
- **30.** (a) Explain the mechanism of a nucleophilic attack on the carbonyl group of an aldehyde or a ketone.
 - (b) An organic compound (A) (molecular formula C₈H₁₆O₂) was hydrolysed with dilute sulphuric acid to give a carboxylic acid (B) and an alcohol (C). Oxidation of (C) with chromic acid also produced (B). On dehydration (C) gives but-1-ene. Write the equations for the reactions involved.

OR

- (a) Give chemical tests to distinguish between the following pairs of compounds:
 - (i) Ethanal and propanal
 - (ii) Phenol and Benzoic acid
- (b) How will you bring about the following conversions?
 - (i) Benzoic acid to benzaldehyde
 - (ii) Ethanal to but-2-enal
 - (iii) Propanone to propene

Give complete reaction in each case.

CBSE (Delhi) SET-II

Questions Uncommon to Set-I

- 1. Which point defect in crystals of a solid does not change the density of the solid?
- **4.** What is the oxidation number of phosphorus in H₃PO₂ molecule?
- **5.** Give an example of coordination isomerism.
- 12. Draw the structural formulae of molecules of following compounds:
 - (i) BrF_3 and (ii) XeF_4

120 Xam idea Chemistry—XII

- 14. Describe the shape and magnetic behaviour of following complexes:
 - (*i*) $[Co(NH_3)_6]^3$
 - (ii) $[Ni(CN_4)]^2$, (At. No. Co = 27, Ni = 28)
- **15.** Explain the following reactions with an example for each:
 - (i) Reimer-Tiemann reaction
 - (ii) Friedel-Crafts reaction.
- **16.** How are the following conversions carried out?
 - (i) Propene to propan-2-ol
 - (ii) Ethylmagnesium chloride to propan-1-ol.
- 22. A solution of glycerol ($C_3H_8O_3$; molar mass = 92 g mol $^{-1}$) in water was prepared by dissolving some glycerol in 500 g of water. This solution has a boiling point of 100.42°C. What mass of glycerol was dissolved to make this solution? K_b for water = 0.512 K kg mol $^{-1}$.
- **24.** Complete the following chemical equations:
 - (i) $C_6H_5N_2Cl + C_6H_5NH_2$
 - (ii) C₆H₅N₂Cl + CH₃CH₂OH
 - (iii) RNH₂ + CHCl₃ + KOH
- **25.** Write the name and structure of the monomer of each of the following polymers:
 - (i) Neoprene
 - (ii) Buna-S
 - (iii) Teflon

CBSE (Delhi) SET-III

Questions Uncommon to Set-I and Set-II.

- 1. Which point defect in crystals of a solid decreases the density of the solid?
- **2.** Define 'rate of a reaction'.
- **3.** Give an example of 'shape-selective catalyst'.
- **4.** Draw the structure of O_3 molecule.
- **5.** Give an example of ionization isomerism.
- **13.** Explain the following observations:
 - (i) Transition elements generally form coloured compounds.
 - (ii) Zinc is not regarded as a transition element.
- **18.** State clearly what are known as nucleosides and nucleotides.
- 19. The density of copper metal is 8.95 g cm⁻³. If the radius of copper atom is 127.8 pm, is the copper unit cell a simple cubic, a body-centred cubic or a face-centred cubic structure? (Given: At. mass of Cu=63.54 g mol⁻¹ and $N_A = 6.022 \times 10^{23}$ mol⁻¹)

- **20.** How are the following colloids different from each other in respect of their dispersion medium and dispersed phase? Give one example of each.
 - (i) Aerosol
- (ii) Emulsion
- (iii) Hydrosol
- **26.** Differentiate between thermoplastic and thermosetting polymers. Give one example of each.
- **27.** Explain the following terms with one suitable example in each case.
 - (i) Cationic detergents
- (ii) Enzymes
- (iii) Antifertility drugs



CBSE (Delhi) SET-I

- 1. Metals are malleable and ductile whereas ionic solids are hard and brittle.
- 2. The order of a reaction can be defined as the sum of the powers of the concentration terms as expressed in rate law.
- An emulsion is a colloidal dispersion of one liquid in another liquid. For example, cod liver oil, milk etc.
- **4.** The dimerisation of NO₂ is because of unpaired electron on nitrogen atom to form N₂O₄ molecule with even number of electrons.

$$\begin{array}{ccc} 2NO_2 & \stackrel{262\,\mathrm{K}}{\longleftarrow} & N_2\,O_4 \\ \text{Paramagnetic} & \text{Colourless and diamagnetic} \end{array}$$

- 5. The example of linkage isomerism: $[Co(NH_3)_5 NO_2]^2$ and $[Co(NH_3)_5 ONO]^2$.
- **6.** CH₃CH₂C*l*CHCH₃

7.
$$CH_3 - CH_2 - C - C$$

- **8.** But-3-en-1-amine.
- **9.** When the vapour pressure of a solution is either higher or lower than that predicted by Raoult's law then the solution exhibits deviation from Raoult's law.

These deviation are caused when solute-solvent molecular interactions $(A \ B)$ are either weak or stronger than solvent-solvent $(A \ A)$ or solute-solute $(B \ B)$ molecular interactions.

Positive Deviation: When $(A \ B)$ molecular interactions are weaker than $(A \ A)$ and $(B \ B)$ molecular interactions. For example a mixture of ethanol and acetone.

Negative deviation: When $(A \ B)$ molecular interactions are stronger than $(A \ A)$ and $(B \ B)$ molecular interactions. For example a mixture of chloroform and acetone.

$$H_{3}C$$
 $C = 0 \cdots H - C Cl$

10. According to the given reaction:

Rate =
$$k [A] [B]^2$$

(i) When the concentration of reactant 'B' is increased three times the rate of reaction becomes 9 times.

i.e., Rate =
$$k [A] [3B]^2 = 9k[A] [B]^2$$

(ii) When the concentration of reactants A and B are doubled, then rate becomes 8 times.

i.e., Rate =
$$k [2A]^1 [2B]^2 = 8k[A] [B]^2$$

11. According to available data:

$$k = 0.0030 \text{ mol } L^{-1} s^{-1},$$
 [R]₀ = 0.10 M, [R] = 0.075 M

We know that


$$[R] = -kt + [R]_0$$

 $0.075 = -0.0030 t + 0.10$
 $3t \quad 100 \quad 75$ $t = 8.33 \text{ second}$

12. White phosphorus is more reactive due to its discrete tetrahedral structure and angular strain.


or

Red phosphorus is less reactive due to its polymeric structure.





- 13. (i) The density of elements from titanium to copper increase in the first series of transition elements. This is due to decrease in metallic radius coupled with increase in atomic mass results in a general increase in the density.
 - (ii) Many transition metals and their components show catalytic properties. This property is due to their ability to exhibit variable oxidation states (incomplete d-orbitals) which enable them to form unstable intermediates.
- 14. (i) Tetraammineaquachloridocobalt (III) chloride
 - (ii) Dichlorido-bis (ethane-1, 2-diamine) chromium(III)chloride

15. (*i*) Kolbe's reaction




(ii) Williamson synthesis





17. (i) Invert Sugar: The dextrorotatory sucrose when hydrolysed by boiling with mineral acid produces an equal number of molecules of dextrorotatory fructose. The resulting mixture is laevorotatory and termed as invert sugar.



(ii) **Polypeptides:** Polypeptides are polymers formed by condensation of more than ten amino acids. They have large number of peptides bonds in them. Polypeptides are amphoteric because of the presence of terminal-ammonium and carboxylate ions as well as the ionized side chains of amino acid residues

OR

Sucrose gives glucose and fructose on hydrolysis. It is a non-reducing sugar as it does not contain free ketone or aldehyde group in its ring form.

18. Essential amino acids: Amino acids which cannot be synthesized in the body but must be obtained through diet, are called essential amino acids. For example: valine, leucine.

Non-essential amino acids: The amino acids, which can be synthesized in the body are known as non-essential amino acids. For example: Glycine, Alanine.

19. Given:
$$d = 3.18 \text{ g cm}^{-3}$$
, $Z = 4$, $a = 5.46 \times 10^{-8} \text{ cm}$

$$M = 78.08 \text{ g mol}^{-1} N_A = ?$$

$$d \frac{Z M}{a^3 N_A}$$

$$N_A \frac{Z M}{a^3 P}$$

Substituting the given values, we get

$$N_A = \frac{4 \cdot 78.08}{(5.46 \cdot 10^{-8})^{-3} \cdot 3.18}$$

$$N_A = \frac{313 \cdot 32 \cdot 10^{24}}{517 \cdot 61}$$

$$N_A = 6.035 \times 10^{23} \text{ mol}^{-1}$$

20. Given
$$T_B = 80.31 - 80.10 = 0.21$$
 °C or 0.21 K

$$W_B = 1.25 \text{g}, K_b = 2.53 \text{°C kg mol}^{-1}$$

 $M_B = ?$
 $W_A = 99 \text{g}$
 $M_B = \frac{K_b W_B}{T_b W_A} = 1000$
 $= \frac{2.53 + 1.25 + 1000}{0.21 + 99} = \frac{3162.5}{20.79}$

$$M_B = 152 \text{ g/mol}$$

21. Multimolecular colloids: In this type of colloids, colloidal particles are aggregates of atoms or molecules each having size less than 1nm, e.g., sulphur sol, gold sol.

Macromolecular colloids: In this type of colloids, colloidal particles are themselves large molecules of colloidal dimensions, e.g., starch, proteins, polyethene, etc.

Associated colloids: There are certain substances which at low concentrations behave as normal electrolyte, but at higher concentrations exhibit colloidal behaviour due to the formation of aggregates. Such colloids are known as associated colloids, e.g., soaps and detergents.

- 22. (i) Pig iron into steel: Pig iron is converted into steel by heating in a converter. A blast of oxygen diluted with carbon dioxide is blown through the converter. Oxygen reacts with impurities and raised the temperature to 2173K. Carbon gets oxidised to CO which burns off at the mouth of the converter. Oxides of silicon and Mg form slag. When the flame is stopped, slag is tapped off and other metals like Mn, Cr, Ni, W may be added in the end.
 - (ii) Metallic zinc can be obtained from zinc oxide. At first calcination of ZnO is done and converted into sinters of oxide.

$$ZnO + C$$
 $Zn + CO$

The oxide is then made into brickettes with coke and clay and heated by producer gas in vertical retorts at 1673 K, zinc, boiling point is 1183 K, distills off and is collected by rapid chilling.

(iii) Impure Titanium into pure Titanium: Impure Titanium is heated with I_2 to form volatile complex (T_iI_4) , which on heating at higher temperature decomposes to give pure titanium.

Ti (impure) +
$$2 I_2$$
 Ti I_4 T_i (pure) + I_2 OR

(i) Role of NaCN in the extraction of gold is to do the leaching of gold ore in the presence of air from which the gold is obtained later by replacement.

$$4Au(s) + 8NaCN(aq) + 2H2O + O2$$
 $4Na[Au(CN)2] + 4KOH$

(ii) SiO₂ is added to copper matte to convert the leftout FeS, FeO into slag.

(iii) Iodine is heated with impure Zr to form volatile compound which on further heating decomposes to give pure zirconium.

$$Zr (Impure) + I_2$$
 ZrI_4 $Zr (pure) + 2I_2$

23. (*i*) This is due to lanthanide contraction.

or

This is due to filling of 4f orbitals which have poor shielding effect.

- (ii) The E° value for the Mn³+/Mn²+ couple is much positive than Cr³+/Cr²+ couple or Fe³+/Fe²+ couple because Mn³+ ion receiving an electron gets d-subshell half-filled which is highly stable. While in case of Fe³+ d-sub shell is already half-filled, so it does not receive electron easily.
- (iii) This is because fluorine and oxygen are highly electronegative elements and have small size.
- **24.** (*i*) **DDT:** It is used as insecticide to control flies, mosquitoes, etc. Iodoform: Iodoform is used as an antiseptic.
 - (ii) 1-Bromopentane, as it is a primary alkyl halide. and
 - (b) 1-Bromo-2-methyl butane, as it is a primary alkylhalide.
- **25.** (i) Increasing order of basic strength is:

$$C_6H_5NH_2 < C_6H_5N(CH_3)_2 < CH_3NH_2 < (C_2H_5)_2NH_3$$

- (ii) p-Toluidine > Aniline > tr-nitroaniline
- (iii) $(C_2H_5)_2 NH_2 < C_2H_5NH_2 < C_6H_5NHCH_3 < C_6H_5NH_2$
- **26.** (*i*) Polythene, PVC.
 - (ii) Nylon 6, Nylon 6,6.
 - (iii) Buna-S, Buna-N.

- 27. Analgesics: They are the drugs used for relieving pain. Analgesics are classified into two categories:
 - (a) Narcotics (addictive) drugs
- (b) Non-narcotics (non-addictive) drugs
- (a) Narcotics: Narcotic drugs are recommended for cardiac pain, post operative pains, terminal cancer and in child birth.
- (b) Non-narcotics: Non-narcotic drugs are effective in relieving skeletal pain, reducing fever, preventing heart attack.
- **28.** (a) Kohlrausch Law: "The molar conductance of an electrolyte at infinite dilution is equal to the sum of the molar conductances of the two ions, i.e., the cation and the anion."

Mathematically,

Expression for the molar conductivity of acetic acid:

or
$${}^{\circ}_{m}$$
 CH₃COOH ${}^{\circ}_{m}$ CH₃COO ${}^{\circ}_{H}$ Or ${}^{\circ}_{m}$ CH₃COOH ${}^{\circ}_{m}$ (CH₃COO)

(b) Given

$$_{m}^{o}$$
 (HCl) 426 S cm² mol⁻¹

$$_{m}^{o}$$
 (CH₃COONa) 91 S cm² mol⁻¹

$$_{m}^{o}\left(\mathrm{CH_{3}COOH}\right) \quad _{m}^{o}\left(\mathrm{CH_{3}COO}\right) \quad _{m}^{o}(\mathrm{H}\right)$$

$$_{m}^{o}$$
 (CH₃COONA) $_{m}^{o}$ (CH₃COO) $_{m}^{o}$ (Na) ...(1)

$${}^{\mathrm{o}}_{m}(\mathrm{NaCl}) \quad {}^{\mathrm{o}}_{m}(\mathrm{Na}) \quad {}^{\mathrm{o}}_{m}(\mathrm{Cl}) \qquad ...(2)$$

$$_{m}^{0}$$
 (HCl) $_{m}^{0}$ (H) $_{m}^{0}$ (Cl) ...(3)

Subtracting (2) from the sum of (1) and (3), we get:

$${}^{o}_{m}(CH_{3}COO) \quad {}^{o}_{m}(Na) + {}^{o}_{m}(H) + {}^{o}_{m}(Cl) \quad {}^{o}_{m}((Na^{+}) \quad {}^{o}_{m}Cl)$$

$$= {}^{o}_{m}(CH_{3}COO) + {}^{o}_{m}(H^{+}) = {}^{o}_{m}(CH_{3}COOH)$$

$${}^{o}_{m}(CH_{3}COOH) \quad 91.0 \quad 426 \quad 126 \ S \ cm^{2} \ mol^{-1}$$

$$= 391.0 \ S \ cm^{2} \ mol^{-1}$$

OR

(a) When lead battery operates, the following cell reactions occur:

Anode half cell reaction:

$$Pb(s) + SO_4^2 (aq)$$
 $PbSO_4(s) + 2e^-$

Cathode half cell reaction:

$$PbO_{2}(s) + 4 H^{+}(aq) + SO_{4}^{2} + 2e^{-}$$
 $PbSO_{4}(s) + 2H_{2}O(l)$

Net reaction, $Pb(s) + PbO_2 + 2H_2SO_4(aq)$

$$2PbSO_4(s) + 2H_2O(l)$$

(b) Given
$$E^{o}_{Ag+/Ag} = 0.80V, \qquad E^{o}_{Cu^{2}/Cu} = +0.34V$$
$$[Cu^{2+}] = 0.10 \text{ M} \qquad [Ag^{+}] = ?$$
$$E^{o}_{cell} = E^{o}_{R} - E^{o}_{L} = 0.80 - 0.34V = 0.46V$$

From Nernst Equation:

$$E_{cell} = E_{cell}^{o} \frac{0.0591}{2} log \frac{[Cu^{2}]}{[Ag]^{2}}$$

$$0.422 = 0.46 \frac{0.591}{2} log \frac{[0.10]}{[Ag]^{2}}$$

$$log \frac{0.10}{[Ag]^{2}} 1.2881$$

$$[Ag^{+}]^{2} = 0.0051$$

$$[Ag^{+}] = 7.1 \times 10^{-2} M$$

29. (*i*) 6 NaOH +
$$3Cl_2$$
 5NaCl + NaClO₃ + $3H_2O$

(ii)
$$XeF_6 + 3H_2O$$
 $XeO_3 + 6HF$ or
$$XeF_6 + H_2O$$
 $XeOF_4 + 2HF$

- (i) Due to smaller size of oxygen the electron cloud is distributed over a small region of space, making electron density high which repels the incoming electrons.
- (ii) This is due to lower bond dissociation enthalpy of F_2 than Cl_2 and comparable size of fluorine and nitrogen.
- (iii) ClF₃ molecule has a T-shaped structure. This is due to presence of two lone pairs in the outer shell of chlorine in ClF₃ molecule which repel the bond pairs.

OR

(a) Complete the following chemical reaction equations:

(i)
$$P_4 + 8SO_2Cl_2$$
 $4PCl_3 + 4SO_2 + 2S_2Cl_2$
(ii) $6XeF_4 + 12H_2O$ $4Xe + 2XeO_3 + 24HF + 3O_2$

- (b) (i) The stability of +5 oxidation state decreases down the group in group 15 of the periodic table. The + 3 oxidation state becomes more and more common on moving down the group from N to Bi. This is because of inert pair effect.
 - (ii) Solid PCl_5 behaves as an ionic compound because it is a salt containing the tetrahedral cation $[PCl_4]^+$ and octahedral anion $[PCl_6]^-$.
 - (iii) Halogens are strong oxidising agents because they have high electron affinities, so, they pick up electrons from other substances.
- **30.** (a) Mechanism of nucleophilic addition reactions:

A nucleophile attacks the electrophilic carbon atom of the polar carbonyl group from a direction approximately perpendicular to the plane of sp^2 hybridized orbitals of carbonyl carbon. The hybridization of carbon changes from sp^2 to sp^3 in this process and a tetrahedral

intermediate is produced. The intermediate captures a proton from the medium to give the neutral product.

(b) The hydrolysis of the given compound with dil H₂SO₄ to give a carboxylic acid and an alcohol suggests that the compound is an ester.

$$\begin{array}{c} O & O \\ \parallel & \parallel \\ R-C-OR' & ^{dil.\,H_2SO_4} & R-C-OH & R-OH \\ \end{array}$$

So, the possible structure of 'A' is

$$\begin{array}{c} \text{O} \\ || \\ \text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{C} - \text{O} - \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{CH}_3 \\ \text{Butyl butanoate} \end{array}$$

$$(i) \ \operatorname{CH}_3 - \operatorname{CH}_2 - \operatorname{CH}_2 - \operatorname{CH}_2 - \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_3$$

$$(A) \qquad \qquad \operatorname{dil. H}_2 \operatorname{SO}_4 \qquad \operatorname{CH}_3 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{COOH} \qquad \operatorname{CH}_3 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{OH}$$

$$\qquad \qquad \operatorname{Butanoic acid} \qquad \operatorname{Butanoic} \qquad \operatorname{Butanoic} \qquad \operatorname{Butanoic} \qquad \operatorname{CH}_3 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{CH}_2 \operatorname{OH} \qquad \operatorname{CH}_3 \operatorname{CH}_2 \operatorname{C$$

$$\begin{array}{ccc} \textit{(ii)} & \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH} & \text{chromic acid} \\ & \text{Butanol} & \text{oxidation} & \text{CH}_3\text{CH}_2\text{CH}_2\text{COOH} \\ & \text{Butanoic acid} & \text{(B)} \end{array}$$

$$\begin{array}{ccc} \textit{(iii)} \ \ \text{CH}_3 - \text{CH}_2 \text{CH}_2 \text{CH}_2 \text{OH} & \text{dehydration} \\ & \text{But-1-ene} \end{array} \\ \text{But-1-ene}$$

OR

(a) Chemical tests to distinguish between ethanal and propanal

Iodoform Test: Ethanol gives iodoform test positive while propanal does not give this test.

$$CH_3CHO + 3I_2 + 4 NaOH$$
 $CHI_3 + HCOONa + 3NaI + 3H_2O$ (yellow)

(b) Phenol and benzoic acid:

Phenol gives violet colour with neutral FeCl₃ solution. Benzoic acid does not give violet colour with FeCl₃.

OH
$$6 \longrightarrow FeCl_{3} \longrightarrow [Fe(OC_{6}H_{5})_{6}]^{3-} + 3H^{+} + HCl$$
Violet complex

COOH
$$CH_2OH$$
 CHO

(i) $Benzoic acid$ CH_2OH CHO

Benzyl alcohol CHO

Benzaldehyde

(ii) Ethanal to but-2-enal

CH₃CHO dil. NaOH
$$\stackrel{|}{\text{CH}_3}$$
 CH₃ - CH- CH₂ - CHO $\stackrel{|}{\text{H}_2\text{O}}$ CH₃ - CH= CH- CHO But 2-en al

(iii) Propanone to propane

$$CH_3$$
 $C = O + 4[H]$ $Zn-Hg/HCl(conc.)$ $CH_3 - CH_2 - CH_3 + H_2O$


Propanone Propano

CBSE (Delhi) SET-II

- 1. Frankel defects do not change the density of the solids.
- **4.** Oxidation number of phosphorous in H_3PO_2 is +1.
- **5.** Example of coordination isomerism.


$$[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6] \ \ \text{and} \ \ [\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$$

12. (i) Structure of BrF_3



Bent T-shaped

(ii) Structure of XeF_4 molecule




Square planar

130 Xam idea Chemistry—XII



Pairing of electrons takes place against the Hund's role



Thus, the complex will be octahedral and diamagnetic.


(ii) [Ni(CN)₄]²

The electronic configuration of the nickel atom and Ni^{2} ion are depicted below:




Since coordination number in the complex is 4. It shows that complex is formed by sp^3 or dsp^2 hybridization.

However, experiments show that the complex is diamagnetic.




The resulting complex is square planar and diamagnetic as it has no unpaired electrons.


15. (i) Reimer Tiemann Reaction



(ii) Friedel - Craft Reactions Acylation:



Alkylation:



132 Xam idea Chemistry—XII

25. (*i*) The monomer of neoprone is chloroprene.

$$CH_2 = CH - C = CH_2$$

$$CH_2 = CH - C = CH_2$$

(ii) The monomer of Buna-S is Buta-1,3-diene and styrene.

$$\mathrm{CH}_2 = \mathrm{CH} - \mathrm{CH} = \mathrm{CH}_2$$
 and $\mathrm{C}_6\mathrm{H}_5\mathrm{CH} = \mathrm{CH}_2$
Buta 1,3 diene Styrene

(iii) The monomer of teflon is Tetrafluoroethene.

CBSE (Delhi) SET-III

- 1. Schottky defect.
- 2. Rate may be defined as the decrease in the concentration of reactant per unit time or increase in the concentration of product per unit time.
- 3. The example of shape selective catalyst is ZSM-5.

5. Example of ionisation isomerism is –

$$[Pt(NH_3)_4Cl_2]Br_2$$
 and $[Pt(NH_3)_4Br_2]Cl_2$

- 13. (i) This is due to d-d transition. When visible (white) light falls on a compound, it absorbs certain radiations of white light and transmit the remaining ones. The transmitted light has the complementary colour to that of the absorbed light.
 - (ii) Because the atoms or simple ions of zinc never have partially filled d orbitals.
- **18.** (*i*) **Nucleoside:** A nucleoside is the condensation product of purine or pyrimidine base with pentose sugar.
 - (ii) Nucleotides: The monomeric unit of nucleic acid is called nucleotide. When a nucleoside is linked to phosphoric acid at 5' position of sugar moiety, we get nucleotide.

19. Given, Radius
$$(r) = 127.8 \text{ pm}$$

For fcc
$$a = 2\sqrt{2}r$$

$$a^3 \quad (2\sqrt{2}r)^3 \quad 4.723 \quad 10^{-23} \text{ cm}^3$$

Atomic mass of
$$Cu = 63.54 \text{ g mol}^{-1}$$

$$N_A = 6.022 \times 1023 \text{ mol}^{-1}$$

Density =
$$8.95 \text{ g cm}^{-3}$$

$$d \quad \frac{Z \quad M}{N_A \quad a^3}$$

$$8.95 = \frac{Z + 63.54}{6.023 + 10^{23} + 4.723 + 10^{-23}}$$

$$Z = \frac{8.95 + 6.022 + 4.723}{63.54} \qquad Z = 4.00$$

Thus copper has fcc structure.

20.

	Aerosol Emulsion		Hydrosol
1.	Sols in which the	1. A colloidal system in which	1. Sols in which the dispersed
	dispersion medium is gas	dispersion medium and	phase is solid and
	and dispersed phase is solid	dispersed phase both are	dispersion medium is
	or liquid.	liquids. For example:	water.
	For example: Smoke, dust	Milk, cod liver oil.	For example: gum sol.

26. Differences between thermosetting and thermoplastic polymers.

Thermoplastics		Thermosetting plastics	
(<i>i</i>)	These polymer are linear or slightly branched chain molecules	(<i>i</i>)	These polymers are cross linked or heavily branched molecules
(ii)	Soften on heating and harden on cooling and can be remoulded.	(ii)	On heating undergo extensive cross linking in moulds and become infusible.
(iii)	Some common examples are polyethene, PVC, polystyrene, etc.	(iii)	Some common examples are bakelite, urea-formaldehyde resins, terylene, etc.

- **27.** (*i*) Cationic detergents: Cationic detergents are mostly acetate or chlorides of quaternary amines. For example: Cetyltrimethyl-ammonium chlorides.
 - (ii) Enzymes: Enzymes are the essential biological catalysts which catalyse specific biological reactions at a very high rate under mild conditions of temperature and pH. For example, invertase, zymase, pepsin, trypsin, etc.
 - (iii) Antifertility drugs: The drugs which are used to control the birth rate are known as antifertility drugs. For example, norethindrone, novestrol, etc.

CBSE EXAMINATION PAPERS

ALL INDIA-2010

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (All India) SET-I

- 1. What type of interactions hold the molecules together in a polar molecular solid?
- 2. What is meant by 'limiting molar conductivity'?
- 3. Fluorine does not exhibit any positive oxidation state. Why?
- **4.** Give the IUPAC name of the following compound:

$$\begin{array}{c} H_3C \\ H \end{array} = \begin{array}{c} CH_3 \\ H \\ CH_3 \end{array}$$

- **5.** Write the structure of the molecule of a compound whose IUPAC name is 1-phenylpropan-2-ol
- **6.** What is Tollen's reagent? Write one usefulness of this reagent.
- 7. What is meant by 'reducing sugars'?
- **8.** What does the designation '6, 6' mean in the name nylon-6, 6?
- **9.** Define the terms, 'osmosis' and 'osmotic pressure'. What is the advantage of using osmotic pressure as compared to other colligative properties for the determination of molar masses of solutes in solutions.
- **10.** Express the relation among the cell constant, the resistance of the solution in the cell and the conductivity of the solution. How is the conductivity of a solution related to its molar conductivity?
- 11. Given that the standard electrode potential (E°) of metals are:

$$K/K = -2.93 \text{ V}, \text{Ag} / \text{Ag} 0.80 \text{ V}, \text{Cu}^2/\text{Cu} = 0.34 \text{ V}, \text{Mg}^{2+}/\text{Mg} 2.37 \text{ V}, \text{Cr}^3/\text{Cr} 0.74 \text{ V},$$

 $Fe^{2+}/Fe 0.44 \text{ V}.$

Arrange these metals in an increasing order of their reducing power.

OR

Two half-reactions of an electrochemical cell are given below:

$$MnO_{4}^{-}(aq) + 8H^{+}(aq) + 5e^{-}$$
 $Mn^{2+}(aq) + 4H_{2}O(l)$, $E^{\circ} = + 1.51 \text{ V}$
 $Sn^{2+}(aq)$ $Sn^{4+}(aq) + 2e^{-}$, $E^{\circ} = + 0.15V$

Construct the redox reaction equation from the two half-reactions and calculate the cell potential from the standard potentials and predict if the reaction is reactant or product favoured.

- **12.** Describe the following:
 - (i) Tyndal effect
- (ii) Shape-selective catalysis
- **13.** What is meant by coagulation of a colloidal solution? Name any method by which coagulation of lyophobic sols can be carried out.
- **14.** Complete the following chemical reaction equations:

(i)
$$I_2 + HNO_3$$
 (ii) $HgCl_2 + PH_3$

- **15.** Draw the structural formulae of the following compounds:
 - (i) H₄P₂O₅
- (ii) XeF₄
- **16.** Give the chemical tests to distinguish between the following pairs of compounds:
 - (i) Ethylamine and Aniline
 - (ii) Aniline and Benzylamine
- 17. Identify A and B in each of the following processes:

(i)
$$CH_3CH_2Cl$$
 NaCN A Reduction B Ni/H₂ (ii) $C_6H_5NH_2$ NaNO₂/HCl A $C_6H_5NH_2$ OH OH

18. Draw the molecular structures of the monomers of

(i) PVC

- (ii) Teflon
- 19. The density of copper metal is 8.95 g cm⁻³. If the radius of copper atom be 127.8 pm, is the copper unit cell simple cubic, body-centred cubic or face-centred cubic?

(Given: atomic mass of $Cu = 63.54 \text{ g mol}^{-1}$ and $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

- **20.** What mass of NaCl (molar mass = 58.5 g mol^{-1}) must be dissolved in $65 \text{ g of water to lower the freezing point by 7.5°C? The freezing point depression constant, <math>K_f$, for water is 1.86 K kg mol⁻¹. Assume van't Hoff factor for NaCl is 1.87.
- **21.** Describe the role of the following:
 - (i) NaCN in the extraction of silver from a silver ore
 - (ii) Iodine in the refining of titanium
 - (ii) Cryolite in the metallurgy of aluminium

OR

Describe the principle involved in each of the following processes of metallurgy:

- (i) Froth floatation method
- (ii) Electrolytic refining of metals
- (ii) Zone refining of metals

136 Xam idea Chemistry—XII

- **22.** Explain the following cases giving appropriate reasons:
 - (i) Nickel does not form low spin octahedral complexes.
 - (ii) The -complexes are known for the transition metals only.
 - (iii) Co²⁺ is easily oxidised to Co³⁺ in the presence of a strong ligand.
- 23. How would you differentiate between S_N1 and S_N2 mechanisms of substitution reactions? Give one example of each.
- **24.** How would you convert the following:
 - (i) Phenol to benzoquinone
 - (ii) Propanone to 2-methylpropan-2-ol
 - (iii) Propene to propan-2-ol
- **25.** How would you account for the following:
 - (i) NCl₃ is an endothermic compound while NF₃ is an exothermic one.
 - (ii) XeF₂ is a linear molecule without a bend.
 - (iii) The electron gain enthalpy with negative sign for fluorine is less than that for chlorine, still fluorine is a stronger oxidising agent than chlorine.
- **26.** Amino acids may be acidic, alkaline or neutral. How does this happen? What are essential and non-essential amino acids? Name one of each type.
- **27.** Explain the following terms with one example in each case:
 - (i) Food preservatives
 - (ii) Enzymes
 - (iii) Detergents
- **28.** (a) Explain the following terms:
 - (i) Rate of a reaction
 - (ii) Activation energy of a reaction
 - (b) The decomposition of phosphine, PH₃, proceeds according to the following equation:

$$4PH_3(g)$$
 $P_4(g) + 6H_2(g)$

It is found that the reaction follows the following rate equation:

Rate =
$$k[PH_3]$$

The half-life of PH₃ is 37.9 s at 120°C.

- (i) How much time is required for 3/4th of PH₃ to decompose?
- (ii) What fraction of the original sample of PH₃ remains behind after 1 minute?

OR

- (a) Explain the following terms:
 - (i) Order of a reaction
 - (ii) Molecularity of a reaction

- (b) The rate of a reaction increases four times when the temperature changes from 300 K to 320 K. Calculate the energy of activation of reaction, assuming that it does not change with temperature. ($R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$)
- **29.** (a) Complete the following chemical equations:
 - (i) $\operatorname{Cr}_2 \operatorname{O}_7^{2-}(aq) + \operatorname{H}_2 \operatorname{S}(g) + \operatorname{H}^+(aq)$
 - (ii) $Cu^{2+}(aq) + \Gamma(aq)$
 - (b) How would you account for the following:
 - (i) The oxidising power of oxoanions are in the order $VO_2^+ < Cr_2O_7^{2-} \quad MnO_4^-$.
 - (ii) The third ionization enthalpy of manganese (Z = 25) is exceptionally high.
 - (iii) Cr²⁺ is a stronger reducing agent than Fe²⁺.

OR

- (a) Complete the following chemical equations:
 - (i) $MnO_4^-(aq) + S_2O_3^{2-}(aq) H_2O(l)$
 - (ii) $\operatorname{Cr}_2 \operatorname{O}_7^{2-}(aq) + \operatorname{Fe}^{2+}(aq) + \operatorname{H}^+(aq)$
- (b) Explain the following observations:
 - (i) La^{3+} (Z = 57) and Lu^{3+} (Z = 71) do not show any colour in solutions.
 - (ii) Among the divalent cations in the first series of transition elements, manganese exhibits the maximum paramagnetism.
 - (iii) Cu⁺ ion is not known in aqueous solutions.
- **30.** (a) Illustrate the following name reactions giving a chemical equation in each case:
 - (i) Clemmensen reaction
 - (ii) Cannizzaro's reaction
 - (b) Describe how the following conversions can be brought about:
 - (i) Cyclohexanol to cyclohexan-1-one
 - (ii) Ethylbenzene to benzoic acid
 - (iii) Bromobenzene to benzoic acid

OR

- (a) Illustrate the following name reactions:
 - (i) Hell-Volhard-Zelinsky reaction
 - (ii) Wolff-Kishner reduction reaction
- (b) How are the following conversions carried out:
 - (i) Ethylcyanide to ethanoic acid
 - (ii) Butan-1-ol to butanoic acid
 - (iii) Methylbenzene to benzoic acid

Write chemical equations for the involved reactions.

CBSE (All India) SET-II

Questions uncommon to Set-I

- 1. What type of semiconductor is obtained when silicon is doped with arsenic?
- 2. Nitrogen is relatively inert as compared to phosphorus. Why?
- **6.** What are monosaccharides?
- **8.** What is meant by 'copolymerisation'?
- **13.** Define the following:
 - (i) Peptization
 - (ii) Reversible sols
- **15.** Complete the following chemical reaction equations:
 - $\begin{array}{ccc} (i) & \text{NaOH} & + & \text{Cl}_2 \\ & \text{(cold and dilute)} \end{array}$
 - (ii) XeF₆ + H_2O (excess)
- 17. Give the chemical tests to distinguish between the following pairs of compounds:
 - (i) Methylamine and Dimethylamine
 - (ii) Aniline and N-methylaniline
- **18.** Draw the structures of the monomers of the following polymers:
 - (i) Bakelite
 - (ii) Nylon-6
- **19.** Silver crystalises in face-centred cubic unit cells. Each side of the unit cell has a length of 409 pm. What is the radius of silver atom?
- **20.** What mass of ethylene glycol (molar mass = 62.0 g mol^{-1}) must be added to 5.50 kg of water to lower the freezing point of water from 0°C to -10.0°C ? (K_f for water = $1.86 \text{ K kg mol}^{-1}$)
- 27. What are analgesic drugs? How are they classified and when are they usually recommended for use?

CBSE (All India) SET-III

Questions uncommon to Set-II and Set-II

- 1. Write a distinguishing feature of metallic solids?
- **3.** Differentiate between molarity and molality of a solution.
- **8.** What are the products of hydrolysis of sucrose?
- 17. Silver crystallises in fcc lattice. If the edge length of the unit cell is 4.07×10^{-8} cm and the density of the crystal is 10.5 g cm⁻³, calculate the atomic mass of silver. (N_A = 6.02×10^{23} atoms mol⁻¹)
- 20. 15 g of an unknown molecular substance was dissolved in 450 g of water. The resulting solution freezes at -0.34°C. What is the molar mass of the substance? (K_f for water = 1.86 K kg mol⁻¹).

- (i) The electron gain enthalpy with negative sign is less for oxygen than that for sulphur.
- (ii) Phosphorus shows greater tendency for catenation than nitrogen.
- (iii) Fluorine never acts as the central atom in polyatomic interhalogen compounds.
- **25.** Write the name, the state of hybridization, the shape and the magnetic behaviour of the following complexes:

$$[CoCl_4]^{2-}$$
, $[Ni(CN)_4]^{2-}$, $[Cr(H_2O)_2(C_2O_4)_2]^{-}$

(Atomic number: Co = 27, Ni = 28, Cr = 24)

- **26.** Differentiate between fibrous proteins and globular proteins. What is meant by the denaturation of a protein?
- **27.** Explain the following terms with an example for each:
 - (i) Antibiotics
- (ii) Antiseptics
- (iii) Analgesics



CBSE (All India) SET-I

- 1. The interactions hold the molecules together in a polar molecular solid is dipole-dipole attractions.
- **2.** When concentration of an electrolyte approaches zero, the molar conductivity is known as limiting molar conductivity $\binom{m}{m}$.
- **3.** Fluorine does not exhibit any positive oxidation state because it is the most electronegative element in the periodic table.
- **4.** 4–Bromo–3–methyl–pent–2–ene
- 5.

6. Tollen's reagent – A solution of AgNO₃ dissolved in NH₄OH is known as Tollen's reagent. This is used to detect the presence of –CHO group in an organic compound. For example:

$$RCHO + 2Ag (NH_3)_2 OH$$
 $RCOONH_4 + 2Ag + H_2O + 3NH_3.$

- **7.** Reducing Sugars: The carbohydrates which reduces Fehling's Reagent and Tollen's Reagent are referred to as reducing sugar. For example, all monosaccharide are reducing sugars.
- **8.** Each monomer of nylon 6, 6 consists of six carbon each.
- **9. Osmosis:** The phenomenon in which solvent molecules flow through a semipermeable membrane from a solution of low concentration to a solution of higher concentration is termed as osmosis.

Osmotic Pressure: The hydrostatic pressure exerted by the column of the solution which is just sufficient to prevent the osmosis is called osmotic pressure.

Advantage of using osmotic pressure as compared to other collegative properties:

- (i) The equilibrium is established very quickly. Hence, results are obtained in a very short time.
- (ii) The concentration of the solution does not change during determination of osmotic pressure.
- (iii) This is ideal method for proteins to find their molar mass since it happens at room temperature and no coagulation occurs.
- 10. Conductivity $(K) = \frac{1}{\text{Resistance } (R)} \times \text{Cell constant}$

Molar Conductivity (
$$_{m}$$
) = $\frac{\text{Conductivity () } 1000}{\text{Molarity (}M\text{)}}$

$$OR$$

$$_{m} = \frac{K \cdot 1000}{M} = \frac{m \cdot M}{1000}$$

$$K = \frac{138.95 \text{ cm}^{2} \text{ mol}^{-1} \cdot 1.5 \text{ mol L}^{-1}}{1000 \text{ cm}^{3} \text{ L}^{-1}} = 0.208355 \text{ cm}^{-1}$$

11. Increasing order of reducing power:

(i)
$$MnO_4 + 8H^+ + Se^ Mn^{2+} + 4H_2O$$
, $E = + 1.51V$ (Reduction)

(ii)
$$\operatorname{Sn}^2$$
 Sn^4 $\operatorname{\bar{2e}}$, E 0.15V

Multiplying equation (i) by 2 and (ii) by 5 we get:

$$Sn^{2+} + 2MnO_4 + 16H^+$$
 5 $Sn^{4+} + 8H_2O + 2Mn^{2+}$
 $E_{cell} = E_R \quad E_L$
= 1.51 - 0.15 = 1.36 V

Since E_{cell} is positive, therefore it will favour product formation.

- **12.** (i) **Tyndall effect:** Tyndall (1869) observed that when a beam of light is passed through a colloidal solution whose particle size is comparable to the wave length of light, the path of the **beam** is illuminated. This phenomenon is called Tyndall effect.
 - (ii) Shape Selective Catalyst: The catalysis reaction which depends upon the pore size of the catalyst and the size of the reactant and product molecules is called shape-selective catalysis. Zeolites are good shape-selective catalysts because of their honey comb-like structures. ZSM-5 used in petroleum industry to convert alcohols directly into gasoline by dehydrating them to give a mixture of hydrocarbons.

$$x \text{ CH}_3 - \text{OH}$$
 ZSM-5 $(\text{CH}_2)_x \quad x\text{H}_2\text{O}$ Gasoline

13. The colloidal particles bear similar electric charge and that is why they stay away from each other due to repulsion. If the charge is removed, they come together and are precipitated.

The precipitation and settling down of the discharged particles is called coagulation or flocculation.

The coagulation of a **lyophobic** sols can be done by mixing of two opposite charged sols or by addition of electrolyte.

14. (*i*) $I_2 + 10HNO_3$


$$2HIO_3 + 10NO_2 + 4H_2O$$


(ii) 3HgCl₂ + 2PH₃

$$Hg_3P_2 + 6HC1$$

(ii)

15. (*i*)





Xe F₄ (Square planar)

16. (*i*) To distinguish Ethylamine and aniline.

Chemical Test	C ₂ H ₅ NH ₂	C ₆ H ₅ NH ₂
Azo dye Test: Add NaNO ₂ + dil. HCl at 273 - 278K followed by NaOH and -naphthol	No reaction	Yellow orange dye is obtained. NH_2 $N=N-Cl$ $N=N-C$

(ii) Aniline and Benzyl amine

Chemical Test	Aniline	Benzyl amine
Test: Add nitrous acid (NaNO ₂ /HCl) to both the samples.	No evolution of N ₂ gas	N_2 gas is evolved CH_2NH_2 CH_2OH (i) $HONO(ii) H_2O + HCI$

- 17.
- (i) CH₃CH₂Cl NaCN
- CH₃CH₂CN Ethyl cyanide (A)
- reduction Ni/H₂
- CH₃CH₂CH₂NH₂ n-Propyl amine (B)

142 Xam idea Chemistry—XII

- 18. (i) PVC structure of monomer: $CH_2 = CH Cl$ (vinylchloride)
 - (ii) Teflon structure of monomer:

$$F \subset C \subset F$$

(Tetrafluoroethylene)

19. Given, Radius
$$(r) = 127.8 \text{ pm}$$

For fcc a $2\sqrt{2}r$

$$a^3 \quad (2\sqrt{2}r)^3 \quad 4.723 \quad 10^{-23} \text{ cm}^3$$

Atomic mass of $Cu = 63.54 \text{ g mol}^{-1}$

$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$

Density =
$$8.95 \text{ g cm}^{-3}$$

$$d \quad \frac{Z \quad M}{N_A \quad a^3}$$

$$8.95 = \frac{Z \quad 63.54}{6.023 \quad 10^{23} \quad 4.723 \quad 10^{-23}}$$

$$Z = \frac{8.95 - 6.022 - 4.723}{63.54}$$

$$Z = 4.00$$

Thus copper has fcc structure.

20. Given:
$$M_B = 58.5 \text{ g mol}^{-1}$$

$$W_A = 65g, W_B = ?$$

$$T_f = 7.5^{\circ}C \text{ or } 7.5 \text{ K},$$

$$K_f = 1.86 \text{ K kg mol}^{-1}$$

$$i = 1.87$$

We know that,

$$T_f = i K_f \times \frac{W_B}{M_B} \quad \frac{1000}{W_A}$$

$$7.5 = 1.87 \times 1.86 \times \frac{W_B}{58.5} \quad \frac{1000}{65}$$

$$W_B = \frac{7.5 \quad 58.5 \quad 65}{1.87 \quad 1.86 \quad 1000}$$

$$W_B = \frac{28518.75}{3478.2}$$

$$W_B = 8.199 \text{ g.}$$

21. (i) NaCN is used in the leaching of argentite (Ag₂S). Argentite is leached with dilute aqueous solution of NaCN in the presence of air

$$Ag_2S + 4NaCN$$
 2 Na $[Ag(CN)_2] + Na_2S$.

(ii) Iodine in the refining of titanium.

The crude metal is heated with iodine in an evacuated vessel to form volatile compound.

$$Ti (impure) + 2I_2$$
 TiI_4

Metal iodide decomposes on heating at 1800°C on a **tungston** filament. The pure metal deposit on the filament.

$$TiI_4$$
 1800 C $Ti(pure) + 2I_2$

(iii) Cryolite in the metallurgy of aluminium: In the metallurgy of Al, purified Al₂O₃ is mixed with cryolite (Na₃AlF₆) which lowers the melting point of the mixture and increase the conductivity.

OR

- (i) Froth floatation method: This method is used to remove gangue from sulphide ores. In this process a suspension of the powdered ore is made with water, some pine oil (collector) and stablisers are added. A rotating paddle agitates the mixture and draws air in it. The froth is light and it skimmed off.
- (ii) Electro refining of metals: In this method, the impure metal is made to act as anode. A strip of the same metal in pure form is used a cathode. They are put in a suitable electrolyte containing soluble salt of the same metal.

Anode: $M M^n ne^-$

Cathode: M^n ne^- M (pure metal)

Sb, Ag, Au, Pt, Cu, Zn may be refined by this process.

- (iii) Zone refining: This method is based on the principle that the impurities are more soluble in the melt than in the solid state of the metal. A circular mobile heater is fixed at one end of a rod of the impure metal. The molten zone moves along with the heater. The pure metal crystallises out of the melt and impurities pass on into the adjacent molten zone.
- 22. (i) Nickel does not form low spin octahedral complexes because 'Ni' has electronic configuration $3d^8 4s^2$, in which two inner *d*-orbitals are not available which are required to form d^2sp^3 hybridization.
 - (ii) The -complexes are known for the transition metals only because they have 'sandwich' structure in which the metal ion lies between two planar C₅H₅ rings. The bonding involves overlap of -electrons of the C₅H₅ rings with unfilled d-orbital of the metal. So that all M-C bonds are identical for their stability.
 - (iii) Co²⁺ is easily oxidised to Co³⁺ in the presence of a strong ligand because in presence of strong ligand, the 3d electrons pair up leaving two orbitals empty to be involved in d²sp³ hybridisation.

144 Xam idea Chemistry—XII

S_N1 Reactions: They are substitution nucleophilic unimolecular reactions in which the rate of the reaction depends on the concentration of only one reactant. Product formation takes place by the formation of carbo cation as reaction intermediate. For example:

$$CH_{3} \xrightarrow{C} CH_{3} \xrightarrow{CH_{3}} CH_{3} CH_{3} \xrightarrow{CH_{3}} CH_{3} CH_{3} CH_{3} CH_{3} CH_{3} C$$

 S_N2 reactions they are substitution nucleophilic bimolecular reactions in which rate of the reaction depends on the concentration of two reactants. For example:

$$R-CI$$
 OH $R-OH$ CI
Rate $[R-CI][OH^-]$

The product formation takes place by formation of transition state.

24. (*i*)

(ii)

(iii)
$$CH_3-CH=CH_2 \xrightarrow{\text{dil } H_2SO_4} CH_3-CH-CH_3$$
Propene Propan-2-ol

- 25. (i) This is because of (a) bond dissociation enthalpy of F₂ is lower than Cl₂ and (b) fluorine forms stronger bond with nitrogen due to comparable size.
 - (ii) As xenon is sp^3 hybridised with three lone pair of electrons.
 - (iii) This is due to (a) low bond dissociation enthalpy of F_2 and (b) high hydration enthalpy of F.
- 26. This is due to presence of carboxylic acid as well as amino groups.
 - Amino acids which contain two —C—O—H group and one —NH₂ group are called acidic amino acid, e.g., aspartic acid.

- $\bullet \quad \text{Amino acids which contain two } -\text{NH}_2 \text{ group and one } -\text{C}-\text{O}-\text{H group are called basic amino acids, } e.g., \\ \text{lysine.}$
- Amino acids which contain one —C—O—H and one —NH₂ group are called neutral amino acids, e.g., glycine.
- Amino acids that cannot be synthesized by the body and must be supplied in the diet are called essential amino acids, *e.g.*, lysine, valine, leucine, etc.
- The amino acids which are synthesized by our body are called non-essential amino acids, *e.g.*, alanine, glycine, etc.
- 27. (i) Food preservatives: Chemical substance which are used to prevent food spoilage due to microbial growth are called preservative. For example, sodium benzoate, potassium sorbate, etc.
 - (ii) Enzymes: Enzymes are the essential biological catalysts which catalyse specific biological reactions at a very high rate under mild conditions of temperature and pH. For example, invertase, zymase etc.
 - (iii) Detergents: They are very similar to the salts of fatty acids found in soap but they are manufactured from materials other than animal fats. For example: Sodium alkylbenzene sulphonates are common ingredients of synthetic detergents.
- 28. (a) (i) Rate of Reaction: It is defined as the decrease in the concentration of reactants per unit time or increase in the concentration of products per unit time.

For a hypothetical reaction R P
$$Rate = \frac{-[R]}{t} Rate = \frac{[P]}{t}$$

(ii) Activation energy: The amount of energy which the reactants must absorb to pass over the energy barrier between reactants and products is known as the activation energy.

(b) (i) We know that
$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]t}$$

$$t_{1/2} = \frac{0.693}{k}$$

$$k = \frac{0.693}{t_{1/2}}$$

$$k = \frac{0.693}{37.9}$$

$$k = 0.0183 \text{ s}^{-1}$$

$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$$

$$t = \frac{2.303}{0.0183} \log \frac{1}{1/4}$$

$$= \frac{2.303}{0.0183} \log 2^2 \quad \frac{2.2303}{0.0183} \log 2$$

$$t = \frac{2 \quad 2.303}{0.0183} \times 0.3010$$

$$t = 75.8 \text{ s}.$$

(ii) 1 minute = 60 second

$$t = \frac{2.303}{k} \log \frac{[A]_o}{[A]}$$
$$60 = \frac{2.303}{0.0183} \log \frac{[A]_o}{[A]}$$
$$60 = 0.0183 \qquad [A]_o$$

$$\frac{60 \quad 0.0183}{2.303} = \log \frac{[A]_0}{[A]}$$

Taking antilog,

$$0.4767 = \log \frac{[A]_0}{[A]}$$

Antilog (0.4767) =
$$\frac{[A]_o}{[A]}$$

$$2.997 = \frac{[A]_o}{[A]}$$

$$\frac{[A]}{[A]_o} = \frac{1}{2.997} = 0.333 = 0.333\%$$

Hence, amount left out = 33.3%

- (a) (i) The order of a reaction can be defined as the sum of the powers of the concentration terms as expressed in rate law.
 - (ii) **Molecularity:** It may be defined as the number of reacting species (molecules, atoms or ions) taking part in an elementary reaction, which must collide simultaneously in order to bring about a chemical reaction.

(b) (i) Given
$$T_1 = 300 \text{ K} \\ T_2 = 320 \text{ K} \\ R = 8.134 \text{ JK}^{-1} \text{ mol}^{-1} \\ E_a = ?$$

We know that,
$$E_a = 2.303 \text{ R} \frac{T_1 T_2}{T_2 T_1} \log \frac{k_2}{k_1}$$

 $E_a = 2.303 \times 8.314 \frac{300 320}{20} \log 4$
 $E_a = 19.15 [4800] \times 0.602$
 $= 55.32 \text{ kJ mol}^{-1}$.

29. (a) (i)
$$Cr_2O_7^{2-} + 3H_2S + 8H^+$$
 $2Cr^{3+} + 3S + 7H_2O$

(*ii*)
$$2Cu^{2+} + 4I^{-}$$
 $Cu_2I_2 + I_2$

- (b) (i) The oxidising power of oxonions are in the order $VO_2^+ < Cr_2 O_7^{2-} < MnO_4^-$, this is due to increase in the oxidation state of the metal ion.
 - (ii) The third ionisation enthalpy of manganese (Z = 25) is exceptionally high because Mn²⁺ ion has $3d^5$ configuration which is highly stable since it is half-filled.
 - (iii) Cr^{2+} stronger reducing agent than Fe^{2+} as its configuration changes from d^4 to d^3 , a more stable half filled t_{2g} configuration.

OR

(a) (i)
$$8\text{Mn O}_4$$
 (aq) $3\text{S}_2\text{O}_3^2$ (aq) H_2O 8MnO_2 6SO_4^2 2OH

(ii)
$$Cr_2O_7(aq)$$
 6Fe² (aq) 14H 2Cr³ 6Fe³ 7H₂O.

- (b) (i) La³⁺ and Lu³⁺ ions do not show any colour in solution because they do not contain any unpaired electrons.
 - (ii) Among the divalent cations in the first series of transition elements, manganese exhibits the maximum paramagnetism because manganese (Mn²⁺) ion has maximum number of 5 unpaired electrons.
 - (iii) Cu^+ ion is not known in aqueous solutions because Cu^{2+} ions are more stable due to more negative H_{hyd} of Cu^{2+} than Cu^+ , which compensates for the second ionisation enthalpy of Cu.

30. (a) (i) Clemmensen reaction

$$\begin{array}{c|c}
CH_3 & C & CH_3 & CH_3 & CH_3 & CH_3 & CH_2 & CH_3 \\
\hline
 Acetone & Propane
\end{array}$$

(ii) Cannizzaro's reaction:

(ii)
$$C_2H_5$$
 COOH $KMnO_4/OH^-$

Ethyl benzene Benzoic acid

(iii) Bromobenzene CN COOH
$$\begin{array}{c} CuCN \\ Py,473K \end{array} \longrightarrow \begin{array}{c} H_2O/H^+ \\ Benzoic acid \end{array}$$

OR

(a) (i) Hell-Volhard-Zelinsky reaction

Carboxylic acid

Halocarboxylic acid

(ii) Wolf-Kishner reduction

(iii)
$$(iii)$$
 (ii) (ii) (ii) (ii) (ii) (iii) $(ii$

CBSE (All India) SET-II

- 1. *n*-type semiconductor.
- 2. Due to presence of triple bond whose dissociation energy is very high.
- **6.** They are the simplest carbohydrates which do not undergo further hydrolysis, for example: glucose.
- **13.** (*i*) **Peptization:** The process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of a suitable electrolyte.
 - (ii) Reversible sols: These are the sols in which dispersion medium can be separated from dispersed phase and reconstituted easily.

$$(ii) \quad \text{XeF}_6 \quad 3\text{H}_2\text{O} \qquad \quad \text{XeO}_3 \quad 6\text{HF}$$

17. (i) To distinguish $CH_3 NH_2$ and $CH_3 - NH - CH_3$

Reaction with HONO

CH₃NH₂ gives methyl alcohol and N₂ gas.

$$\text{CH}_3\text{NH}_2$$
 Hono $^{4\text{ C}}$ CH_3OH N_2 H_2O

Dimethyl amine forms nitroso amine which is water insoluble yellow oil.

$$(CH_3)_2 N - N O H_2 O$$
N-Nitroso

dimethyl amine (yellow oil)

(iii) Aniline and N-methyl aniline

Test: Carbyl amine test:

Aniline on warming with chloroform and **KOH** gives offensive smell of **carbylamine** while N-methyl aniline does not.

$$C_6H_5NC$$

Phenyl carbylamine

18. (i) Bakelite

Monomers: Phenol and formaldehyde

Structure:

and

Formaldehyde

(ii) Nylon-6

Monomers: Caprolactum

$$H_2C$$
 N $C = O$

$$\begin{array}{ccc} \operatorname{H}_2 \operatorname{C} & \operatorname{CH}_2 \\ \operatorname{H}_2 \operatorname{C} & \operatorname{CH}_2 \end{array}$$

Caprolactum

19. Atomic mass (M) = 108 g mol^{-1} , a = 409 pm

We know that for fcc lattice:

$$r = \frac{a}{2\sqrt{2}}$$

$$r = \frac{409}{2 \cdot 1.414} = \frac{409}{2.828}$$

$$r = 144.62 \text{ pm}$$

20.
$$M_B = 62.0 \text{ g mol}^{-1}$$
, $W_A = 5.50 \text{ kg or } 5500 \text{ g}$

$$K_f = 1.86 \text{ K kg mol}^{-1}, \quad T_f = 10 \text{ K}$$

We know that,

$$T_f = K_f \times \frac{W_B}{M_B} \quad \frac{1000}{W_A}$$

$$10 = 1.86 \times \frac{W_B}{62} \quad \frac{1000}{5500}$$

$$W_B = \frac{62 \quad 5500 \quad 10}{1000 \quad 1.86}$$

$$W_B = \frac{3410000}{1860} = 1833.3 \text{ g}$$

- 27. Analgesics: They are the drugs used for relieving pain. Analgesics are classified into two categories:
 - (a) Narcotics (addictive) drugs
- (b) Non-narcotics (non-addictive) drugs
- (a) Narcotics: Narcotic drugs are recommended for cardiac pain, post operative pains, terminal cancer and in child birth.
- (b) Non-narcotics: Non-narcotic drugs are effective in relieving skeletal pain, reducing fever, preventing heart attack.

CBSE (All India) SET-III

- 1. Metallic solids have high melting and boiling points.
- 3. Molality of a solution does not change on changing the temperature of the solution while molarity does change.

8.
$$C_{12}H_{22}O_{11}$$
 H_2O H $C_6H_{12}O_6$ $C_6H_{12}O_6$ (Glucose) (Fructose)

19. For fcc, Z = 4

$$a = 4.07 \times 10^{-8}$$
 cm, d = 10.5 g cm⁻³, M = atomic mass = ?, N_A = 6.02×10^{23}

We know that,

$$d = \frac{Z \quad M}{N_A \quad a^3} \qquad M \quad \frac{d \quad N_A \quad a^3}{Z}$$

$$M = \frac{10.5 \quad 6.022 \quad 10^{23} \quad (4.07 \quad 10^{-8})^3}{4}$$

$$M = 106.5 \text{ g mol}^{-1}$$

20. Given:
$$W_B = 15 \text{ g}, W_A = 450 \text{ g}$$

$$T_f = [273 - (273 - 0.34)]K = 0.34 \text{ K}$$

$$M_B = ?, K_f = 1.86 \text{ K kg mol}^{-1}$$

$$M_B = K_f \times \frac{W_B - 1000}{T_f - W_A}$$

= 1.86 K kg mol⁻¹ ×
$$\frac{15 \text{ g}}{0.34 \text{ K}} = 182.3 \text{ g mol}^{-1}$$

- 22. (i) This is due to smaller size of oxygen the electron cloud is distributed over a small region of space, making electron density high which repels the incoming electrons.
 - (ii) Because P—P bond is stronger than N—N bond.
 - (iii) Fluorine never acts as the central atom in polyatomic inter-halogen compounds since it is the most electronegative element of the group.
- 25. (i) Tetrachloridocobaltate (II) ion

sp³ hybridisation

Tetrahedral

Paramagnetic

(ii) Tetracyanonickelate (II) ion

dsp² hybridisation

Square plannar

Diamagnetic.

(iii) Diaquadioxalatochromate (III) ion

 d^2sp^3 hybridisation

Octahedral

Paramagnetic.

26.

Fibrous Proteins	Globular Proteins
1. Consist of linear thread-like molecules which tend to lie side by side to form fibre like structure.	
2. Insoluble in water.	2. Soluble in water.
3. Keratin in hair, fibroin in silk, etc.	3. Albumin in eggs, insulin, etc.

Denaturation of Proteins

When a protein in its native form, is subjected to a change in temperature or change in pH, the hydrogen bond are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called denaturation of protein. During denaturation 2° and 3° structures are destroyed but 1° structures remain intact, e.g., coagulation of egg white on boiling, curdling of milk, etc.

27. (i) Antibiotics

The chemical substances produced from some micro-organism and are used to inhibit the growth of other microorganism or even kill them are called antibiotics. For example, penicillin, chloramphenicol.

(ii) Antiseptics

Antiseptics are the chemicals which prevent or destroy the growth of the harmful microorganism. For example: Savlon, dettol.

(iii) Analgesics

These are the substances which are used to get relief from pain. For example: Aspirin.

CBSE EXAMINATION PAPERS

FOREIGN-2010

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Foreign) SET-I

- 1. What is the number of atoms in a unit cell of a simple cubic crystal?
- 2. Identify the order of reaction from the following unit for its rate constant:

 $L \text{ mol}^{-1} \text{s}^{-1}$

- **3.** Give an example of shape-selective catalyst.
- **4.** Which is a stronger acid in aqueous solution—HCl or HI, and why?
- **5.** What is an ambidentate ligand? Give an example.
- **6.** Draw the structure of the following compound:

4-Bromo-3-methylpent-2-ene

7. Give IUPAC name of the following compound:

- **8.** What happens when glucose is treated with bromine water?
- **9.** Write the anode and cathode reactions occurring in a commonly used mercury cell. How is the overall reaction represented?
- **10.** Define:
 - (i) Elementary reaction in a process
 - (ii) Rate of a reaction.
- 11. The rate constant for a zero order reaction is 0.0030 mol L⁻¹ s⁻¹. How long will it take for the initial concentration of the reactant to fall from 0.10 M to 0.075 M?
- **12.** Draw the structures of the following molecules:
 - (i) BrF₃

- (ii) XeOF₄
- **13.** Describe the preparation of
 - (i) Potassium dichromate from sodium chromate, and
 - (ii) KMnO₄ from K₂MnO₄

- **14.** Describe the state of hybridization, the shape and the magnetic behaviour of the following complexes:
 - (i) $[Cr(H_2O)_2(C_2O_4)_2]^-$
 - (ii) $[Co(NH_3)_2(en)_2]^{3+}$, (en = ethane-1,2-diamine)

(At. Nos :
$$Cr = 24$$
, $Co = 27$)

OR

Explain the following terms:

- (i) Crystal field splitting in an octahedral field
- (ii) Spectrochemical series.
- **15.** Write the mechanisms of the following reactions:
 - (i) Hydration of ethene to ethanol
 - (ii) Dehydration of ethanol, giving ethene.
- **16.** Illustrate the following name reactions giving chemical equations:
 - (i) Reimer-Tiemann reaction
 - (ii) Williamson synthesis.
- **17.** Explain the following giving a reason in each case:
 - (i) Why is an alkylamine more basic than ammonia?
 - (ii) Why do primary amines have higher boiling points than the tertiary amines?
- **18.** Complete the following reaction equations:
 - (i) $C_6H_5NH_2 + CHCl_3 + KOH$ (alc)
 - $(ii) C_6H_5N_2Cl + H_3PO_2 + H_2O$
- 19. Iron has a body-centred cubic unit cell with a cell edge of 286.65 pm. The density of iron is 7.874 g cm^{-3} . Use this information to calculate Avogadro's number. (At. mass of iron = 56 g mol⁻¹)
- 20. One half-cell in a voltaic cell is constructed from a silver wire dipped in silver nitrate solution of unknown concentration. The other half-cell consists of a zinc electrode in a 0.10 M solution of Zn (NO₃)₂. A voltage of 1.48 V is measured for this cell. Use this information to calculate the concentration of silver nitrate solution.

(Given
$$E_{Zn^2/Zn} = -0.763 \text{ V}, E_{Ag^2/Ag}$$
 0.80 V)

- 21. What is the difference between multimolecular and macromolecular colloids? Give one example of each type. How are associated colloids different from the above two types of colloids?
- **22.** Explain the following observations:
 - (i) The enthalpies of atomisation of transition metals are quite high.
 - (ii) There is a close similarity in physical and chemical properties of the 4 d and 5 d series of the transition elements, much more than expected on the basis of usual family relationship.
 - (iii) The members in the actinoid series exhibit larger number of oxidation states than the corresponding members in the lanthanoid series.
- **23.** Answer the following:
 - (i) Identify chiral in CH₃CHOHCH₂CH₃ and CH₃CHOHCH₃.

(ii) Among the following compounds, which one is more easily hydrolysed and why?

CH₃CHCICH₂CH₃ or CH₃CH₂CH₂CH₂Cl

(iii) Which of these will react faster in S_N² displacement and why?

1-bromopentane or 2-bromopentane

- **24.** Describe how the following changes are brought about:
 - (i) Pig iron into steel
- (ii) Bauxite into pure alumina
- (iii) Impure copper into pure copper.
- **25.** Explain the meanings of the following terms:
 - (i) Invert sugar

- (ii) Peptide linkage
- (iii) Denaturation of proteins.

OR

Mention three such facts/reactions about glucose which cannot be explained by its open end structure. What is meant by pyranose structure of glucose?

- **26.** Identify the four groups into which the polymers are classified on the basis of the magnitude of intermolecular forces present in them. To which group or groups do polythene and bakelite belong?
- 27. How do antiseptics differ from disinfectants? Name a substance that can be used as an antiseptic as well as a disinfectant.
- **28.** (*a*) Non-ideal solutions exhibit either positive or negative deviations from Raoult's law. What are these deviations and how are they caused?
 - (b) What mass of NaCl (molar mass = 58.5 g mol^{-1}) must be dissolved in 65g of water to lower the freezing point by 7.50°C ? The freezing point depression constant, K_f for water is $1.86 \text{ K kg mol}^{-1}$. Assume van't Hoff factor for NaCl is 1.87.

- (a) Define the terms osmosis and osmotic pressure. What is the advantage of using osmotic pressure as compared to other colligative properties for the determination of molar masses of solutes in solution?
- (b) A solution prepared from 1.25 g of oil of wintergreen (methyl salicylate) in 99.0 g of benzene has a boiling point of 80.31°C. Determine the molar mass of this compound. (Boiling point of pure benzene = 80.10°C and K_b for benzene = 2.53°C kg mol⁻¹).
- **29.** (a) Complete the following chemical reaction equations:
 - (i) $HgCl_2(aq) + PH_3(g)$
 - (ii) $SiO_2(s) + HF(g)$
 - (b) Explain the following observations:
 - (i) Sulphur in vapour state exhibits paramagnetic behaviour.
 - (ii) The stability of +3 state increases down the group in group 15 of the periodic table.
 - (iii) XeF₂ has a linear shape and not a bent structure.

OR

- (a) Complete the following chemical reaction equations:
 - (i) $AgCl(s) + NH_3(aq)$
 - (ii) $P_4(s) + NaOH(aq) + H_2O(l)$
- (b) Explain the following observations:
 - (i) H₂S is less acidic than H₂Te
 - (ii) Fluorine is a stronger oxidising agent than chlorine
 - (iii) Noble gases are the least reactive elements.
- **30.** (a) How will you prepare the following compounds starting with benzene?
 - (i) Benzaldehyde
 - (ii) Acetophenone.
 - (b) Give chemical tests to distinguish between the following pairs of compounds:
 - (i) Ethanal and Propanal
 - (ii) Benzaldehyde and Acetophenone
 - (iii) Phenol and Benzoic acid.

OR

- (a) Explain the mechanism of nucleophilic attack on a carbonyl group of an aldehyde or a ketone.
- (b) How would you bring about the following conversions?
 - (i) Propanone to Propene
 - (ii) Ethanol to 3-hydroxybutanal
 - (iii) Benzaldehyde to Benzophenone.

CBSE (Foreign) SET-II

Questions uncommon to Set I

- 1. State a feature to distinguish a metallic solid from an ionic solid.
- 2. Which one of these compounds is more easily hydrolysed by KOH solution and why?

CH₃ CHCICH₂CH₃

Or

CH₃CH₂CH₂CH₂Cl

- **9.** Draw the structures of O_3 and S_8 molecules.
- **14.** How are the following conversions carried out?
 - (i) Propene to Propan-2-ol
 - (ii) Ethyl magnesium chloride to Propan-1-ol
- 19. Silver crystallises in a fcc lattice. If the edge length of the unit cell is 4.07×10^{-8} cm and density of silver is 10.5 g cm⁻³, calculate the atomic mass of silver. (NA = 6.02×10^{23} atoms mol⁻¹)
- **20.** A copper-silver cell is set up. The copper ion concentration is 0.10 M. The concentration of silver ions is not known. The cell potential was found to be 0.422 V. Determine the concentration of silver ions in the cell.

(Given
$$E_{Ag/Ag} = 0.80 V$$
, $E_{Cu^2/Cu} = 0.34 V$)

156 Xam idea Chemistry—XII

- 21. State what is observed when
 - (i) an electrolyte NaCl, is added to hydrated ferric oxide sol.
 - (ii) an electric current is passed through a colloidal solution.
 - (iii) a beam of light is passed through a colloidal solution.
- **26.** What are biodegradable and non-biodegradable detergents? Give one example of each. Is there any structural difference between the two?
- **27.** Write the structure of the monomer of each of the following polymers:
 - (i) Polyvinyl chloride
 - (ii) Teflon
 - (iii) Bakelite.

CBSE (Foreign) SET-III

Ouestions uncommon to Set I and Set II

- 2. Identify the order of reaction for which the rate constant is expressed in units of L^{-1} mol s^{-1} .
- **3.** What is meant by 'reversible sols'?
- **4.** Give an example of coordination isomerism.
- **8.** What happens when glucose reacts with nitric acid?
- **12.** Draw the structure of the following molecules:
 - (i) XeF₂
 - (ii) HClO₄
- **16.** Illustrate the following name reactions giving a chemical reaction equation for each:
 - (i) Kolbe's reaction of phenol
 - (ii) Friedel-Crafts' acetylation of anisole.
- 19. The well-known mineral fluorite is calcium fluoride. It is known that in the unit cell of this mineral there are 4 Ca^{2+} ions and $8F^-$ ions and the Ca^{2+} ions are arranged in a face-centred cubic lattice. The F^- ions fill all the tetrahedral holes in the fcc lattice of Ca^{2+} ions. The edge of the unit cell is 5.46×10^{-8} cm in length. The density of the mineral is 3.18 g cm^{-3} . Use this information to calculate Avogadro's number. (Molar mass of $CaF_2 = 78.08 \text{ g mol}^{-1}$)
- **20.** A voltaic cell is set up at 25°C with the following half-cells, Al³⁺ (0.0010 M) and Ni²⁺ (0.50 M). Write an equation for the reaction that occurs in the cell when the cell generates an electric current and determine the cell potential.

(Given E
$$_{\text{Ni}^2}$$
 /Ni 0.25V, E $_{\text{Al}^3}$ /Al 1.66V)

- **21.** Write three features of chemisorption which are not found in physisorptions. Illustrate your answer with suitable examples.
- **22.** Explain the following:
 - (i) The transition elements have great tendency for complex formation.
 - (ii) There is a gradual decrease in the atomic sizes of transition elements in a series with increasing atomic numbers.
 - (iii) Lanthanum and Lutetium do not show colouration in solutions. (At. No.: La = 57, Lu = 71)



CBSE (Foreign) SET-I

- 1. One atom.
- 2. Second order
- **3.** ZSM-5
- 4. HI is stronger acid then HCl because dissociation energy of H–I bond is less than that of H–Cl.
- 5. An ambidentate ligand is one which can link through either of the two donor atoms to the central metal ion. For example: NO_2^- and \overline{O} NO, \overline{CN} and \overline{NC} .

6.
$$CH_3 - CH - CH = CH - CH_3$$

$$CH_3 - CH - CH = CH - CH_3$$

$$CH_3$$

- 7. 4-Methyl-pent-3-en-2-one
- **8.** When glucose is treated with bromine water it forms gluconic acid.

9. Chemical reactions of mercury cell

Anode:
$$Zn(s)$$
 $2OH^ ZnO(s)$ H_2O $2e^ Cathode: HgO H_2O $2e^ Hg(l)$ $2OH^-$$

The overall reaction is represented by

$$Zn(s)$$
 HgO(s) $ZnO(s)$ + Hg(l)

- **10.** (*i*) Some reactions occur by a series of elementary steps and such simple steps are known as elementary reactions in a process.
 - (ii) Rate of a reaction: The change in the concentration of reactants products of or per unit time is called rate of the reaction.


Rate =
$$\frac{R}{t}$$
 or $+\frac{P}{t}$

11. Given:
$$[R_0] = 0.10 \text{ M}$$
 $[R] = 0.075 \text{ M}$ $k = 0.0030 \text{ mol } L^{-1} \text{ s}^{-1}, t = ?$


We know that
$$k = \frac{[R_o] - [R]}{t}$$

or $t = \frac{[R_o] [R]}{k} = \frac{0.10 \ 0.075}{0.0030} = 8.33 \text{ s.}$

12. (i) Structure of BrF₃ molecule



(ii) Structure of XeOF₄ molecule.



Square pyramidal

- 13. (i) Preparation of potassium dichromate from sodium chromate
 - (a) $2Na_2CrO_4$ H_2SO_4

(b) Na₂CrO₇ 2KCl K₂Cr₂O₇ 2NaCl

(ii) Preparation of KMnO₄ from K₂MnO₄

$$2K_2MnO_4 + Cl_2$$

$$2KMnO_4 + 2KCl$$

OR

$$2K_2MnO_4 + H_2O + O_3$$
 $2KMnO_4 + 2KOH + O_2$


- **14.** (*i*) $[Cr (H_2O)_2 (C_2O_4)_2]^-$
 - (a) Hybridization d^2sp^3
 - (b) Shape octahedral
 - (c) Magnetic behaviour Paramagnetic
 - (ii) $[\text{Co (NH}_3)_2 (\text{en})_2]^{3+}$
 - (a) Hybridisation d^2sp^3
 - (b) Shape Octahedral
 - (c) Magnetic behaviour diamagnetic

OR

(i) Crystal field splitting in an octahedral field.

According to C.F.T at first there is an increase in the energy of d-orbitals relative to that of the free ion just as would be the case in spherical field. The two orbitals lying along the axis get repelled more strongly than other three (d_{xy}, d_{yz}, d_{xz}) . The degenerate set of d-orbitals get split into two sets: the lower energy t_{2g} set and e_g set. The energy separation is denoted by o. The actual configuration adopted is decided by the relative values of on and P. P-represents the energy required for electron pairing in a single orbital.

- (a) If $_{0}$ < P then fourth electron goes to e_{q}
- (b) When $_{0} > P$, then fourth electron pairs the t_{2g} orbital.



(ii) Spectrochemical Series: It is a series in which ligands can be arranged in the order of increasing field strength or in order of increasing magnitude of splitting they produce.

The order is

I Br SCN Cl S 2 F OH $C_2O_4^2$ $H_2O < NCS < edta$ $^4 < NH_3$ < en < CN CO

15. (*i*) Mechanism of hydration of ethene to ethanol


Step-I: Protonation of alkene

Step-II: Nucleophilic attack of water as carbocation

Step-III: Deprotonation of alcohol

(ii) Mechanism of dehydration of ethanol to ethene.


Step-I: Formation of protonated alcohol



Step-II: Formation of carbocation

Setp-III. Formation of ethene


16. (i) Reimer-Tiemann reaction



(ii) Williamson synthesis

$$C_2H_5ONa + ICH_3$$
 $C_2H_5OCH_3 + NaI$
Sodium ethoxide Methyl-ethyl ether

17. (i) Alkylamine is more basic than ammonia because in aliphatic amines, the electron releasing alkyl group stabilize their ammonium cations by dispersing the positive charge and in parent amine make the nitrogen unshared electrons more available for sharing with a proton.



(ii) The boiling points of primary amine are higher than the tertiary amines because strong intermolecular hydrogen bonding takes place between the molecules of primary amine.

19. **Given:**
$$Z = 2$$

$$a = 286.65 \text{ pm} \qquad d = 7.874 \text{ g cm}^{-3}$$

$$M = 56 \text{ g mol}^{-1} \qquad N_A = ?$$

$$d = \frac{ZM}{(a^3) N_A}$$

$$7.874 = \frac{2 \quad 56}{(286.65)^3 \quad 10^{-30} N_A}$$

$$N_A = \frac{2 \quad 56}{7.874 \quad (286.65)^3 \quad 10^{-30}} = \frac{112}{18.5 \quad 10^{-23}} \quad \frac{112 \quad 10^{23}}{18.5}$$

$$N_A = 6.054 \times 10^{23}$$

20. Electrochemical cell

$$Zn(s)|Zn^{2+} (.10 \text{ M})|Ag^{+} (conc.)|Ag (s)$$

 $E_{cell}^{o} = E_{R}^{o} E_{L}^{o}$
 $= 0.80 (-0.763)V = 1.563 \text{ V}$

We know that

$$E_{cell}^{o} = (E_{R}^{o} - E_{L}^{o}) - \frac{0.0591}{n} \log \frac{[Zn^{2}]}{[Ag]^{2}}$$

$$1.48 = 1.563 - \frac{0.0591}{2} \log \frac{[0.10]}{[Ag^{+}]^{2}}$$

$$\log \frac{0.10}{[Ag]^{2}} = \frac{0.083}{0.02955} = 2.8087$$
Or
$$\frac{0.10}{[Ag]^{2}} = \text{antilog of } 2.8087 = 643.7$$

$$[Ag^{+}]^{2} = \frac{0.10}{643.7} = 1.553 \times 10^{-4}$$

$$[Ag^{+}] = 1.247 \times 10^{-2} \text{ M}$$

21. (i) Macro molecular colloids: Macromolecules in suitable solvents form solution. Where in size of the macromolecules may fall within the colloidal range. The system showing such characteristics are called macromolecular colloids. The colloids are quite stable and in many respect they resemble true solutions. Examples of naturally occurring macro molecules are starch, cellulose, proteins and those of manmade macromolecules are nylon, polythene, polystyrene, synthetic rubber etc.

- (ii) Multimolecular colloids: A large number of atoms or smaller molecules (diameter < 1 nm) of a substance on dissolution aggregate together to form species having size in the colloidal range. Such species are called *multimolecular colloids*. Example: A sulphur sol contains thousands of S_8 sulphur molecules, a platinum or gold sol may have particles of various sizes having many atoms.
- (iii) Associated colloides (Micelles): There are certain substances which at low concentrations behave like normal strong electrolytes but at higher concentration behave as colloidal solutions due to the formation of aggregated particles. Such colloids are associated colloids and aggregated particles as micelles. Examples: soaps and detergents.
- **22.** (*i*) This is because transition metals have strong metallic bonds as they have large number of unpaired electrons.
 - (ii) There is a close similarity in physical and chemical properties of the 4d and 5d series of the transition elements much more than expected on the basis of usual family relationship. This is because 5d and 4d series elements have virtually the same atomic and ionic radii due to lanthanide contraction. Due to equality in size of Zr and Hf, Nb and Ta, Mo and W, etc., the two elements of pair have the same properties.
 - (iii) The members in the actinoid series exhibit larger number of oxidation states than the corresponding members in the lanthanide series due to the fact that the 5f, 6d and 7s levels are of comparable energies.

- (ii) CH₃CHClCH₂CH₃, in more easily hydrolysed as it forms secondary carbocation which is more stable than primary carbocation.
- (iii) 1-bromopentane will react faster in S_N² displacement reaction as it is a 1° alkyl halide.
- 24. (i) Pig iron into steel: Pig iron is converted into steel by heating in a converter. A blast of oxygen diluted with carbon dioxide is blown through the converter. Oxygen reacts with impurities and raised the temperature to 2173K carbon gets oxidised to CO which burns off at the mouth of the converter. Oxides of silicon and Mg form slag. When the flame is stopped, slag is tapped off and other metals like Mn, Cr, Ni, W may be added in the end.
 - (ii) Bauxite into pure alumina.

$$\begin{array}{lll} (a) & {\rm Al_2O_3}(s) + 2{\rm NaOH}(aq) + 3{\rm H_2O}(l) & 2{\rm Na~[Al~(OH)_4~]}(aq) \\ \\ (b) & 2{\rm Na~[Al~(OH)_4~]}(aq) + {\rm CO_2}(g) & {\rm Al_2O_3.}x{\rm H_2O}(s) + 2{\rm NaHCO_3}(aq) \\ \\ (c) & {\rm Al_2O_3.}x{\rm H_2O} & {\rm Al_2O_3}(s) & x{\rm H_2O}(g) \\ \end{array}$$

(iii) Impure copper into pure copper

Pure copper is obtained by electro-refining process. In this method, the impure metal is made to act as anode. A strip of the same metal in pure form is used as cathode. They are put in a suitable electrolyte bath containing soluble salt of the same metal.

Anode: Cu
$$Cu^2$$
 $2\bar{e}$ Cathode: Cu² $2\bar{e}$ Cu (Pure copper)

25. (*i*) **Invert Sugar:** The dextrorotatory sucrose when hydrolysed by boiling with mineral acid produces an equal number of molecules of dextrorotatory fructose. The resulting mixture is laevorotatory and termed as invert sugar.

0

- (ii) **Peptide linkage:** The amide linkage (— C NH —) formed between two —amino acid molecules with the loss of a water molecule in a polypeptide is called a peptide linkage.
- (iii) **Denaturation of Proteins:** When a protein in its native form, it is subjected to a change, change in temperature or change in pH, the hydrogen bond are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called denaturation of protein. During denaturation 2° and 3° structures are destroyed but 1° structures remain intact, *e.g.*, coagulation of egg white on boiling, curdling of milk, etc.

OR

The following reactions cannot be explained by the open chain structure of glucose.

- (i) Despite having the aldehyde group, glucose does not give 2, 4– DNP test, Schiffs test and it does not form the hydrogensulphite addition product with NaHSO₃.
- (ii) The penta acetate of glucose does not react with hydroxyl amine indicating the absence of free —CHO group.
- (iii) D-Glucose on treatment of methyl alcohol in the presence of dry HCl gas gives two isomeric nonomethyl derivatives known as -D-glucoside and methyl D-glucoside. These glucosides do not reduce fehling solution and also do not react with hydrogen cyanide indicating the absence of free —CHO group.

A ring structure called pyranose structure (- or -) is proposed for the glucose molecule.

- 26. Polymers are classified into four groups based on the magnitude of intermolecular forces. They are—
 - (i) Elastomers

- (ii) Fibers
- (iii) Thermoplastic polymers
- (iv) Thermosetting polymers

Polythene—Polythene is a thermoplastic polymer.

Bakelite— It is a thermosetting polymer.

27. Antiseptics: Antiseptics are the chemicals which prevent the growth of the harmful microorganism. They do not harm the living tissues. For example dettol.

Disinfectants: These are the chemicals which completely destroy the microorganism: They are toxic to living tissues. 1% solution of phenol is a disinfectant whereas 0.2% solution of phenol is antiseptic.

28. (a) **Positive deviation from Raoult's law:** When the partial vapour pressure of each component (A and B) consequently the total vapour pressure is greater than the pressure expected on the basis of Raoult's law then the deviation is termed as positive deviation.

Cause of positive deviation: This type of deviation is observed by solution in which the forces of attraction between A–A molecules and between B–B molecules is greater then the forces of attraction between A–B molecules.

Negative deviation from Raoult's law: When the partial vapour pressure of each component of solution is less than the vapour pressure expected on the basis of Raoult's law then the deviation is called as negative deviation.

Causes of negative deviation: This type of deviation is shown by solutions in which the forces of attraction between A–A and B–B molecules is less than the forces of attraction between A and B molecules.

A B A A Or B B

(b) Given:
$$M_B = 58.5 \text{ g mol}^{-1}$$
 $W_A = 65\text{ g}$, $W_B = ?$
 $T_f = 7.5^{\circ}\text{C}$ or 7.5 K ,

 $K_f = 1.86 \text{ K kg mol}^{-1}$
 $i = 1.87$

We know that,

$$T_f = iK_f \times \frac{W_B}{M_B} \frac{1000}{W_A}$$

$$7.5 = 1.87 \times 1.86 \times \frac{W_B}{58.5} \frac{1000}{65}$$

$$W_B = \frac{7.5 \quad 58.5 \quad 65}{1.87 \quad 1.86 \quad 1000}$$

$$W_B = \frac{28518.75}{3478.2}$$

$$W_B = 8.199 \text{ g.}$$

OR

(a) **Osmosis:** The phenomenon in which solvent molecules flow through a semipermeable membrane from a solution of low concentration to a solution of higher concentration is termed as osmosis.

Osmotic Pressure: The hydrostatic pressure exerted by the column of the solution which is just sufficient to prevent the osmosis is called osmotic pressure.

Advantage of using osmotic pressure as compared to other collegative properties:

- (i) The equilibrium is established very quickly. Hence, results are obtained in a very short time.
- (ii) The concentration of the solution does not change during determination of osmotic pressure.
- (iii) This is ideal method for proteins to find their molar mass since it happens at room temperature and no coagulation occurs.

(b) Given
$$T_B = 80.31 - 80.10 = 0.21 \text{ °C or } 0.21 \text{ K}$$

$$W_B = 1.25 \text{ g}, K_b = 2.53 \text{ °C kg mol}^{-1}$$

$$M_B = ?, W_A = 99 \text{ g}$$

$$M_B \frac{K_b W_B}{T_b W_A} 1000$$

$$= \frac{2.53 + 25 + 1000}{0.21 + 99} \frac{3162.5}{20.79}$$

$$M_B = 152 \text{ g/mol}$$

29. (a) (i)
$$3HgCl_2 + 2PH_3$$
 $Hg_3P_2 + 6HCl_3$

(ii)
$$SiO_2 + 6HF$$
 $H_2SiF_6 + 2H_2O$

- (b) (i) Sulphur in vapour state exhibits paramagnetic behaviour because it forms S_2 molecules like O_2 which contains two unpaired electrons.
 - (ii) The stability of +3 state increases down the group in group 15 due to inert pair effect.
 - (iii) This is due to presence of two bond pairs and three lone pairs of electrons in XeF₂.

OR

(a) (i)
$$AgCl + 2NH_3$$
 $[Ag (NH_3)_2]^+ Cl^-$

(ii)
$$P_4 + 3NaOH + 3H_2O$$
 $PH_3 + 3NaH_2PO_2$

- (b) (i) This is because bond dissociation enthalpy of H-Te bond is less than H-S as the size of Te is larger than S.
 - (ii) Fluorine is a stronger oxidising agent than chlorine due to low dissociation enthalpy of F-F bond and high hydration enthalpy of F-ions.
 - (iii) Noble gases are the least reactive elements due to fully filled outermost shells and zero electron gain enthalpy.

30. (a) (i)
$$CH_3$$
 CH_3 $COCH_3$ CH_3 $COCH_3$ $COCH_3$ CH_3 $COCH_3$ CH_3 $COCH_3$ $COCH_3$

(b) (i) Ethanol and propanal

Benzene

Iodoform test: Ethanol will give iodoform test positive while propanal will not.

Acetophenone

(ii) Benzaldehyde and Acetophenone

Iodoform test: Treat both the compounds with I_2 and NaOH. A yellow ppt. of iodoform will be obtained in acetophenone.

$$COCH_3$$
 $I_{2, NaOH}$ CHI_3 $+$ $3NaI + $3H_2O$$

(iii) Phenol and Benzoic acid

Neutral FeCl₃ test: Add few drops of FeCl₃ solution to both the samples of the compounds. A violet colouration will be obtained in case of phenol.

(a) Mechanism of nucleophilic addition reactions:

A nucleophile attacks the electrophilic carbon atom of the polar carbonyl group from a direction approximately perpendicular to the plane of sp^2 hybridized orbitals of carbonyl carbon. The hybridization of carbon changes from sp^2 to sp^3 in this process and a tetrahedral intermediate is produced. The intermediate captures a proton from the medium to give the neutral product.

(b) (i)
$$CH_3 - C - CH_3$$
 $CH_4 - CH_3 - CH - CH_3$ CH_2SO_4 $CH_3 - CH_2CH_3$ $CH_3 - CH_4$ $CH_4 - CH_5$ $CH_5 - CH_5$ $CH_5 - CH_5$ $CH_6 - CH_6$ $CH_7 - CH_7$ $CH_8 - CH_8$ $CH_$

(ii)
$$C_2H_5OH$$
 PCC CH_3 CHO dil NaOH CH_3 — CH — CH_2 — CHO Ethanol CH_3 — CH — CH_2 — CHO — CH_3 — CH —

(iii)
$$KMnO_4/H^+$$
 $COCI$ $C=O$ $Renzene$ $Anh. AlCl_3$ $Renzene$ $Anh. AlCl_3$ $Renzene$ $Renz$

CBSE (Foreign) SET-II

- 1. Metallic solids have high melting and boiling points.
- 2. CH₃ CHCICH₂CH₃ because it forms secondary carbocation which is more stable.
- **9.** (a) (i) Ozone molecule:

$$(ii) \text{ Structure of } S_8 \text{ molecule}$$

$$S_8 S_8 S_8$$

14. (*i*) Propan-2-ol can be prepared from propene by hydration as shown below:

$$CH_3$$
 — $CH = CH_2 + conc. H_2SO_4$ CH_3 — CH — CH_3 —

Isopropyl hydrogen sulphate

$$CH_3$$
 — CH — CH_3 H_2O Boiling CH_3 — CH — CH_3 + H_2SO_4 OH OH Propan -2-ol

19.
$$M = \frac{d \quad a^3 \quad N_A}{Z} = \frac{10.5 \text{ g cm}^{-3} \quad (4.077 \quad 10^{-8} \text{ cm})^3 \quad (6.02 \quad 10^{23} \text{ mol}^{-1})}{4}$$

$$= \frac{10.5 \text{ g cm}^{-3} \quad 67.767 \quad 10^{-24} \text{ cm}^3 \quad 6.02 \quad 10^{23} \text{ mol}^{-1}}{4}$$

 $= 107.09 \text{ g mol}^{-1}$

20. Given
$$E^{o} Ag^{+}/Ag = 0.80V$$
, $E^{o}_{Cu^{2}}/Cu} = +0.34V$
 $[Cu^{2+}] = 0.10 M$ $[Ag^{+}] = ?$
 $E^{o}_{cell} = E^{o}_{R} - E^{o}_{L} = 0.80 - 0.34V = 0.46V$

From Nernst Equation:

$$E_{cell} = E_{cell}^{o} \quad \frac{0.0591}{2} \log \frac{[Cu^{2}]}{[Ag]^{2}}$$

$$0.422 = 0.46 \quad 0 \quad [0.10]$$

$$\log \frac{0.10}{[Ag]^{2}} \quad 1.2881$$

$$[Ag^{+}]^{2} = 0.0051$$

- $[Ag^+] = 7.1 \times 10^{-2} M$
- (i) The positively charged colloidal particles of Fe(OH)3 get coagulated by the oppositely 21. charged Cl⁻ ions provided by NaCl.
 - (ii) On passing direct current, colloidal particles move towards the oppositely charged electrode where they lose their charge and get coagulated.
 - (iii) Scattering of light by the colloidal particles takes place and the path of light becomes visible (Tyndall effect).
- 26. Biodegradable detergents: Detergents having straight hydrocarbon chains are easily degraded by micro-organism and hence called biodegradable detergents, e.g., sodium-4-(1-dodecyl) benzene sulphonate.

$$CH_3$$
 $(CH_2)_{11}$ SO_3 Na

Non-biodegradable detergents: Detergents having branched hydrocarbon chains are not easily degraded by micro-organisms and hence are called non-biodegradable detergents, e.g., sodium-4-(1, 3, 5, 7-**tetramethyl octyl**) benzenesulphonate.


$${
m CH_3}$$
 ${
m CH_3}$ ${
m CH_3}$ ${
m CH_3}$ ${
m CH}$ ${
m SO_3}$ ${
m Na}$

Non-biodegradable detergents accumulate in rivers and waterways thereby causing water pollution.

- 27. (i) $CH_2 = CHC1$ Vinyl chloride
 - (ii) CF₂ = CF₂ Tetrafluoroethene
 - (iii) Phenol and formaldehyde

CBSE (Foreign) SET-III

- Second order. 1.
- Those colloids which can be separated back into dispersed phase and dispersion medium. 3.
- 4. [Co(NH₂)₆] [Cr(CN)₆] and [Cr(NH₂)₆] [Co(CN)₆]
- Glucose gets oxidised to succinic acid 8.



XeF₂ (Linear shape)

16. (i) Kolbe's reaction

(ii) Friedel-Crafts' acetylation of anisole

19. Given:
$$d = 3.18 \text{ g cm}^{-3}$$
, $Z = 4$, $a = 5.46 \times 10^{-8} \text{ cm}$

$$M = 78.08 \text{ g mol}^{-1} N_A = ?$$

$$d \frac{Z M}{a^3 N_A}$$

$$N_A \frac{Z M}{a^3 P}$$

Substituting the given values, we get

$$N_A = \frac{4 \cdot 78.08}{(5.46 \cdot 10^{-8})^{-3} \cdot 3.18}$$

$$N_A = \frac{313 \cdot 32 \cdot 10^{24}}{517 \cdot 61}$$

$$N_A = 6.035 \times 10^{23} \text{ mol}^{-1}$$

Here,
$$n = 6$$
, $[Al^{3+}] = 0.001M = 1 \times 10^{-3}M$, $[Ni^{2+}] = 0.5M$
 $E_{cell}^{0} = E_{Ni^{2}/Ni}^{0}$ $E_{Al^{3}/Al}^{0}$ 0.25 V 1.66 V
 $E_{cell}^{0} = 1.41 \text{ V}$
 $E_{cell} = 1.41 \frac{0.0591}{6} \log \frac{(10^{-3})^{2}}{(0.5)^{3}}$ $1.41 \frac{0.0591}{6} \log \frac{10^{-6}}{0.125}$
 $= 1.41 \frac{0.0591}{6} \log (10^{-6} 8)$ $1.41 - \frac{0.0591}{6} (\log 10^{-6} \log 2^{3})$
 $= 1.41 \frac{0.0591}{6} (-6 \log 10 - 3 \log 2)$ $1.41 - \frac{0.0591}{6} (-6 - 3 - 0.3010)$
 $= 1.41 \frac{0.0591}{6} (5.097) = 1.41 \frac{0.3012}{6}$
 $= 1.41 + 0.0502 = 1.4602 \text{ V}$
 $E_{cell} = 1.46 \text{ V}$

21. Three features of chemisorption

- (i) It is caused by chemical bond formation and enthalpy of adsorption is high (80-240 kJ mol⁻¹)
- (ii) It forms unimolecular layer.
- (iii) It is highly specific in nature.
- **22.** (*i*) The transition elements have great tendency for complex formation due to presence of vacant d-orbitals of suitable energy, small size of cations and higher nuclear charge.
 - (ii) There is a gradual decrease in the atomic sizes of transition elements in a series with increasing atomic numbers due to poor shielding effect of d-electrons, the net electrostatic attraction between the nucleus and the outermost electrons increase.
 - (iii) Lanthanum and Lutetium do not show colouration in solutions because both the element exhibit +3 oxidation state in their compound thus their cations do not possess any unpaired electrons in them.

CBSE EXAMINATION PAPERS

DELHI-2011

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question numbers 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question numbers 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question numbers 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question numbers 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables, if necessary. Use of calculators is not allowed.

CBSE (Delhi) SET-I

- 1. 'Crystalline solids are anisotropic in nature.' What does this statement mean?
- 2. Express the relation between conductivity and molar conductivity of a solution held in a cell?
- 3. Define 'electrophoresis'.
- **4.** Draw the structure of XeF_2 molecule.
- **5.** Write the IUPAC name of the following compound:

- **6.** Draw the structure of 3-methylbutanal.
- 7. Arrange the following compounds in an increasing order of their solubility in water:

$$C_6H_5NH_2$$
, $(C_2H_5)_2NH$, $C_2H_5NH_2$

- **8.** What are biodegradable polymers?
- **9.** The chemistry of corrosion of iron is essentially an electrochemical phenomenon. Explain the reactions occurring during the corrosion of iron in the atmosphere.
- **10.** Determine the values of equilibrium constant (K_C) and G° for the following reaction:

$$Ni(s) + 2Ag^{+}(aq)$$
 $Ni^{2+}(aq) + 2Ag(s)$, $E^{\circ} = 1.05 \text{ V}$
(1 F = 96500 C mol⁻¹)

- 11. Distinguish between 'rate expression' and 'rate constant' of a reaction.
- **12.** State reasons for each of the following:
 - (i) The N-O bond in NO_2^- is shorter than the N-O bond in NO_3^- .
 - (ii) SF₆ is kinetically an inert substance.

OR

State reasons for each of the following:

- (i) All the P—Cl bonds in PCl₅ molecule are not equivalent.
- (ii) Sulphur has greater tendency for catenation than oxygen.

172 Xam idea Chemistry—XII

- **13.** Assign reasons for the following:
 - (i) Copper (I) ion is not known in aqueous solution.
 - (ii) Actinoids exhibit greater range of oxidation states than lanthanoids.
- **14.** Explain the following giving one example for each:
 - (i) Reimer-Tiemann reaction.
 - (ii) Friedel-Crafts acetylation of anisole.
- 15. How would you obtain
 - (i) Picric acid (2, 4, 6-trinitrophenol) from phenol,
 - (ii) 2-Methylpropene from 2-methylpropanol?
- **16.** What is essentially the difference between -form of glucose and -form of glucose? Explain.
- 17. Describe what you understand by primary structure and secondary structure of proteins.
- **18.** Mention two important uses of each of the following:
 - (i) Bakelite
 - (ii) Nylon-6
- 19. Silver crystallizes in face-centered cubic unit cell. Each side of this unit cell has a length of 400 pm. Calculate the radius of the silver atom. (Assume the atoms just touch each other on the diagonal across the face of the unit cell. That is, each face atom is touching the four corner atoms.)
- **20.** Nitrogen pentoxide decomposes according to equation:

$$2N_2O_5(g)$$
 $4NO_2(g) + O_2(g)$.

This first order reaction was allowed to proceed at 40°C and the data below were collected:

[N ₂ O ₅] (M)	Time (min)
0.400	0.00
0.289	20.0
0.209	40.0
0.151	60.0
0.109	80.0

- (a) Calculate the rate constant. Include units with your answer.
- (b) What will be the concentration of N_2O_5 after 100 minutes?
- (c) Calculate the initial rate of reaction.
- 21. Explain how the phenomenon of adsorption finds application in each of the following processes:
 - (i) Production of vacuum
 - (ii) Heterogeneous catalysis
 - (iii) Froth floatation process

OR

Define each of the following terms:

- (i) Micelles
- (ii) Peptization
- (iii) Desorption

- 22. Describe the principle behind each of the following processes:
 - (i) Vapour phase refining of a metal.
 - (ii) Electrolytic refining of a metal.
 - (iii) Recovery of silver after silver ore was leached with NaCN.
- **23.** Complete the following chemical equations:
 - (i) $MnO_4^ C_2O_4^2$ H
 - (ii) KMnO₄ heated
 - (iii) $Cr_2O_7^2$ H_2S H
- 24. Write the name, stereochemistry and magnetic behaviour of the following:
 - (At. nos. Mn = 25, Co = 27, Ni = 28)
 - (i) $K_4[Mn(CN)_6]$
 - (ii) [Co(NH₃)₅Cl]Cl₂
 - (iii) K₂[Ni(CN)₄]
- **25.** Answer the following:
 - (i) Haloalkanes easily dissolve in organic solvents, why?
 - (ii) What is known as a racemic mixture? Give an example.
 - (iii) Of the two bromoderivatives, C₆H₅CH(CH₃)Br and C₆H₅CH(C₆H₅)Br, which one is more reactive in S_N1 substitution reaction and why?
- **26.** (a) Explain why an alkylamine is more basic than ammonia.
 - (b) How would you convert
 - (i) Aniline to nitrobenzene
 - (ii) Aniline to iodobenzene?
- **27.** Describe the following giving one example for each:
 - (i) Detergents
 - (ii) Food preservatives
 - (iii) Antacids
- **28.** (a) Differentiate between molarity and molality for a solution. How does a change in temperature influence their values?
 - (b) Calculate the freezing point of an aqueous solution containing 10.50 g of MgBr₂ in 200 g of water. (Molar mass of MgBr₂= 184 g)

$$(K_f \text{ for water} = 1.86 \text{ K kg mol}^{-1})$$

- (a) Define the terms osmosis and osmotic pressure. Is the osmotic pressure of a solution a colligative property? Explain.
- (b) Calculate the boiling point of a solution prepared by adding 15.00 g of NaCl to 250.00 g of water. (K_b for water = 0.512 kg mol⁻¹),

(Molar mass of NaCl =
$$58.44$$
 g)

174 Xam idea Chemistry—XII

- **29.** (a) Give chemical tests to distinguish between
 - (i) Propanal and propanone,
 - (ii) Benzaldehyde and acetophenone.
 - (b) How would you obtain
 - (i) But-2-enal from ethanal
 - (ii) Butanoic acid from butanol,
 - (iii) Benzoic acid from ethylbenzene?

OR

- (a) Describe the following giving linked chemical equations:
 - (i) Cannizzaro reaction
 - (ii) Decarboxylation
- (b) Complete the following chemical equations:

$$(i) \begin{tabular}{c} $\operatorname{CH}_2\operatorname{CH}_3$ \\ & \operatorname{KMnO}_4$ \\ & \operatorname{KOH, heat} \end{tabular}$$

(iii)
$$C_6H_5CONH_2$$
 H_3O heat

- **30.** (a) Explain the following:
 - (i) NF₃ is an exothermic compound whereas NCl₃ is not.
 - (ii) F₂ is most reactive of all the four common halogens.
 - (b) Complete the following chemical equations:

(i) C
$$H_2SO_4$$
(conc.)

$$(ii)$$
 P₄ + NaOH + H₂O

- (a) Account for the following:
 - (i) The acidic strength decreases in the order $HCl > H_2S > PH_3$.
 - (ii) Tendency to form pentahalides decreases down the group in group 15 of the periodic table.
- (b) Complete the following chemical equations:

(i)
$$P_4 SO_2Cl_2$$

CBSE (Delhi) SET - II

Questions Uncommon to Set-I

- 1. Which stoichiometric defect in crystals increases the density of a solid?
- 3. What is meant by 'shape-selective catalysis' of reactions?
- **4.** Draw the structure of XeF₄ molecule.
- **9.** Explain what is meant by (i) a peptide linkage, (ii) a glycosidic linkage.
- 10. Name the bases present in RNA. Which one of these in not present in DNA?
- **22.** Explain the role of each of the following in the extraction of metals from their ores:
 - (i) CO in the extraction of nickel.
 - (ii) Zinc in the extraction of silver.
 - (iii) Silica in the extraction of copper.
- **24.** For the complex $[Fe(en)_2 Cl_2] Cl$, identify the following:
 - (i) Oxidation number of iron.
 - (ii) Hybrid orbitals and shape of the complex.
 - (iii) Magnetic behaviour of the complex.
 - (iv) Number of its geometrical isomers.
 - (v) Whether there may be optical isomer also.
 - (vi) Name of the complex.
- **27.** Explain the following terms with one suitable example for each:
 - (i) A sweetening agent for diabetic patients.
 - (ii) Enzymes
 - (iii) Analgesics
- **28.** (a) State the following:
 - (i) Henry's law about partial pressure of a gas in a mixture.
 - (ii) Raoult's law in its general form in reference to solutions.
 - (b) A solution prepared by dissolving 8.95 mg of a gene fragment in 35.0 mL of water has an osmotic pressure of 0.335 torr at 25°C. Assuming the gene fragment is a non-electrolyte, determine its molar mass.

- (a) Differentiate between molarity and molality in a solution. What is the effect of temperature change on molarity and molality in a solution?
- (b) What would be the molar mass of a compound if 6.21g of it dissolved in 24.0 g of chloroform form a solution that has a boiling point of 68.04°C. The boiling point of pure chloroform is 61.7°C and the boiling point elevation constant, K_b for chloroform is 3.63°C/m.

CBSE (Delhi) SET-III

Questions Uncommon to Set-I and II

- 2. Express the relation among the conductivity of solution in the cell, the cell constant and the resistance of solution in the cell.
- **4.** Draw the structure of BrF₃ molecule.
- **8.** In nylon-6, 6 what does the designation '6, 6' mean?
- **9.** What type of a battery is lead storage battery? Write the anode and the cathode reactions and the overall reactions occurring in a lead storage battery.
- **10.** Two half-reactions of an electrochemical cell are given below:

$$MnO_4^-(aq)$$
 8H (aq) 5e⁻ Mn^2 (aq) 4H₂O_(l), E^o 1.51 V
Sn² (aq) Sn^4 (aq) 2e⁻, E^o 0.15 V

Construct the redox equation from the standard potential of the cell and predict if the reaction is reactant favoured or product favoured.

- 13. Assign reasons for each of the following:
 - (i) Transition metals generally form coloured compounds.
 - (ii) Manganese exhibits the highest oxidation state of +7 among the 3d series of transition elements.
- **18.** Name the sub-groups into which polymers are classified on the basis of magnitude of intermolecular forces.
- **19.** The density of lead is 11.35 g cm⁻³ and the metal crystallizes with fcc unit cell. Estimate the radius of lead atom. (At. mass of lead = 207 g mol⁻¹ and $N_A = 6.02 \times 10^{23}$ mol⁻¹)
- **26.** Complete the following chemical equations :

(i)
$$CH_3CH_2Cl$$
 NaCN (A) reduction Ni/ H_2 (B)

$$\begin{array}{ccc} (ii) & \mathrm{C_6H_5N_2Cl} + \mathrm{H_3PO_2} + \mathrm{H_2O} \\ & & \mathrm{O} \\ & || \\ (iii) & \mathrm{R-C-NH_2} & \mathrm{LiAlH_4} \\ & & \mathrm{H_2O} \end{array}$$

- **27.** Answer the following questions :
 - (i) Why do soaps not work in hard water?
 - (ii) What are the main constituents of dettol?
 - (iii) How do antiseptics differ from disinfectants?




CBSE (Delhi) SET - I


- 1. Crystalline solids are anisotropic in nature means some of their physical properties like electrical conductivity, refractive index, etc., are different in different directions.
- $\frac{1000}{C}$, where is the conductivity and C is the molar concentration.
- 3. The movement of colloidal particles under an applied electric potential is called electrophoresis.
- 4. Total no. of electron pairs in the valence shell of central Xe atom = $\frac{8}{2}$ = 5

No. of bond pairs = 2

No. of lone pairs = 3

Therefore, the shape of molecule would be linear.





6.
$$CH_3 O | CH_3 - CH - CH_2 - C - H$$

7. Increasing order of solubility of amines in water

$$C_6H_5NH_2 < (C_2H_5)_2NH < C_2H_5NH_2$$

- 8. Polymers which get disintegrated by themselves in biological systems during a certain period of time by enzymatic hydrolysis or to some extent by oxidation are called biodegradable polymers. An example of biodegradable polymer is PHBV, i.e., poly-hydroxy butyrate-co-hydroxyvalerate.
- 9. According to electrochemical theory of rusting the impure iron surface behaves like small electrochemical cell in the presence of water containing dissolved oxygen or carbon dioxide. In this cell pure iron acts as anode and impure iron surface acts as cathode. Moisture having dissolved CO₂ or O_2 acts as an electrolyte. The reactions are given below.

At anode:

Fe Fe²
$$2e^{-}$$
; $E^{o}_{Fe^{2}/Fe}$ 0.44 V

At cathode: 2H
$$\frac{1}{2}$$
O₂ 2e⁻ H₂O; $E_{\text{H}^+/\text{O}_2/\text{H}_2\text{O}}$ 1.23V

Overall reaction: Fe + 2H $\frac{1}{2}$ O₂ Fe² H₂O; E_{cell} 1.67V

Fe²
$$H_2O$$
; E_{cell} 1.67

The Fe²⁺ ions are further oxidised by atmospheric oxygen to Fe³⁺ ions, which comes out in the form of hydrated ferric oxide (rust).

$$2Fe^2 \quad \frac{1}{2}O_2 \quad 2H_2O \qquad Fe_2O_3 \quad 4H$$

$$Fe_2O_3 + xH_2O$$
 $Fe_2O_3 xH_2O$ (Rust)

10. Ni(s) + 2Ag⁺(aq) Ni²⁺ (aq) + 2Ag(s);
$$E^{\circ} = 1.05 \text{ V}$$

Here, $n = 2$

$$\log K_c = \frac{n}{0.059} \quad E_{\text{cell}}^{\circ}$$

$$\log K_c = \frac{2}{0.059} \quad 1.05 = 39.5932$$

$$K_c = \text{antilog } 39.5932 = 3.919 \times 10^{39}$$

$$K_c = 3.92 \times 10^{39}$$

$$G^{\circ} = -nFE_{\text{cell}}^{\circ}$$

$$G^{\circ} = -2 \times 96500 \times 1.05 = -202650 \text{ J}$$

$$G^{\circ} = -202.65 \text{ kJ}$$

11. Rate expression is an experimentally determined expression which relates the rate of reaction with the concentration of the reactants whereas rate constant is the rate of reaction when concentration of each reactant in the reaction is unity.

Consider a general reaction

$$aA + bB$$
 $cC + dD$

The rate expression for this reaction is

Rate =
$$k [A]^m [B]^n$$

Where the proportionality constant *k* is called rate constant.

- 12. (i) This is because the N—O bond in NO₂ is an average of a single bond and a double bond whereas the N—O bond in NO₃ is an average of two single bonds and a double bond.
 - (ii) In SF₆, S atom is sterically protected by six F atoms and does not allow water molecules to attack the S atom. Further, F does not have d-orbitals to accept the electrons denoted by H₂O molecules. Due to these reasons, SF₆ is kinetically an inert substance.

OR

- (i) In PCl₅ the two axial bonds are longer than the three equatorial bonds. This is due to the fact that the axial bond pairs suffer more repulsion as compared to equatorial bond pairs.
- (ii) The property of catenation depends upon bond strength of the element. As S—S bond is much stronger (213 kJ mol⁻¹) than O—O bond (138 kJ mol⁻¹), sulphur has greater tendency for catenation than oxygen.
- **13.** (i) In aqueous solution Cu⁺ undergoes disproportionation to form a more stable Cu²⁺ ion.

$$2 \text{ Cu}^+ \text{ (aq)}$$
 $2 \text{ Cu}^{2+} \text{ (aq)} + \text{ Cu(s)}$

The higher stability of Cu^{2+} in aqueous solution may be attributed to its greater negative $_{hyd}H^{\circ}$ than that of Cu^{+} . It compensates the second ionization enthalpy of Cu involved in the formation of Cu^{2+} ions.

- (ii) Actinoids exhibit greater range of oxidation states than lanthanoids. This is because there is less energy difference between 5f and 6d orbitals belonging to actinoids than the energy difference between 4f and 5d orbitals in case of lanthanoids.
- **14.** (*i*) **Reimer–Tiemann Reaction:** Treatment of phenol with chloroform in the presence of aqueous alkali at 340 K followed by hydrolysis of resulting product gives salicylaldehyde as a major product.

(ii) Friedel-Crafts acylation of anisole: Anisole on treatment with acylchloride in the presence of anhydrous AlCl₃ undergo electrophilic substitution in the ring at ortho and para positions. For example,

OH OH OH

15. (i)
$$H_2 SO_4 (conc.)$$
 $SO_3 H$ $HNO_3 (conc.)$ $O_2 N$ NO_2 $O_3 H$ $O_4 MO_2$ $O_3 H$ $O_4 MO_2$ $O_5 MO_2$ $O_6 MO_2$ $O_7 MO_2$ $O_8 MO_2$ O_8

In -D Glucose, the —OH group at C_1 is toward right whereas in -glucose, the -OH group at C_1 is towards left. Such a pair of stereo-isomers which differ in the configuration only around C_1 are called anomers.

17. Primary Structure: The specific sequence in which the various -amino acids present in a protein are linked to one another is called its primary structure. Any change in the primary structure creates a different protein.

Secondary Structure: The conformation which the polypeptide chain assumes as a result of hydrogen bonding is known as secondary structure. The two types of secondary structures are -helix and -pleated sheet structure.

In -helix structure, the polypeptide chain forms all the possible hydrogen bonds by twisting into a right handed screw (helix) with the —NH groups of each amino acid residue hydrogen bonded to the C=Ogroups of an adjacent turn of the helix. In -pleated structure, all peptide chains are stretched out nearly to maximum extension and then laid side by side which are held together by hydrogen bonds.

- **18.** (*i*) Bakelite is used for making combs, electrical switches, handles of various utensils and phonograph records.
 - (ii) Nylon-6 is used for making tyre cords, ropes and fabrics.
- **19.** a = 400 pm

For fcc,
$$r = \frac{a}{2\sqrt{2}}$$

 $r = \frac{400}{2\sqrt{2}} = \frac{400}{2\sqrt{2}} = \frac{\sqrt{2}}{\sqrt{2}} = \frac{400\sqrt{2}}{4} = 100\sqrt{2}$
 $r = 100 \times 1.414 = 141.4 \text{ pm}$

20. (a) When t = 20 min, [A] = 0.289 mol L⁻¹

Also,
$$[A]_0 = 0.400 \text{ mol } L^{-1}$$

For a first order reaction

$$k = \frac{2.303}{t} \log \frac{[A]}{[A]}$$

$$k = \frac{2.303}{20} \log \frac{0.400}{0.289}$$

$$k = \frac{2.303}{20} \log \frac{4.00}{2.89}$$

$$k = \frac{2.303}{20} [\log 4.00 - \log 2.89]$$

$$k = \frac{2.303}{20} [0.6021 - 0.4609]$$

$$k = \frac{2.303}{20} 0.1412$$

$$k = 2.303 \times 0.00706 = 0.016259 \text{ min}^{-1}$$

$$k = 1.6259 \times 10^{-2} \text{ min}^{-1}$$

(b)
$$t = \frac{2303}{k} \log \frac{[A]_0}{[A]}$$

Here,
$$[A]_0 = 0.400 \text{ mol}^{-1}$$
, $t = 100 \text{ min}$, $k = 1.626 \times 10^{-2} \text{ min}^{-1}$
$$100 = \frac{2.303}{1.626 \cdot 10^{-2}} \log \frac{0.400}{[A]}$$

$$\frac{100 \quad 1.626 \quad 10^{-2}}{2.303} = \log \frac{0.4}{[A]}$$

$$0.7060 = \log \frac{0.4}{[A]}$$

Antilog
$$(0.7060) = \frac{0.4}{[A]}$$

$$5.082 = \frac{0.4}{[A]}$$

$$[A] = \frac{0.4}{5.082} = 0.0787 \text{ M}$$

(c) Initial rate, i.e., rate of reaction when t = 0

When,
$$t = 0.00 \text{ min}$$
, [A] = 0.400 mol L⁻¹
Also, $k = 1.626 \times 10^{-2} \text{ min}^{-1}$

Initial rate =
$$k$$
 [A] = $1.626 \times 10^{-2} \text{ min}^{-1} \times 0.400 \text{ mol L}^{-1}$
= $6.504 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$.

- **21.** (i) **Production of Vacuum:** Adsorption can be successfully applied to create conditions of high vacuum. For this a bulb of charcoal cooled in liquid air, is connected to vessel which has already been exhausted as far as possible by vacuum pump. The remaining traces of air inspite of low pressure are adsorbed by the charcoal almost completely.
 - (ii) Heterogeneous Catalysis: There are many gaseous reactions of industrial importance involving solid catalyst. Manufacture of ammonia using iron as a catalyst, manufacture of H_2SO_4 by contact process using V_2O_5 catalyst and use of finely divided nickel in the hydrogenation of vegetable oils are the excellent examples. The gaseous reactants are adsorbed on the surface of the solid catalyst. As a result, the concentration of the reactants increases on the surface of the catalyst and hence the rate of reaction increases.
 - (iii) Froth Floatation Process: In froth floatation process the powdered ore is mixed with water. It is then mixed with pine oil (a frother). The oil particles are adsorbed on the surface of ore particles. Now, a stream of air is blown through the mixture from below when froth is formed at the water surface. The ore particles stick to the bubbles of the air rises to surface along with the foam while the gangue particles which are wetted by water settle at the bottom. The foam is separated out and is collected and in the course, the ore particles also settle down.

OR

(i) Micelles: There are some substances which at low concentration behave as normal strong electrolytes but at higher concentration exhibit colloidal behaviour due to formation of aggregated particles. The aggregated particles thus formed are called micelles. The formation

- of micelles takes place only above a particular temperature called **Kraft temperature** and above a particular concentration called **critical micelle concentration** (CMC). Surface active agents such as soap and synthetic detergents belong to this class.
- (ii) **Peptization:** The process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of a small amount of suitable electrolyte is called peptization. During peptization, the precipitate absorbs one of the ions of the electrolyte on its surface. This causes development of positive or negative charge on precipitates, which ultimately break up into particles of colloidal dimension.
- (iii) Desorption: The process of removing an adsorbed substance from a surface on which it is adsorbed is called desorption.
- **22.** (i) Vapour phase refining of a metal: In this method, the metal is converted into its volatile compound and collected elsewhere. It is then decomposed to give pure metal. For example, refining of nickel by Mond process.

(ii) Electrolytic refining of a metal: In this method, the impure metal is made to act as anode. A strip of same metal in pure form is used as cathode. They are put in a suitable electrolytic bath containing soluble salt of same metal. When electric current is passed the metal from the anode goes into solution as ions due to oxidation while pure metal gets deposited at the cathode due to reduction of metal ions. The voltage applied for electrolysis is such that impurities of more electropositive metals remains in the solution as ions while impurities of the less electropositive metals settle down under the anode as anode mud.

At anode:
$$M$$
 M^n $n \bar{e}$ At cathode: M^n $n \bar{e}$ M

(iii) Recovery of silver after silver ore was leached with NaCN: During leaching Ag is oxidised to Ag⁺ which then combines with CN⁻ ions to form soluble complex, [Ag(CN)₂]⁻. Silver is then recovered from this complex by displacement method using more electropositive zinc metal.

$$2[Ag(CN)_{2}]^{-}(aq) + Zn(s) 2Ag(s) + [Zn(CN)_{4}]^{2-}(aq)$$
23. (i) $MnO_{4}^{-} + 8H^{+} + 5\bar{e}$ $Mn^{2+} + 4H_{2}O] \times 2$

$$C_{2}O_{4}^{2-} 2CO_{2} + 2\bar{e}] \times 5$$

$$2MnO_{4}^{-} + 5C_{2}O_{4}^{2-} + 16H^{+} 2Mn^{2+} + 10CO_{2} + 8H_{2}O$$

(ii)
$$2KMnO_4$$
 heat $K_2MnO_4 + MnO_2 + O_2$
(iii) $Cr_2O_7^{2-} + 14H^+ + 6\bar{e}$ $2Cr^{3+} + 7H_2O$
 H_2S $2H^+ + S + 2e] \times 3$
 $Cr_2O_7^{2-} + 3H_2S + 8H^+$ $2Cr^{3+} + 3S + 7H_2O$

Iodo benzene

Name of the Complex	Hybridisation of metal ion involved	Geometry of complex	Magnetic behaviour
Potassium hexacyano manganate (II)	d^2sp^3	Octahedral	Paramagnetic
Pentaammine Chloridocobalt (III) Chloride	d^2sp^3	Octahedral	Diamagnetic
Potassium tetra cyanonikelate (II)	dsp^2	Square planar	Diamagnetic

- **25.** (*i*) Haloalkanes dissolve in organic solvents because the new intermolecular attractions between haloalkanes and organic solvent molecules have much the same strength as ones being broken in the separate haloalkanes and solvent molecules.
 - (ii) An equimolar mixture of a pair of enantiomers is called racemic mixture. For example, butan-2-ol. A racemic mixture is optically inactive due to external compensation.
 - (iii) Of the two bromo derivatives, $C_6H_5CH(CH_3)Br$ and $C_6H_5CH(C_6H_5)Br$, the intermediate obtained from C_6H_5CH (C_6H_5) Br is more stable than obtained from $C_6H_5CH(CH_3)$ Br because it is stabilised by two phenyl groups due to resonance. Therefore, C_6H_5CH (C_6H_5)Br is more reactive than $C_6H_5(CH_3)$ Br.

26.

$$(a) \ \mathbf{R} - \mathbf{N}; \quad + \quad \mathbf{H}^{+} \quad \longrightarrow \quad \mathbf{R} - \mathbf{N}^{+} - \mathbf{H}$$

Aniline

Due to electron releasing nature, the alkyl group (R) pushes electrons towards nitrogen in alkyl amine and thus makes the unshared electron pair more available for sharing with the proton of the acid. Therefore alkyl amine are more basic than ammonia.

$$(b) \quad (i) \qquad \begin{array}{c} NH_2 \\ NaNO_2/HCl \\ \hline \\ Nitro benzene \\ \hline \\ Nit$$

27. (*i*) **Detergents:** Detergents are cleansing agents which have all the properties of soaps, but actually do not contain any soap *e.g.*, sodium dodecylbenzene sulphonate. These can be used both in soft and hard water as they give foam even in hard water. Detergents are mainly classified into three categories:

- (i) Anionic detergents
- (ii) Cationic detergents
- (iii) Non-ionic detergents
- (ii) Food Preservatives: These are the chemical substances which are added to the food materials to prevent their spoilage due to microbial growth and to retain their nutritive value for long time. Sodium benzoate, sodium metabisulphite are some common preservatives.

Preservatives prevent rancidity and kill or inhibit the growth of microorganism.

- (iii) Antacids: These are the chemical substances which neutralize the excess acid and raise the pH to an appropriate level in the stomach. Sodium hydrogen carbonate or a mixture of aluminium and magnesium hydroxide are some common antacids.
- **28.** (a) Molarity is the number of moles of solute dissolved in one litre of solution whereas molality is the number of moles of solute per kilogram of the solvent. Molarity decreases with increase in temperature as volume increases with increase in temperature. Molality is independent of temperature because mass does not depend on temperature.

(b)
$$T_f = \frac{i \quad K_f \quad W_B \quad 1000}{M_B \quad W_A}$$

 $i = 3, K_f = 1.86 \text{ K kg mol}^{-1}, W_B = 10.5 \text{ g}$
 $M_B = 184 \text{ g mol}^{-1}, W_A = 200 \text{ g}$

Substituting these values in above equation, we get

$$T_f = \frac{3 - 1.86 \,\mathrm{K \ kg \ mol}^{-1} - 10.5 \,\mathrm{g}^{-1000 \,\mathrm{g \ kg}^{-1}}}{184 \,\mathrm{g \ mol}^{-1} - 200 \,\mathrm{g}}$$

$$T_f = 1.59 \text{ K}$$

Therefore, freezing of aqueous solution

$$= 273.15 \text{ K} - 1.59 \text{ K}$$

 $= 271.56 \text{ K}$

OR

(a) **Osmosis:** The spontaneous movement of the solvent molecules either from the pure solvent to the solution or from a less concentrated solution to a more concentrated solution through a semi-permeable membrane.

Osmotic Pressure: The minimum excess pressure that has to be applied on the solution to prevent the passage of solvent molecules into it through semipermeable membrane is called osmotic pressure.

Osmotic pressure is a colligative property as it depends on the number of moles of solute particles and not on their identity.

(b)
$$T_{b} = \frac{i \quad K_{b} \quad W_{B} \quad 1000}{M_{B} \quad W_{A}}$$

$$i = 2, K_{b} = 0.512 \text{ K kg mol}^{-1}, W_{B} = 15g$$

$$M_{B} = 58.44 \text{ g mol}^{-1}, W_{A} = 250 \text{ g}$$
Substitution the selection was a service.

Substituting these values in the above equation, We get

$$T_b = \frac{2 - 0.512 \text{ K kg mol}^{-1} - 15 \text{ g}^{-1000 \text{ g Kg}^{-1}}}{58.44 \text{ g mol}^{-1} - 250 \text{ g}} = 1.05 \text{ K}$$

Therefore, boiling point of aqueous solution

29. (a) (i) Propanal and propanone

Tollen's reagent test: Propanal being an aldehyde reduces Tollen's reagent to silver mirror but propanone being a ketone does not.

$$CH_{3}CH_{2}CHO + 2[Ag(NH_{3})_{2}]^{+} + 3\overline{O}H \qquad CH_{3}CH_{2} \quad CO\overline{O} \quad 2Ag \quad 4NH_{3} \quad 2H_{2}O$$
 Propanal
$$CH_{3}COCH_{3} \quad \frac{Tollen's}{reagent} \quad No \quad silver \quad mirror$$

(ii) Benzaldehyde and acetophenone

Iodoform test: Acetophenone being a methyl ketone on treatment with I₂/NaOH undergoes Iodoform reaction to give yellow ppt. of iodoform but benzaldehyde does not.

$$\begin{array}{ccccc} C_6H_5COCH_3 & 3NaOI & C_6H_5COONa & CHI_3 & 2NaOH\\ & & & & & & & & \\ Acetophenone & & & & & & \\ OH & & & & & & \\ & & & & & & \\ OH & & & & & \\ \end{array}$$

(b) (i) CH₃CHO
$$\stackrel{\text{dil. NaOH}}{\longleftrightarrow}$$
 CH₃CH—CH₂CHO $\stackrel{\Delta}{\longrightarrow}$ CH₃CH—CHCHO But-2-enal

 $CrO_3 - H_2SO_4$

(ii)
$$CH_3CH_2CH_2CH_2OH$$
 $CrO_3 - H_2SO_4$ $CH_3CH_2CH_2COOH$ Butanoic acid

$$CH_2CH_3$$
 $COOK$

Ethyl benzene

$$COOK$$

(i) Cannizzaro reaction: Aldehydes which do not have an -hydrogen atom undergoes (a) disproportionation reactions on treatment with concentrated alkali to give a mixture of carboxylic acid salt and alcohol.

OR

(ii) Decarboxylation: Carboxylic acids lose carbon dioxide to form hydrocarbons when their sodium salts are heated with sodalime.

- **30.** (a) (i) This is because bond dissociation energy of F₂ is lower than Cl₂ Moreover, fluorine forms stronger bond with nitrogen due to comparable size.
 - (ii) Because of low F—F bond dissociation enthalpy.

(excess)

(b) (i)
$$C + 2H_2SO_4$$
 (conc.) $CO_2 + 2SO_2 + 2H_2O$
(ii) $P_4 + 3NaOH + 3H_2O$ $PH_3 + 3NaH_2PO_2$
Phosphine
(iii) $Cl_2 3F_2 573K$ $2ClF_3$

OR

- (a) (i) As bond dissociation enthalpy of H—Cl bond is lower than H—S which is lower than P—H. The acidic strength decreases because of decrease in polarity of E—H bond from H—Cl to P—H which is due to decrease in electonegativity of E.
 - (ii) This is due to inert pair effect. The stability of +5 oxidation state decreases down the group in group 15.

(b) (i)
$$P_4 + 10SO_2Cl_2$$
 $4PCl_5 + 10SO_2$
(ii) $2Xe F_2 + 2H_2O$ $2Xe + 4HF + O_2$
(iii) $2HNO_3$ (conc.) $H_2O + 2NO_2 + (O)] \times 5$
 $I_2 + 5(O)$ I_2O_5
 $I_2O_5 + H_2O$ $2HIO_3$
 $I_2 + 10HNO_3$ (conc.) $2HIO_3 + 10NO_2 + 4H_2O$

CBSE (Delhi) SET - II

- 1. Interstitial defect
- **3.** The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecules is called shape-selective catalysis.


4. Total number of electron pairs in the valence shell of central

Xe atom =
$$\frac{8}{2} = 6$$

No. of bond pairs = 4

No. of lone pairs = 2

Therefore, the shape of molecule would be square planar.



- (i) Peptide linkage: The amide linkage (C NH —) formed between two —amino acid
- molecules with the loss of a water molecule in a polypeptide is called a peptide linkage.

 (ii) Glycosidic linkage: The oxide linkage between two monosaccharides in oligosaccharides or polysaccharides is called glycosidic linkage.
- 10. The bases present in RNA are adenine (A), guanine (G), cytosine (C) and uracil (U). Uracil is not present in DNA.
- **22.** (*i*) **CO in the extraction of nickel:** Impure nickel is heated in a stream of carbon monoxide when volatile nickel tetracarbonyl is formed and the impurities are left behind in the solid state. The vapour of nickel tetra carbonyl is taken to a decomposer chamber maintained at 450–470 K where it decomposes to give pure nickel metal and carbon monoxide.

(ii) Zinc in the extraction of silver: Silver present in the ore is leached with dilute solution of NaCN in the presence of air or oxygen to form a soluble complex.

$$4Ag(s) + 8CN^{-}(aq) + 2H_{2}O(l) + O_{2}(g)$$
 4 [Ag (CN)₂]⁻ + 4 OH⁻(aq)

Silver is then recovered from the complex by displacement method using complex using a more electropositive zinc metal.

$$2[Ag(CN)_2]^-(aq) + Zn(s)$$
 $2Ag(s) + [Zn(CN)_4]^{2-}(aq)$

(iii) Silica in the extraction of copper: During smelting and bessemerisation the impurity ferrous sulphide oxidised to ferrous oxide which is then reacted with silica (flux) to form slag ferrous silicate.

$$\begin{array}{ccc} \text{FeO} & \text{SiO}_2 & \text{FeSiO}_3 \\ & \text{Flux} & \text{Ferrous silicate(slag)} \end{array}$$

24. (*i*) [Fe (en)₂ Cl_2]Cl

$$x + 0 \times 2 + (-1) \times 2 + (-1) \times 1 = 0$$
 $x = 3$

Oxidation number of iron = 3

- (ii) d^2sp^3 hybridization, Octahedral
- (iii) Paramagnetic
- (iv) Two
- (v) Yes, cis-[Fe(en)₂Cl₂]
- (vi) Bis-(ethane-1, 2-diamine)-dichlorido iron (III) chloride
- **27.** (*i*) Artificial sweetening agents are chemical substances which are sweet in taste but do not add calories to our body. For example, saccharin. Its use is of great value to diabetic persons as it is excreted from the body through urine as such.

- (ii) **Enzymes:** Enzymes are catalysts of biological origin which accelerate various cellular reactions without themselves undergoing any apparent change during the course of action. Enzymes are highly specific in their action on substrate. Almost all the enzymes are globular proteins.
- (iii) Analgesics: Analgesics are chemical compounds which are used for relieving pain. Analgesics relieves pain by acting on central nervous system or on peripheral pain mechanism, without significantly affecting consciousness. There are two types of analgesics:

Narcotics — Morphine, cocaine, heroine

Non-narcotics — Aspirin, ibuprofen etc.

28. (a) (i) **Henry's law:** It states that the partial pressure of the gas in vapour phase (p) is proportional to the mole fraction of the gas (x) in the solution and is expressed as:

$$P = K_H x$$

Where K_H is Henry's law constant

(ii) **Raoult's law:** It states that for any solution the partial pressure of each volatile component in the solution is directly proportional to its mole fraction.

$$M_{\rm B} = \frac{W_{\rm B} - R - T}{V}$$

Here, $W_B = 8.95 \text{ mg} = 8.95 \times 10^{-3} \text{g}, R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$ $T = (25 + 273) \text{ K} = 298 \text{ K}, = \frac{0.335}{760} \text{ atm}$ $V = 35 \times 10^{-3} \text{ L}$

Substituting these values in the above equation

We get,
$$M_{\rm B} = \frac{8.95 \cdot 10^{-3} \,\mathrm{g} \cdot 0.0821 \,\mathrm{L} \,\mathrm{atm} \,\mathrm{K}^{-1} \,\mathrm{mol}^{-1}}{\frac{0.335}{760} \,\mathrm{atm} \cdot 35 \cdot 10^{-3} \,\mathrm{L}}$$
 $M_{\rm B} = 14193. \, 29 \,\mathrm{g} \,\mathrm{mol}^{-1}$

OR

(a) Molarity (M): It is defined as number of moles of solute dissolved in one litre of solution.

Molarity
$$\frac{\text{Moles of solute}}{\text{Volume of solution in litre}}$$

Unit of molarity is mol L^{-1} or M (molar). Molarity changes with change in temperature as volume changes with change in temperature.

Molality (*m*): It is defined as the number of moles of the solute dissolved in one kilogram (kg) of the solvent and is expressed as:

Unit of molality is mol kg⁻¹.

Molality is independent of temperature.

(b)
$$M_{B} = \frac{K_{b} \quad W_{B}^{1} \quad 1000}{T_{b} \quad W_{A}}$$
Here,
$$K_{b} = 3.63K \text{ kg mol}^{-1}, \quad W_{B} = 6.21 \text{ g}$$

$$T_b = 68.04$$
°C $- 61.7$ °C $= 6.34$ °C, $W_A = 24.0$ g

Substituting these values in the above equation, we get

$$M_{\rm B} = \frac{3.63 \text{ K kg mol}^{-1} \quad 6.21 \text{ g}^{-1000 \text{ g kg}^{-1}}}{6.34 \text{ K} \quad 24.0 \text{ g}}$$

$$M_{\rm B} = 148.15 \text{ g mol}^{-1}$$

CBSE (Delhi) SET - III


- (conductivity) = $\frac{1}{R \text{ (Resistance)}}$ G (Cell constant)
- 4. Total number of electron pairs in the valence shell of central

Br atom =
$$\frac{7}{2}$$
 = 5

Number of bond pairs = 3

Number of lone pairs = 2

Therefore, the shape of molecule would be that of slightly bent-T.



- 8. In nylon-6, 6 both the monomers hexamethylene diamine and adipic acid contain six carbon atoms each.
- **9.** Secondary cell.

$$Pb(s)$$
 SO_4^2 (aq)

$$PbSO_4(s) + 2\overline{e}$$

$$PbO_{2}(s)$$
 SO_{4}^{2} (aq) 4H (aq) $2\bar{e}$ $PbSO_{4}(s) + 2H_{2}O(l)$

$$PbSO_4(s) + 2H_2O(l)$$

Cell reaction :
$$Pb(s) + PbO_2(s) + 2H_2SO_4(aq)$$

$$2 \text{ PbSO}_4(s) + 2 \text{H}_2 \text{O}(l)$$

$$\operatorname{Sn}^{2+}(aq)$$

$$Sn^{4+}(aq) + 2e^{-1} \times 5$$

$$MnO_4^-(aq) + 8H^+(aq) + 5e^-$$

$$Mn^{2+}(aq) + 4H_2O(l)] \times 2$$

Cell reaction:
$$2\text{MnO}_4^-(aq) + 5\text{Sn}^{2+}(aq) + 16\text{ H}^+(aq)$$
 $2\text{Mn}^{2+}(aq) + 5\text{Sn}^{4+}(aq) + 8\text{H}_2\text{O}(l)$

$$2Mn^{2+}(aq) + 5Sn^{4+}(aq) + 8H_2O(l)$$

$$E_{\text{Cell}}^{\text{o}} = E_{\text{cathode}}^{\text{o}} - E_{\text{anode}}^{\text{o}} = 1.51 \text{ V} - 0.15 \text{ V}$$

= 1.36 V

As cell potential is positive therefore the reaction is product favoured.

- 13. (i) This is due to d-d transition. When visible (white) light falls on a transition metal compounds, they absorb certain radiation of visible light and transmit the remaining ones. The colour observed corresponds to complementary colour of the light absorbed.
 - (ii) As manganese has maximum number of unpaired electrons (5) in 3d subshell in addition to 2 electrons in the 4s subshell.
- 18. On the basis of magnitude of intermolecular forces polymers are classified into following four sub-groups:
 - (i) Elastomers

- (ii) Fibers
- (iii) Thermoplastic polymers
- (iv) Thermosetting polymers

Let

19.
$$d = \frac{Z - M}{a^3 - N_A}$$
 $a^3 = \frac{Z - M}{d - N_A}$...(i)

For fcc unit cell, Z = 4

For fcc.

M = 207 g mol⁻¹ N_A =
$$6.02 \times 10^{23}$$
 mol⁻¹
 $d = 11.35$ g cm⁻³

Substituting these values in equation (i), we get

$$a^{3} = \frac{4 \quad 207 \text{ g mol}^{-1}}{11.35 \text{ g cm}^{-3} \quad 6.02 \quad 10^{23} \text{ mol}^{-1}}$$

$$a^{3} = \frac{4 \quad 207 \quad 10}{11.35 \quad 6.02 \quad 10^{24}} \text{ cm}^{3} \qquad a = \frac{8280}{11.35 \quad 6.02} \quad 10^{-8} \text{ cm}$$

$$x = \frac{8280}{11.35 \quad 6.02}$$

$$\log x = \frac{1}{3} \left[\log 8280 - \log 11.35 - \log 6.02 \right] = \frac{1}{3} \left[3.9180 \quad 1.0549 \quad 0.7796 \right]$$

$$\log x = \frac{1}{3} \left[2.0835 \right] = 0.6945$$

$$x = \text{Antilog} (0.6945) \qquad x = 4.949$$

$$a = 4.949 \times 10^{-8} \text{ cm} \qquad a = 494.9 \text{ pm}$$

$$r = \frac{a}{2\sqrt{2}}$$

$$r = \frac{494.9}{2\sqrt{2}} \text{ pm} = \frac{494.9\sqrt{2}}{4} \text{ pm} = \frac{494.9 \quad 1.414}{4} \text{ pm}$$

r = 174.95 pm **26.** (i) CH₃CH₂Cl NaCN CH₃CH₂CN Reduction Ni/H₂ CH₃CH₂CH₂NH₂

- 27. (i) Hard water contains calcium and magnesium salts. In hard water, soaps get precipitated as calcium and magnesium soaps which being insoluble stick to the clothes as gummy mass.
 - (ii) Chloroxylenol and -terpineol in a suitable solvent.
 - (iii) Antiseptics are chemical substances which prevent the growth of microorganisms and may even kill them but are not harmful to human or animal tissues. Examples are dettol and savlon. They are generally applied on wounds, ulcers, cuts and diseased skin surfaces. Furacin and soframycin are well-known antiseptic creams.

Disinfectants are chemical substances which kill microorganisms but are not safe to be applied to the living tissues. These are generally used to kill microorganisms present in the toilets, drains, floors, etc. Some common examples of disinfectants are phenol (1% solution) and chlorine (0.2 to 0.4 ppm).

CBSE EXAMINATION PAPERS

ALL INDIA-2011

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question numbers 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question numbers 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question numbers 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question numbers 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables, if necessary. Use of calculators is not allowed.

CBSE (All India) SET-I

- 1. Define 'order of a reaction'.
- 2. What is meant by 'shape selective catalysis'?
- 3. Differentiate between a mineral and an ore.
- **4.** What is meant by 'lanthanoid contraction'?
- **5.** Write the IUPAC name of the following compound:

 $CH_2 = CHCH_2Br$

- **6.** Draw the structure of 4-chloropentan-2-one.
- 7. How would you convert ethanol to ethene?
- **8.** Rearrange the following in an increasing order of their basic strengths:

 $C_6H_5NH_2$, $C_6H_5N(CH_3)_2$, $(C_6H_5)_2$ NH and CH_3NH_2

- **9.** Explain how you can determine the atomic mass of an unknown metal if you know its mass density and the dimensions of unit cell of its crystal.
- **10.** Calculate the packing efficiency of a metal crystal for a simple cubic lattice.
- 11. State the following:
 - (i) Raoult's law in its general form in reference to solutions.
 - (ii) Henry's law about partial pressure of a gas in a mixture.
- **12.** What do you understand by the rate law and rate constant of a reaction? Identify the order of a reaction if the units of its rate constant are:
 - (i) $L^{-1} \text{ mol s}^{-1}$
 - (ii) L mol^{-1} s⁻¹.
- 13. The thermal decomposition of HCO₂H is a first order reaction with a rate constant of 2.4×10^{-3} s⁻¹ at a certain temperature. Calculate how long will it take for three-fourths of initial quantity of HCO₂H to decompose. (log 0.25 = -0.6021)

- 14. Describe the principle controlling each of the following processes:
 - (i) Vapour phase refining of titanium metal.
 - (ii) Froth floatation method of concentration of a sulphide ore.
- **15.** How would you account for the following:
 - (i) Cr^{2+} is reducing in nature while with the same d-orbital configuration (d⁴) Mn^{3+} is an oxidising agent.
 - (ii) In a transition series of metals, the metal which exhibits the greatest number of oxidation occurs in the middle of the series.
- **16.** Complete the following chemical equations:
 - (i) $MnO_4^-(aq) + S_2O_3^{2-}(aq) + H_2O(l)$
 - (ii) $Cr_2O_7^{2-}(aq) + Fe^{2+}(aq) + H^+(aq)$

OR

State reasons for the following:

- (i) Cu(I) ion is not stable in an aqueous solution.
- (ii) Unlike Cr^{3+} , Mn^{2+} , Fe^{3+} and the subsequent other M^{2+} ions of the 3d series of elements, the 4d and the 5d series metals generally do not form stable cationic species.
- **17.** Explain what is meant by the following:
 - (i) peptide linkage
 - (ii) pyranose structure of glucose.
- **18.** Write the main structural difference between DNA and RNA. Of the four bases, name those which are common to both DNA and RNA.
- **19.** A solution prepared by dissolving 8.95 mg of a gene fragment in 35.0 mL of water has an osmotic pressure of 0.335 torr at 25°C. Assuming that the gene fragment is a non-electrolyte, calculate its molar mass.
- **20.** Classify colloids where the dispersion medium is water. State their characteristics and write an example of each of these classes.

OR

Explain what is observed when

- (a) an electric current is passed through a sol.
- (b) a beam of light is passed through a sol.
- (c) an electrolyte (say NaCl) is added to ferric hydroxide sol.
- **21.** How would you account for the following:
 - (i) H₂S is more acidic than H₂O.
 - (ii) The N—O bond in NO_2 is shorter than the N—O bond in NO_3 .
 - (iii) Both O₂ and F₂ stabilise high oxidation states but the ability of oxygen to stablise the higher oxidation state exceeds that of fluorine.
- **22.** Explain the following terms giving a suitable example in each case:
 - (i) Ambident ligand
 - (ii) Denticity of a ligand
 - (iii) Crystal field splitting in an octahedral field.

- **23.** Rearrange the compounds of each of the following sets in order of reactivity towards S_N² displacement:
 - (i) 2-Bromo-2-methylbutane, 1-Bromopentane, 2-Bromopentane
 - (ii) 1-Bromo-3-methylbutane, 2-Bromo-2methylbutane,
 - 3-Bromo-2-methylbutane
 - (iii) 1-Bromobutane, 1-Bromo-2, 2-dimethylpropane,
 - 1-Bromo-2-methylbutane
- **24.** How would you obtain the following:
 - (i) Benzoquinone from phenol
 - (ii) 2-Methylpropan-2-ol from methylmagnesium bromide
 - (iii) Propan-2-ol from propane
- **25.** State reasons for the following:
 - (i) pK_b value for aniline is more than that for methylamine.
 - (ii) Ethylamine is soluble in water whereas aniline is not soluble in water.
 - (iii) Primary amines have higher boiling points than tertiary amines.
- **26.** Draw the structures of the monomers of the following polymers:
 - (i) Polythene
 - (ii) PVC
 - (iii) Teflon
- 27. What are the following substances? Give one example of each.
 - (i) Food preservatives
 - (ii) Synthetic detergents
 - (iii) Antacids
- **28.** (a) What type of a battery is lead storage battery? Write the anode and cathode reactions and the overall cell reaction occurring in the operation of a lead storage battery.
 - (b) Calculate the potential for half-cell containing

$$0.10 \text{M K}_2 \text{Cr}_2 \text{O}_7 \text{ (aq)}, 0.20 \text{Mcr}^{3+} \text{ (aq)} \text{ and } 1.0 \times 10^{-4} \text{MH}^+ \text{(aq)} \text{ The half cell reaction is}$$

 $\text{Cr}_2 \text{O}_7^{2-} (aq) + 14 \text{H}^+ (aq) + 6 \text{e}^- \qquad 2 \text{Cr}^{3+} (aq) + 7 \text{H}_2 \text{O}(l)$

and the standard electrode potential is given as
$$E^{\circ} = 1.33V$$
.

OR

- (a) How many moles of mercury will be produced by electrolysing 1.0 M. Hg(NO₃)₂ solution with a current of 2.00 A for 3 hours?
- (b) A voltaic cell is set up at 25° C with the following half-cells Al³⁺ (0.001 M) and Ni²⁺ (0.50 M). Write an equation for the reaction that occurs when the cell generates an electric current and determine the cell potential.

(Given:
$$E_{\text{Ni}^2/\text{Ni}}^{\text{o}} = -0.25 \text{ V}, \quad E_{\text{Al}^3/\text{Al}}^{\text{o}} = -1.66 \text{ V})$$

- **29.** (a) Draw the structures of the following molecules:
 - (i) (HPO₃)₃
 - (ii) BrF₃

- (b) Complete the following equations:
 - (i) $HgCl_2 + PH_3$
 - (ii) SO₃ + H₂SO₄
 - (iii) XeF₄ + H₂O

OR

- (a) What happens when
 - (i) chlorine gas is passed through a hot concentrated solution of NaOH?
 - (ii) sulphur dioxide gas is passed through an aqueous solution of a Fe (III) salt?
- (b) Answer the following:
 - (i) What is the basicity of H₃PO₃ and why?
 - (ii) Why does fluorine not play the role of a central atom in interhalogen compounds?
 - (iii) Why do noble gases have very low boiling points?
- **30.** (a) Illustrate the following name reactions:
 - (i) Cannizzaro's reaction
 - (ii) Clemmensen reduction
 - (b) How would you obtain the following:
 - (i) But-2-enal from ethanal
 - (ii) Butanoic acid from butanol
 - (iii) Benzoic acid from ethylbenzene

OR

- (a) Give chemical tests to distinguish between the following:
 - (i) Benzoic acid and ethyl benzoate
 - (ii) Benzaldehyde and acetophenone.
- (b) Complete each synthesis by giving missing reagents or products in the following:

COOH
$$COOH$$

$$COOH$$

$$COOH$$

$$(ii) C_6H_5CHO$$

$$CH_2$$

$$CH_2$$

$$CHO$$

CBSE (All India) SET-II

Questions Uncommon to Set-I

- **2.** What are lyophobic colloids? Give one example for them.
- **3.** Why is it that only sulphide ores are concentrated by froth floatation process?

5. Write the IUPAC name of the following compound:

$$H$$
 H
 H
 H
 H
 H
 H

- **6.** Draw the structure of 2, 6-Dimethylphenol.
- **9.** Define the following terms in relation to crystalline solids:
 - (i) Unit cell
 - (ii) Coordination number

Give one example in each case.

- **12.** A reaction is of second order with respect to a reactant. How is the rate of reaction affected if the concentration of the reactant is reduced to half? What is the unit of rate constant for such a reaction?
- **14.** Describe the principle controlling each of the following processes:
 - (i) Zone refining of metals.
 - (ii) Electrolytic refining of metals.
- **15.** Explain giving a suitable reason for each of the following:
 - (i) Transition metals and their compounds are generally found to be good catalysts.
 - (ii) Metal-metal bonding is more frequent for the 4d and the 5d series of transition metals than that for the 3d series.
- 19. What mass of NaCl must be dissolved in 65.0 g of water to lower the freezing point of water by 7.50°C? The freezing point depression constant (K_f) for water is 1.86 C/m. Assume van't Hoff factor for NaCl is 1.87. (Molar mass of NaCl = 58.5 g).
- 22. Write the structures and names of all the stereoisomers of the following compounds:
 - (i) $[Co(en)_3]Cl_3$
 - (ii) $[Pt(NH_3)_2Cl_2]$
 - (iii) [Fe(NH₃)₄Cl₂]Cl

CBSE (All India) SET-III

Questions Uncommon to Set-I and Set-II

- 1. Define 'activation energy' of a reaction.
- **2.** What is meant by 'reverse osmosis'?
- 3. What type of ores can be concentrated by magnetic separation method?
- 11. Differentiate between molality and molality values for a solution. What is the effect of change in temperature on molarity and molality values?
- **14.** Describe the principle controlling each of the following processes:
 - (i) Preparation of cast iron from pig iron.
 - (ii) Preparation of pure alumina (Al₂O₃) from bauxite ore.

- **15.** Explain giving reasons:
 - (i) Transition metals and their compounds generally exhibit a paramagnetic behaviour.
 - (ii) The chemistry of actinoids is not as smooth as that of lanthanoids.
- 18. Write such reactions and facts about glucose which cannot be explained by open chain structure.
- 21. How would you account for the following:
 - (i) NF₃ is an exothermic compounds but NCl₃ is not.
 - (ii) The acidic strength of compounds increases in the order:

$$PH_3 < H_2S < HCl$$

- (iii) SF₆ is kinetically inert.
- **22.** Write the state of hybridization, the shape and the magnetic behaviour of the following complex entities:
 - (i) [Cr(NH₃)₄ Cl₂]Cl
 - (ii) [Co(en)₃]Cl₃
 - (iii) K₂ [Ni (CN)₄]
- **26.** Write the names and structures of the monomers of the following polymers:
 - (i) Buna-S
 - (ii) Dacron
 - (iii) Neoprene



CBSE (All India) SET-I

- 1. Order of a reaction may be defined as the sum of the powers of the concentration terms of the reactants in the rate law expression.
- 2. The catalysis in which the pore structure of the catalyst and the size of the reactant and product molecules are comparable is called shape selective catalysis.
- The naturally occurring chemical substances present in the earth's crust which can be obtained by mining are called minerals, while mineral from which metals can be extracted economically are called ores.
- **4.** The steady decrease in the atomic and ionic radii (having the same charge) with increase in atomic number as we move across the series from lanthanum to lutetium is known as lanthanoid contraction.
- **5.** 3-Bromoprop-l-ene

6.
$$\begin{array}{cccc} & & \text{Cl} & \text{O} \\ & & | & | \\ \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{C} - \text{CH}_3 \end{array}$$

7.
$$CH_3CH_2OH$$
 conc. H_2SO_4 $CH_2 = CH_2 + H_2O$ Ethanol Ethene

8. $(C_6H_5)_2 NH < C_6H_5NH_2 < C_6H_5N(CH_3)_2 < CH_3NH_2$

9. We can determine the atomic mass of an unknown metal by using the formula of density of its unit cell.

$$d ext{ (density)} = \frac{Z ext{ (No. of atoms per unit cell)} ext{ } M ext{ (atomic mass)}}{a^3 ext{ (cell edge)} ext{ } N_A ext{ (Avogadro number)}$$

By knowing d, Z, a and N_A , we can calculate M, the atomic mass of metal.

10. Packing efficiency = $\frac{Z \text{ volume of one atom}}{\text{Volume of cubic unit cell}}$ 100

$$= \frac{Z + \frac{4}{3}r^3}{a^3} = 100$$

For a simple cubic lattice, a = 2r and Z = 1

Packing efficiency =
$$\frac{1}{3} = \frac{r^3}{(2r)^3} = 100 = \frac{1}{6} = 100$$

= $52.36\% = 52.4\%$

- 11. (i) Raoult's law: It states that for any solution, the partial pressure of each volatile component in the solution is directly proportional to its mole fraction.
 - (ii) Henry's law: It states that the partial pressure of a gas in vapour phase (P) is proportional to its mole fraction (x) in the solution.
- 12. An experimentally determined expression which relates the rate of reaction with the concentration of reactants is called rate law while the rate of reaction when concentration of each reactant is unity in a rate law expression is called rate constant.
 - (i) Comparing power of mole in L^{-1} mol s^{-1} and (mol L^{-1})¹⁻ⁿ s^{-1} ,

We get

$$1 = 1 - n$$
 $n = 0$ *i.e.*, zero order reaction

(ii) Again comparing power of mole in L mol^{-1} s⁻¹ and $(\text{mol L}^{-1})^{1-n}$ s⁻¹, we get

$$-1 = 1 - n$$
 $n = 2$, *i.e.*, second order reaction

13. For a first order reaction

$$t = \frac{2.303}{k} \log \frac{[A]_o}{[A]}$$

Here,
$$k = 2.4 \times 10^{-3} \text{ s}^{-1}$$
, $[A] = [A]_0 \frac{3}{4} [A]_0 = \frac{[A]_0}{4}$, $t = ?$

Substituting these values in the equation, we get

$$t = \frac{2.303}{2.4 \cdot 10^{-3} \text{ s}^{-1}} \log \frac{\text{[A]o}}{\text{[A]}_{\frac{9}{4}}}$$

$$t = \frac{2.303}{2.4 \cdot 10^{-3} \text{ s}^{-1}} \log 4$$

$$= \frac{2.303}{2.4 \cdot 10^{-3}} \cdot 0.6021 \text{ s}$$

$$t = 577.7 \text{ s} = 578 \text{ s}$$

14. (i) In this method, the metal is converted into its volatile compound and collected elsewhere. It is then decomposed to give pure metal.

$$\begin{array}{cccc} {\rm Ti} & {\rm ZI}_2 & {\rm ^{500\,K}} & {\rm Ti\,I_4} \\ {\rm Impure} & & & & & & \\ {\rm Ti\,I_4} & {\rm ^{1700\,K}} & {\rm Ti} & {\rm _{Pure\,metal}} \end{array}$$

- (ii) This method of concentration of ore is based upon the principle that the surface of sulphide ores is preferentially wetted by oils while that a gangue is preferentially wetted by water.
- (i) Cr^{2+} is reducing as its configuration changes from d^4 to d^3 , a more stable half filled t_{2g} configuration while Mn³⁺ is oxidising as Mn³⁺ to Mn²⁺results a more stable half filled d⁵ configuration.
 - (ii) It is due to greater number of unpaired electrons in (n-1)d and ns orbitals at the middle of the series.
- **16.** (i) $8MnO_4^-(aq) + 3S_2O_3^{2-}(aq) + H_2O(l)$ $8MnO_2(s) + 6SO_4^{2-}(aq) + 2OH^-(aq)$
 - (ii) $Cr_2O_7^{2-}(aq)$ 14H (aq) 6Fe² (aq) 2Cr³ (aq) 6Fe³ (aq) 7H₂O(l)

OR


(i) This is because Cu(I) ion is unstable in aqueous solution and undergo disproportionation.

2Cu
$$(aq)$$
 Cu 2 (aq) Cu (s)

(ii) This is because due to lanthanoid contraction the expected increase in size does not occur.



- 17. (i) Peptide linkage: The amide (-C-NH-) linkage between two -amino acids formed with the loss of a water molecule is called a peptide linkage.
 - (ii) The six membered cyclic structure of glucose is called pyranose structure (), in analogy with heterocylic compound pyran.



 $\alpha - D - (+) - Glucopyranose$

18. Structural difference between DNA and RNA

DNA	RNA	
1. The sugar present in DNA is 2-deoxy D-(–) ribose.	1. The sugar present in RNA is D-(–)-ribose.	
2. DNA has double stranded -helix structure.	2. RNA has single -helix structure.	

The common bases present in both DNA and RNA are adenine (A), guanine (G) and cytosine (C).

19.
$$M_{B} = \frac{W_{B} \quad R \quad T}{V} \qquad(i)$$

Here,
$$W_B = 8.95 \text{ mg} = 8.95 \times 10^{-3} \text{g}$$
 , $R = 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$

$$T = 25^{\circ}C = (25 + 273) K = 298 K$$
, $= 0.335 torr = \frac{0.335}{760} atm$

$$V = 35 \text{ mL} = 35 \times 10^{-3} \text{ L}$$

Substituting these values in the equation (i), we get

$$M_B = \frac{8.95 \cdot 10^{-3} \text{ g} \cdot 0.0821 \text{ L atm mol}^{-1} \text{K}^{-1} \cdot 298 \text{K}^{-7} \cdot 760}{0.335 \text{ atm}^{-3} \cdot 35^{-1} \cdot 10^{-3} \text{L}} = 14193.3 \text{ g mol}^{-1}$$

20. These are of two types

(i) Hydrophilic

Stability: More stable as the stability is due to charge and water envelope surrounding the sol particles.

Nature: Reversible

Examples: Starch, gum, etc.

(ii) Hydrophobic

Stability: Less stable as the stability is due to charge only.

Nature: Irreversible

Examples: Metal hydroxide like Fe(OH)₃ and metal sulphide like As₂S₃.


OR

- (i) On passing electric current through a sol, colloidal particles start moving towards oppositely charged electrode where they lose their charge and get coagulated (electrophoresis).
- (ii) Scattering of light by the colloidal particles takes place and the path of light becomes visible (Tyndall effect).
- (iii) The positively charged colloidal particles of ferric hydroxide sol get coagulated by the oppositely charged Cl⁻ ions provided by NaCl.
- 21. (i) This is because bond dissociation enthalpy of H—S bond is lower than that of H—O bond.
 - (ii) This is because two bonds share a double bond in the resonance hybrid structure of NO_2^- while three bonds share a double bond in the resonance hybrid structure of NO_3^- . This is because NO_2^- has bond order 1.5 while NO_3^- has bond order 1.33.
 - (iii) This is due to tendency of oxygen to form multiple bonds with metal atom.
- 22. (i) Ambidentate ligand: A ligand which can coordinate to central metal atom through two different atoms is called ambidentate ligand. For example NO₂ ion can coordinate either through nitrogen or through oxygen to a central metal atom/ion.
 - (ii) Denticity: The number of coordinating groups present in ligand is called the denticity of ligand. For example, bidentate ligand ethane-1, 2-diamine has two donor nitrogen atoms which can link to central metal atom.

$$H_2 N - CH_2 - CH_2 - NH_2$$

Ethane-1, 2-diamine

- (iii) The splitting of the degenerated d-orbitals into three orbitals of lower energy, t_{2g} set and two orbitals of higher energy eg, set due to the presence of ligand in a octahedral crystal field is known as crystal splitting in a octahedral field.
- **23.** (*i*) 1-Bromopentane > 2-Bromopentane > 2-Bromo-2methyl butane.
 - (ii) 1-Bromo-2-methyl butane > 3-Bromo-2-methyl butane > 2-Bromo-2-methyl butane
 - (iii) 1-Bromobutane > 1-Bromo-2-methylbutane > 1-Bromo-2,2-dimethyl butane.



- 25. (i) In aniline, the lone pair of electrons on N-atom are delocalised over benzene ring due to resonance. As a result, electron density on the nitrogen atom decreases. In contrast, in methylamine, +I-effect of CH₃⁻ group increases electron density on the nitrogen atom. Therefore, aniline is a weaker base than methylamine, hence its pK_b value is more than that for methylamine.
 - (ii) Ethylamine is soluble in water due to formation of inter-molecular hydrogen bonds with water molecules. However, in aniline due to large hydrophobic part, *i.e.*, hydrocarbon part, the extent of hydrogen bonding decreases considerably and hence aniline is insoluble in water.
 - (iii) Due to the presence of two H-atoms on N-atom, primary amines undergo extensive intermolecular hydrogen bonding whereas tertiary amines have no H-atoms on the nitrogen atom, do not undergo H-bonding. As a result, primary amines have higher boiling points than tertiary amines.

26.	(i) Name of monomer	Structure	
	Ethene	$CH_2 = CH_2$	
	Vinyl chloride	$CH_2 = CH-Cl$	
	Tetrafluoroethene	$F_2C = CF_2$	

27. (i) Food Preservatives: These are the chemical substances which are added to the food materials to prevent their spoilage due to microbial growth and to retain their nutritive value for long time. Sodium benzoate, sodium metabisulphite are some common preservatives.

Preservatives prevent rancidity and kill or inhibit the growth of microorganism.

- (ii) Synthetic Detergents: Detergents are cleansing agents which have all the properties of soaps, but actually do not contain any soap *e.g.*, sodium dodecylbenzene sulphonate. These can be used both in soft and hard water as they give foam even in hard water. Detergents are mainly classified into three categories:
 - (i) Anionic detergents
 - (ii) Cationic detergents
 - (iii) Non-ionic detergents
- (iii) Antacids: These are the chemical substances which neutralize the excess acid and raise the pH to an appropriate level in the stomach. Sodium hydrogen carbonate or a mixture of aluminium and magnesium hydroxide are some common antacids.
- **28.** (a) The lead storage battery is a secondary cell.

The cell reactions when the battery is in use are given below

At anode: Pb(s) SO_4^2 (aq) $PbSO_4(s)$ 2e At cathode: $PbO_2(s)$ SO_4^2 (aq) 4H (aq) 2e $PbSO_4(s)$ 2H₂O(l) Overall cell reaction: Pb(s) $PbO_2(s)$ 2H₂SO₄(aq) 2PbSO₄(s) 2H₂O(l)

(b) For half cell reaction

$$\begin{aligned} &\text{Cr}_2 \text{ O}_7^2 \ (aq) & 14\text{H} \ \ (aq) & 6\bar{\text{e}} & 2\text{Cr}^{3+}(aq) & 7\text{H}_2\text{O}(\textit{l}) \\ &E_{\text{cell}} = E_{\text{cell}}^{\text{o}} & \frac{0.0591}{n} \log \frac{[\text{Cr}^3]^2}{[\text{Cr}_2 \text{ O}_7^{2-}] [\text{H}]^{14}} \\ &\text{Here, E}_{\text{Cell}}^{\text{o}} = 1.33 \text{ V, n} = 6, [\text{Cr}^{3+}] = 0.2 \text{ M} \\ &[\text{Cr}_2 \text{ O}_7^2] = 0.1 \text{ M, } [\text{H}^+] = 1 \times 10^{-14} \text{ M} \end{aligned}$$

Substituting these values in the above given expression, we get

$$E_{\text{cell}} = 1.33 \text{ V} - \frac{0.0591}{6} \log \frac{(0.20)^2}{(0.1)(10^{-4})^{14}}$$

$$= 1.33 \text{ V} - \frac{0.0591}{6} \log (4 \times 10^{-55})$$

$$= 1.33 \text{ V} - \frac{0.0591}{6} [\log 4 + \log 10^{55}]$$

$$= 1.33 \text{ V} - \frac{0.0591}{6} [2 \log 2 + 55 \log 10]$$

$$= 1.33 \text{ V} - \frac{0.0591}{6} [2 \times 0.3010 + 55] = 1.33 \text{ V} - 0.548 \text{ V}$$

$$E_{\text{cell}} = 0.782 \text{ V}$$

OR

(a) Mass of mercury produced at the cathode,

$$m = Z \times I \times t = \frac{M - I - t}{n - F}$$

$$m = \frac{\text{M g mol}^{-1} \quad 2\text{A} \quad 3 \quad 60 \quad 60 \text{ s}}{2 \quad 96500 \text{ C mol}^{-1}} = 0.119 \times \text{M g}$$
$$m = 0.119 \times \text{M g}$$

No. of moles of mercury produced = $\frac{0.119 \text{ Mg}}{\text{Mg mol}^{-1}} = 0.119 \text{ mol}$

 $A1^3$ (aq) $3e^-$] 2 (*b*) At anode: Al(s) Ni^2 $2e^-$ Ni (s) At cathode: $2Al(s) + 3Ni^{2+}(aq)$ $2Al^{3+}(aq) + 3Ni(s)$ Cell reaction:


Applying Nerst equation to the above cell reaction

$$E_{cell} = E_{cell}^{o} - \frac{0.0591}{n} \log \frac{[Al^3]^2}{[Ni^2]^3}$$

 $E_{\text{cell}}^{\text{o}} = E_{\text{Ni}^2 / \text{Ni}}^{\text{o}} \quad E_{\text{Al}^3 / \text{Al}}^{\text{o}} = -0.25 \text{ V} - (-1.66 \text{V}) = 1.41 \text{V}, n = 6$ Here, $[Al^{3+}] = 1 \times 10^{-3} M$, Ni = 0.5 M

$$\begin{split} & E_{cell} = 1.41 \text{ V} - \frac{0.0591}{6} \log \frac{(10^{-3})^2}{(0.5)^3} = 1.41 \text{ V} - \frac{0.0591}{6} \log (8^{-10^{-6}}) \\ & = 1.41 \text{ V} - \frac{0.0591}{6} (\log 2^3 + \log 10^{-6}) \\ & = 1.41 \text{ V} - \frac{0.0591}{6} [3 \times \log 2 + (-67) \log 10] \\ & = 1.41 \text{ V} - \frac{0.0591}{6} [3 \times 0.3010 - 6] \\ & = 1.41 \text{ V} + 0.050 \text{ V} \end{split}$$

 $E_{cell} = 1.46 \text{ V}$



(b) (i) $3HgCl_2 + 2PH_3$

 $Hg_3P_2 + 6HCl$


(ii) SO₃ + H₂SO₄

 $H_2S_2O_7$

(iii) $6XeF_4 + 12H_2O$ $4Xe + 2XeO_3 + 24HF + 3O_2$

OR

(a) (i)
$$3\text{Cl}_2 + 6\text{NaOH (conc.)}$$
 $5\text{NaCl} + \text{NaClO}_3 + 3\text{H}_2\text{O}$
(ii) $2\text{Fe}^{3+} + \text{SO}_2 + 2\text{H}_2\text{O}$ $2\text{Fe}^{2+} + \text{SO}_4^{2-} + 4\text{H}^+$



Two, as the structure of H₃PO₄ has two P—OH bonds.

- (ii) This is due to absence of d-orbitals in fluorine.
- (iii) Noble gases being mono atomic gases have no interatomic forces except weak dispersion forces, therefore they have low boiling points.
- **30** (a) (i) Cannizzaro reaction: Aldehydes which do not have an -hydrogen atom undergoes disproportionation reactions on treatment with concentrated alkali to give a mixture of carboxylic acid salt and alcohol.


2 HCHO Formaldehyde Conc. NaOH HCOO Na Sod. formate CH₃ OH Methyl alcohol

(ii) Clemmensen reduction: The carbonyl group of aldehydes and ketones is reduced to CH₂ group on treatment with zinc-amalgam and concentrated hydrochloric acid.

$$\begin{array}{c} C = O \,+\, 4(H) & \begin{array}{c} \operatorname{Zn-Hg} \\ \operatorname{Aldehyde} \\ \operatorname{or} \\ \operatorname{Ketone} \end{array} & CH_2 \,\,+\,\, H_2O \end{array}$$

(b) (i) CH₃CHO
$$\stackrel{\text{dil. NaOH}}{\rightleftharpoons}$$
 CH₃CH—CH₂CHO $\stackrel{\Delta}{\longrightarrow}$ CH₃CH—CHCHO But-2-enal

 $\begin{array}{ccc} \textit{(ii)} & \text{CH}_{3}\text{CH}_{2}\text{CH}_{2}\text{CH}_{2}\text{OH} & & \text{CrO}_{3}-\text{H}_{2}\text{SO}_{4} & & \text{CH}_{3}\text{CH}_{2}\text{CH}_{2}\text{COOH} \\ & \text{Butanoic acid} & & \text{Butanoic acid} \end{array}$



OR

(a) (i) Benzoic acid on warming with sodium hydrogen carbonate gives brisk effervescence of CO₂ gas while ethyl benzoate does not respond to this test

$$C_6H_5COOH$$
 + NaHCO $_3$ $C_6H_5^-$ COONa CO_2 H_2O Benzoic acid Brisk effervescence

(ii) Benzaldehyde and acetophenone

Iodoform test: Acetophenone being a methyl ketone on treatment with I₂/NaOH undergoes Iodoform reaction to give yellow ppt. of iodoform but benzaldehyde does not.

$$C_{6}H_{5}COCH_{3} \quad 3NaOI \quad C_{6}H_{5}COONa \quad CHI_{3} \quad 2NaOH$$

$$Acetophenone \quad SOCI_{2} \quad COCI \quad +SO_{2}+HCI$$

$$(ii) \quad C_{6}H_{5}-CHO \quad H_{2}NCONHNH_{2} \quad C_{6}H_{5}CH = NNHCONH_{2} + H_{2}O$$

$$(iii) \quad CH_{2} \quad (ii) \quad B_{2}H_{6}H_{2}O_{2}/\bar{O}H \quad CHO$$

CBSE (All India) SET-II

- **2.** Lyophobic sols are those sols in which the particles of the disperse phase have little affinity for the particles of the dispersion medium, *e.g.*, sols of metal and their sulphides and hydroxides.
- **3.** This is because the sulphide ore particles are preferentially wetted by oil and gangue particles are preferentially wetted by water.
- **5.** 3-Bromo-2-methylpropene

- **9.** (*i*) **Unit cell:** A unit cell is the smallest portion of a crystal lattice which, when repeated in different direction, generates the entire lattice.
 - (ii) Coordination number: The number of nearest neighbours of any constituent particle in a packing is called its coordination number. The coordination number of an atom in the bcc structure is 8.
- **12.** Consider the reaction nR Products

As the reaction is of second order

Rate,
$$r = k [R]^2$$
 ...(i)

If the concentration of the reactant reduced to half, then

Rate,
$$r = k \frac{R}{2}$$
 ...(ii)

Dividing equation (ii) by (i), we get

$$\frac{r}{r} = \frac{K [R]^2}{4K [R]^2} = \frac{1}{4}$$

$$r = \frac{1}{4}r$$
 i.e., rate of reaction becomes $\frac{1}{4^{th}}$ of the initial rate.

The unit of rate constant is mol⁻¹ L s⁻¹.

- **14.** (*i*) Zone refining is based on the principle that the impurities are more soluble in the melt than in the solid state of the metal.
 - (ii) Electrolytic refining of a metal: In this method, the impure metal is made to act as anode. A strip of same metal in pure form is used as cathode. They are put in a suitable electrolytic bath containing soluble salt of same metal. When electric current is passed the metal from the anode goes into solution as ions due to oxidation while pure metal gets deposited at the cathode due to reduction of metal ions. The voltage applied for electrolysis is such that impurities of more electropositive metals remains in the solution as ions while impurities of the less electropositive metals settle down under the anode as anode mud.

At anode: M Mⁿ n e

At cathode: Mⁿ n e M

15. (*i*) The catalytic activity of transition metals and their compounds is attributed to the following reasons:

Because of their variable oxidation states transition metals form unstable intermediate compounds and provide a new path with lower activation for the reaction.

In some cases, the transition metal provides a suitable large surface area with free valencies on which reactants are adsorbed.

(ii) In general in the same group of d block elements, the 4d and 5d transition element has larger size than that of 3d elements. Thus, the valence electrons are less lightly held and hence can form metal-metal bond more frequently.

19. $T_f = \frac{i - K_f - W_B - 1000}{M_B - W_A}$

Here, i = 1.87, $W_A = 65.0$ g, $T_f = 7.50$ K


 $K_f = 1.86 \text{ K kg mol}^{-1}, M_B = 58.5 \text{ g mol}^{-1}$

Substituting these values in the above equation, we get

$$7.5 \text{ K} = \frac{1.87 - 1.86 \text{ K kg mol}^{-1} - \text{W}_{\text{B}} - 1000 \text{ g kg}^{-1}}{58.5 \text{ g mol}^{-1} - 65.0 \text{ g}}$$

$$W_{\text{B}} = \frac{7.5 - 58.5 - 65.0}{1.87 - 1.86 - 1000} \text{ g} = 8.199 \text{ g}$$

$$W_B = 8.2 g$$



CBSE (All India) SET-III

- 1. The energy required to form the intermediate called activated complex is known as activation energy.

 Activation energy = Threshold energy Average energy of the reactants
- **2. Reverse osmosis:** If a pressure larger than the osmotic pressure is applied on the solution side, the solvent starts to flow from the solution into the pure solvent through the semipermeable membrane. This phenomenon is called reverse osmosis.
- **3.** If either the ore or the gangue is capable of being attracted by magnetic field, then such separation are carried out by magnetic separation.

- 11. Molarity is the number of moles of solute dissolved in one litre of solution whereas molality is the number of moles of solute per kilogram of the solvent. Molarity decreases with increase in temperature as volume increases with increase in temperature. Molality is independent of temperature because mass does not depend on temperature.
- **14.** (*i*) Pig iron is melt with scrap iron and coke using hot air blast. Due to this, impurities such as C, S and P present in the pig iron are removed as CO₂, SO₂ and P₂O₅ and carbon content reduced to about 3%.
 - (ii) Bauxite is soluble in concentrated NaOH solution whereas impurities are not.
- **15.** (i) This is due to presence of unpaired electrons in the (n-1) d orbitals of transition elements.
 - (ii) The chemistry of actinoids is not as smooth as lanthanoid because they are radioactive and show greater number of oxidation states due comparable energies of 5f, 6d and 7s orbitals.
- 18. The following facts and reactions cannot be explained by open chain structure of glucose.
 - (i) Despite having the aldehyde group, glucose does not give 2, 4-DNP test, Shiff's test and it does not form the hydrogen sulphite addition product with NaHSO₃.
 - (ii) The penta-acetate of glucose does not react with hydroxylamine indicating the absence of free aldehydic group.
- **21.** (*i*) This is because bond dissociation energy of F₂ is lower than Cl₂ Moreover, fluorine forms stronger bond with nitrogen due to comparable size.
 - (ii) The acidic strength of compounds increases because of increase in polarity of E—H bond from P—H to H—Cl, which is due to increase in electronegativity of E.
 - (iii) In SF₆, S atom is sterically protected by six F atoms and does not allow water molecules to attack the S atom. Further, F does not have d-orbitals to accept the electrons denoted by H_2O molecules. Due to these reasons, SF₆ is kinetically an inert substance.

22.

Complex ion	Central metal ion/atom	Hybridisation of metal ion involved	Geometry of the complex	Magnetic behaviour
[Cr (NH ₃) ₄ Cl ₂]Cl	Cr ³⁺	d^2sp^3	Octahedral	Paramagnetic
[Co(en) ₃]Cl ₃	Co ³⁺	d^2sp^3	Octahedral	Diamagnetic
[Ni (CO) ₄]	Ni	sp ³	Tetrahedral	Diamagnetic

26. (i)
$$CH_2$$
— CH — CH_2 and CH CH_2 Buta-1, 3-diene Styrene

(iii)
$$CH_2 = C - CH = CH_2$$
 CI
2-chloro-1, 3-butadiene

CBSE EXAMINATION PAPERS

FOREIGN-2011

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Foreign) SET-I

- 1. What is meant by an 'intrinsic semiconductor'?
- 2. State Henry's law about partial pressure of a gas in a mixture.
- **3.** What do you understand by 'denticity of a ligand'?
- **4.** Which will react faster in S_N2 displacement, 1-bromopentane or 2-bromopentane, and why?
- 5. Give the IUPAC name of the following compound:

- **6.** Write the structure of the following compound : 3-oxopentanal.
- 7. Why is an alkylamine more basic than ammonia?
- **8.** What is meant by a 'broad spectrum antibiotic'?
- **9.** Differentiate between molarity and molality of a solution. Explain how molarity value of a solution can be converted into its molality.
- **10.** A 0.561 m solution of an unknown electrolyte depresses the freezing point of water by 2.93°C. What is Van't Hoff factor for this electrolyte? The freezing point depression constant (K_f) for water is 1.86°C kg mol⁻¹.
- 11. Determine the values of equilibrium constant (K_c) and G° for the following reaction:

- **12.** Define the following terms giving an example of each:
 - (i) Emulsion (ii)
 - (ii) Hydrosol
- 13. Explain how to phenomenon of adsorption finds application in the following processes:
 - (i) Production of vacuum
- (ii) Heterogeneous catalysis
- **14.** How would you account for the following:
 - (i) The following order of increase in strength of acids:

$$PH_3 < H_2S < HCI$$

(ii) The oxidising power of oxoacids of chlorine follows the order:

$$HClO_4 < HClO_3 < HClO_2 < HClO$$

- **15.** Name the following coordination compounds and draw their structures:
 - (i) $[CoCl_2 (en)_2]Cl$
 - (ii) $[Pt(NH_3)_2Cl(NO_2)]$ (At. no. Co = 27, Pt = 78)
- **16.** Explain what is meant by the following:
 - (i) Peptide linkage
 - (ii) Pyranose structure of glucose
- **17.** Name the products of hydrolysis of (*i*) sucrose and (*ii*) lactose.

OR

Mention three such properties of glucose which cannot be explained by its open chain structure.

- **18.** State the reason in each of the following cases:
 - (i) Soaps do not work well in hard water.
 - (ii) Synthetic detergents are better than soaps.
- 19. Aluminium crystallises in a cubic close-packed structure. Radius of the atom in the metal is 125 pm.
 - (i) What is the length of the side of the unit cell?
 - (ii) How many unit cells are there in 1 cm³ of aluminium?
- **20.** A voltaic cell is set up at 25°C with the following half-cells, Al³⁺ (0.001 M) and Ni²⁺ (0.50 M). Write the cell reaction when the cell generates an electric current and determine the cell potential.

(Given:
$$E_{Ni^2/Ni}^o = 0.25 \text{ V}, E_{Al^3/Al}^o = 1.66 \text{ V}$$
)

- **21.** State the principle on which each of the following processes operates:
 - (i) Recovery of silver after the silver ore has been leached with NaCN.
 - (ii) Electrolytic refining of a metal.
 - (iii) Vapour phase refining of a metal.
- **22.** Complete the following chemical equations:
 - (i) NaOH Cl₂
 - (ii) XeF₄ + O₂F₂ 143 K
 - (iii) Br₂ F₂ (excess)
- 23. (a) Mention the optimum conditions for the industrial manufacture of ammonia by Haber's process.
 - (b) Explain the following giving appropriate reasons:
 - (i) Sulphur vapour exhibits paramagnetic behaviour.
 - (ii) Red phosphorus is less reactive than white phosphorus.

OR

Draw the structures of the following molecules:

- (*i*) NF₃
- (ii) H₂S₂O₈
- (iii) H₃PO₃

24. Complete the following reaction equations:

$$(i) \begin{array}{c} \text{CH}_3 \\ + \text{ HI} \\ \\ (ii) \end{array} \begin{array}{c} \text{H} \\ + \text{ HBr} \end{array}$$

- (iii) CH₃CH₂CH = CH₂ + HBr
- 25. Illustrate the following reactions giving a chemical equation in each case:
 - (i) Gabriel phthalimide synthesis
 - (ii) A coupling reaction
 - (iii) Hoffmann's bromamide reaction
- 26. How would you obtain
 - (i) Benzoquinone from phenol?
 - (ii) Propan-2-ol from propene?
 - (iii) 2-Methylpropan-2-ol from methyl magnesium bromide?
- **27.** Mention two important uses for each of the following polymers:
 - (i) Bakelite
- (ii) Nylon 6,6
- (iii) PVC
- 28. (a) Express clearly what you understand by 'rate expression' and 'rate constant' of a reaction.
 - (b) Nitrogen pentoxide decomposes according to the equation

$$2\mathrm{N}_2\mathrm{O}_5(g)$$

$$4NO_2(g)$$

$$O_2(g)$$

This first order reaction was allowed to proceed at 40°C and the data given below were collected:

$[N_2O_5](M)$	Time (min)
0.400	0.00
0.289	20.00
0.209	40.00
0.151	60.00
0.109	80.00

- (i) Calculate the rate constant for the reaction. Include units with your answer.
- (ii) Calculate the initial rate of reaction.
- (iii) After how many minutes will $[N_2O_5]$ be equal to 0.350 M?

OR

- (a) Define
 - (i) Order of a reaction.
 - (ii) Elementary step in a reaction
- (b) A first order reaction has a rate constant value of 0.00510 min⁻¹.

If we begin with 0.10 M concentration of the reactant, how much of the reactant will remain after 3.0 hours?

- **29.** (a) Complete the following reactions in an aqueous medium:
 - (i) $MnO_4^- C_2O_4^2 H$
 - (ii) $Cr_2O_7^2$ H_2S H
 - (b) How would you account for the following:
 - (i) Metal-metal bonding is more extensive in the 4d and 5d series of transition elements than the 3d series.
 - (ii) Mn (III) undergoes disproportionation reaction easily.
 - (iii) Co (II) is easily oxidised in the presence of strong ligands.

OR

- (a) Complete the following chemical equations:
 - (i) $Fe^{3+} + I^{-}$
 - (ii) CrO_4^2 H
- (b) Explain the following:
 - (i) Copper (I) ion is not stable in an aqueous solution.
 - (ii) With same (d⁴) configuration Cr (II) is reducing whereas Mn (III) is oxidising.
 - (iii) Transition metals in general act as good catalysts.
- **30.** (a) Give simple chemical tests to distinguish between the following:
 - (i) Propanal and propanone
 - (ii) Benzaldehyde and acetophenone
 - (b) How would you obtain
 - (i) But-2-enal from ethanal?
 - (ii) Butanoic acid from butanol?
 - (iii) Benzoic acid from ethylbenzene?

OR

- (a) Describe the following reactions giving a chemical equation in each case:
 - (i) Cannizzaro's reaction
 - (ii) Decarboxylation reaction
- (b) Complete the following chemical equations:

$$(i) \begin{tabular}{c} $\operatorname{CH}_2\operatorname{CH}_3$ \\ & & & & & \\ \operatorname{KOH, Heat} \\ \\ (ii) \begin{tabular}{c} COOH \\ & & & & \\ \operatorname{COOH} \\ \\ \end{tabular}$$

(iii) C₆H₅CONH₂ H₃O heat

CBSE (Foreign) SET-II

Questions uncommon to Set-I

- 1. "Crystalline solids are anisotropic in nature." What does it mean?
- 2. State Henry's law about partial pressure of a gas in a mixture.
- **3.** Why is CO a stronger ligand than Cl⁻?
- **8.** What are antiseptics? Give an example.
- 12. Write four distinguishing features operative between chemisorption and physisorption.
- 16. State what you understand by primary and secondary structures of proteins.
- 17. Draw the structures of isomers, if any, and write the names of the following complexes:
 - (i) $[Cr (NH_3)_4 Cl_2]^+$
 - (*ii*) $[Co(en)_3]^{3+}$
- **18.** Explain the following terms giving an example of each:
 - (i) Antacids
 - (ii) Sweetening agents
- **21.** Write the reactions involved in the following processes:
 - (i) Leaching of bauxite ore to prepare pure alumina.
 - (ii) Refining of zirconium by van Arkel method.
 - (iii) Recovery of gold after gold ore has been leached with NaCN solution.
- **22.** Complete the following chemical equations:
 - $(i) P_4 + SO_2Cl_2$
 - (*ii*) $Fe^{3+} + SO_2 + H_2O$
 - (iii) XeF_6 H_2O (excess)
- **27.** (a) What does the designation '6,6' in nylon 6, 6 polymer mean?
 - (b) Which polymer is obtained when free radical polymerisation of chloroprene occurs? Write the structure of the polymer thus obtained.

CBSE (Foreign) SET-III

Questions uncommon to Set-I and Set-II

- 1. How many atoms are there in one unit cell of a body centred cubic crystal?
- 2. State Raoult's law in its general form with respect to solutions.
- **3.** Name the following coordination compound:

$$K_3[CrF_6]$$

- **4.** Which will react faster in S_N1 displacement, 1-bromobutane or 2-bromobutane, and why?
- **12.** Define the following terms:
 - (i) Aerosol
 - (ii) Coagulation of colloids

- **14.** Explain giving a reason for each of the following situation:
 - (i) In aqueous medium, HCl is a stronger acid than HF.
 - (ii) White phosphorus is more reactive than red phosphorous.
- 15. Give the name, the stereochemistry and the magnetic behaviour of the following complexes:
 - (i) [Co (NH₃)₅ Cl]Cl₂
 - (ii) K₂[Ni (CN)₄]
- 19. Silver crystallises with face-centred cubic unit cell. Each side of this unit cell has a length of 409 pm. What is the radius of silver atom? Assume the atoms just touch each other on the diagonal across the face of the unit cell.
- 22. Complete the following chemical equations:
 - (i) $C + H_2SO_4$ (conc.)
 - (ii) P₄ + NaOH + H₂O
 - (iii) Cl₂ F₂ (excess
- **26.** How would you obtain
 - (i) Benzoquinone from phenol?
 - (ii) Propan-2-ol from propene?
 - (iii) 2- Methylpropan-2-ol from methyl magnesium bromide?
- **27.** What are addition polymers? How are the two types of addition polymers different from each other? Give one example of each type.



CBSE (Foreign) SET-I

- 1. Pure substances exhibiting conductivity similar to that of silicon and germanium are called intrinsic semiconductors.
- **2. Henry's law:** It states that the partial pressure of a gas in vapour phase (P) is proportional to its mole fraction (x) in the solution.
- **3.** Denticity: The number of coordinating groups present in ligand is called the denticity of ligand. For example, bidentate ligand ethane-1, 2-diamine has two donor nitrogen atoms which can link to central metal atom.

$$H_2 N - CH_2 - CH_2 - NH_2$$

Ethane-1, 2-diamine

- **4.** 1–Bromopentane, as it is a primary alkyl halide.
- **5.** 2–Bromo –3– methyl-but –2–en–1–ol.

6.
$$CH_3 - CH_2 - C - CH_2 - C - H$$

7. R N H
$$\rightleftharpoons$$
 R $=$ R $=$ H $=$ H $=$ H $=$ H $=$ H

Due to +I effect alkyl group pushes electron towards nitrogen and thus makes the lone pair of electrons more available for sharing with the proton of the acid. Hence alkyl amine is more basic than ammonia.

- **8.** The antibiotic which is effective against a wide range of microorganisms is known as broad spectrum antibiotic. For example, chlorophenicol.
- **9.** Molarity (M) is the number of moles of solute dissolved in one litre of solution whereas molality (m) is the number of moles of the solute per thousand grams of solvent.

If M_B is the molar mass of solute, d is the density of solution then molarity (M) value of a solution can be converted into its molality (m) by using the following for formula.

$$m = \frac{1000 \text{ M}}{1000 \text{ d} \text{ M} \text{ M}_{B}}$$

10.
$$T_f = K_f m$$

$$K_f = 1.86$$
°C kg mol⁻¹, m = 0.561 mol kg⁻¹

Substituting these values in the above equation,

We get,

$$T_f = 1.86 \,^{\circ}\text{C kg mol}^{-1} \times 0.561 \,\text{mol kg}^{-1} = 1.04 \,^{\circ}\text{C}$$
(T_f) Calculated = 1.04 $^{\circ}\text{C}$, (T_f) Observed = 2.93 $^{\circ}\text{C}$

$$i = \frac{\text{Observed colligative property}}{\text{Calculated colligative property}} = \frac{(T_f) \,\text{observed}}{(T_f) \,\text{calculated}}$$

$$i = \frac{2.93 \,^{\circ}\text{C}}{1.04 \,^{\circ}\text{C}} = 2.82.$$

11.
$$Ni(s) + 2Ag^{+}(aq)$$
 $Ni^{2+}(aq) + 2Ag(s)$; $E^{\circ} = 1.05 \text{ V}$

Here, n = 2

$$\log K_c = \frac{n}{0.059} \quad E_{\text{cell}}^{\text{o}}$$

$$\log K_c = \frac{2}{0.059} \quad 1.05 = 39.5932$$

$$K_c = \text{antilog } 39.5932 = 3.919 \times 10^{39}$$

$$K_c = 3.92 \times 10^{39}$$


$$G^{\circ} = -nFE_{\text{cell}}^{\circ}$$

$$G^{\circ} = -2 \times 96500 \times 1.05 = -202650 \text{ J}$$


$$G^{\circ} = -202.65 \text{ kJ}$$

- **12.** (*i*) **Emulsion:** An emulsion is a colloidal system in which both the dispersed phase and the dispersion medium are liquids (e.g., milk, Cod liver, oil, etc.)
 - (ii) **Hydrosol:** A sol in which solid is the dispersed phase and water in dispersion medium is called hydrosol.

- 13. (i) Production of Vacuum: Adsorption can be successfully applied to create conditions of high vacuum. For this a bulb of charcoal cooled in liquid air, is connected to vessel which has already been exhausted as far as possible by vacuum pump. The remaining traces of air inspite of low pressure are adsorbed by the charcoal almost completely.
 - (ii) Heterogeneous Catalysis: There are many gaseous reactions of industrial importance involving solid catalyst. Manufacture of ammonia using iron as a catalyst, manufacture of H₂SO₄ by contact process using V₂O₅ catalyst and use of finely divided nickel in the hydrogenation of vegetable oils are the excellent examples. The gaseous reactants are adsorbed on the surface of the solid catalyst. As a result, the concentration of the reactants increases on the surface of the catalyst and hence the rate of reaction increases.
- **14.** (*i*) Larger the difference in electronegativity greater will be polarity and hence greater will be acidic character.
 - (ii) As HClO₄ is most stable and tendency to give oxygen is least while HClO is least-stable and gives oxygen most easily.
- **15.** (i) Dichloridobis (ethane–1, 2–diamine) cobalt (III) chloride



(ii) Diamminechloridonitrito- N platinum (II)



0

- **16.** (*i*) Peptide linkage: The amide (—C—NH—) linkage between two -amino acids formed with the loss of a water molecule is called a peptide linkage.
 - (ii) The six membered cyclic structure of glucose is called pyranose structure (or), in analogy with heterocylic compound pyran.

 $\alpha - D - (+) - Glucopyranose$

17. (i)
$$C_{12}H_{22}O_{11} + H_2O$$
 H^+ $C_6H_{12}O_6 + C_6H_{12}O_6$ Sucrose Glucose Fructose

OR

The following facts and reactions cannot be explained by open chain structure of glucose.

- (i) Despite having the aldehyde group, glucose does give 2, 4-DNP test, Shiff's test and it does not form the hydrogen sulphite addition product with NaHSO₃.
- (ii) The penta-acetate of glucose does not react with hydroxylamine indicating the absence of free aldehydic group.
- 18. (i) Hard water contains calcium and magnesium salts. In hard water, soaps get precipitated as calcium and magnesium soaps which being insoluble stick to the clothes as gummy mass.
 - (ii) As synthetic detergents can be used in hard water as well as acidic solutions.

19. (*i*) For fcc (or ccp),
$$a = 2\sqrt{2}r = 2 \times 1.414 \times 125 \text{ pm}$$

= 354 pm

(ii)
$$a = 354 \text{ pm} = 3.54 \times 10^{-8} \text{ cm}$$

= 354 pm = 354 pm(ii) $a = 354 \text{ pm} = 3.54 \times 10^{-8} \text{ cm}$ Volume of one unit cell = $a^3 = (3.54 \times 10^{-8} \text{ cm})^3 = 4.436 \times 10^{-23} \text{cm}^3 = 4.44 \text{ cm}^3$

Number of unit cells = $\frac{\text{Total volume}}{\text{Volume of one unit cell}}$

$$= \frac{1 \text{ cm}^3}{4.44 \quad 10^{-23} \text{ cm}^3} = 2.25 \times 10^{22}$$

20. At anode: Al (s) Al
3
 (aq) 3 e $^-$] 2 At cathode: Ni 2 2 e $^-$ N_i(s)] × 3

Cell reaction:
$$2Al(s) + 3Ni^{2+}(aq)$$
 $2Al^{3+}(aq) + 3Ni(s)$

Applying Nerst equation to the above cell reaction

$$E_{cell} = E_{cell}^{o} - \frac{0.0591}{n} \log \frac{[Al^3]^2}{[Ni^2]^3}$$

Here,
$$E_{\text{cell}}^{o} = E_{\text{Ni}^{2}/\text{Ni}}^{o}$$
 $E_{\text{Al}^{3}/\text{Al}}^{o} = -0.25 \text{ V} - (-1.66 \text{V}) = 1.41 \text{V}, n = 6$

$$[Al^{3+}] = 1 \times 10^{-3} M$$
, Ni = 0.5 M

$$\begin{split} E_{\text{cell}} &= 1.41 \text{ V} - \frac{0.0591}{6} \log \frac{(10^{-3})^2}{(0.5)^3} = 1.41 \text{ V} - \frac{0.0591}{6} \log (8^{-10^{-6}}) \\ &= 1.41 \text{ V} - \frac{0.0591}{6} (\log 2^3 + \log 10^{-6}) \\ &= 1.41 \text{ V} - \frac{0.0591}{6} [3 \times \log 2 + (-67) \log 10] \\ &= 1.41 \text{ V} - \frac{0.0591}{6} [3 \times 0.3010 - 6] \\ &= 1.41 \text{ V} + 0.050 \text{ V} \end{split}$$

 $E_{cell} = 1.46 \text{ V}$

21. (i) Recovery of silver after silver ore was leached with NaCN: During leaching Ag is oxidised to Ag⁺ which then combines with CN⁻ ions to form soluble complex, [Ag(CN)₂]⁻. Silver is then recovered from this complex by displacement method using more electropositive zinc metal.

$$2[Ag(CN)_2]^-(aq) + Zn(s)$$
 $2Ag(s) + [Zn(CN)_4]^{2-}(aq)$

(ii) Electrolytic refining of a metal: In this method, the impure metal is made to act as anode. A strip of same metal in pure form is used as cathode. They are put in a suitable electrolytic bath containing soluble salt of same metal. When electric current is passed the metal from the anode goes into solution as ions due to oxidation while pure metal gets deposited at the cathode due to reduction of metal ions. The voltage applied for electrolysis is such that impurities of more electropositive metals remains in the solution as ions while impurities of the less electropositive metals settle down under the anode as anode mud.

At anode: $M M^n n\bar{e}$ At cathode: $M^n n\bar{e} M$

(iii) Vapour phase refining of a metal: In this method, the metal is converted into its volatile compound and collected elsewhere. It is then decomposed to give pure metal. For example, refining of nickel by Mond process.

$$Ni + 4CO$$
 $^{330\ 350\ K}$ $Ni(CO)_4$ $Ni(CO)_4$ $Ni + 4CO$

22. (*i*) 6NaOH + 3Cl₂ 5NaCl + NaClO₃ + 3H₂O Hot and conc.

23. (a) Optimum conditions for the industrial manufacture of ammonia by Haber's process.

Pressure: 200 bar Temperature: 723–773 K

Catalyst: Finely divided iron and molybdenum as promoter.

(b) (i) In vapour state sulphur partly exists as S_2 molecule which has two unpaired electrons in the antibonding * molecular orbitals like O_2 and, hence exhibits paramagnetic behaviour.

(ii) White phosphorus is more reactive than red phosphorus due to its discrete tetrahedral structure and angular strain. Red phosphorus is less reactive due its polymeric structure.

OR


(i) Total no. of electron pairs around

The central N atom =
$$\frac{1}{2}(5 \ 3) \ 4$$

No. of bond pairs = 3


No. of lone pairs = 1

Therefore, according to VSEPR theory; NF₃ should be pyramidal.



25. (i) Gabriel phthalimide synthesis: This reaction is used for the preparation of aliphatic primary amines. In this reaction, phthalimide is first of all treated with ethanolic KOH to form potassium phthalimide. Potassium phthalimide on treatment with alkyl halide gives N-alkyl phthalimide, which on hydrolysis with dilute hydrochloric acid gives a primary amine as the product.

Br



(ii) Coupling reactions:

Diazonium salts react with aromatic amines in weakly acidic medium and phenols in weakly alkaline medium to form coloured compounds called azo dyes by coupling at *p*-position of amines or phenol. The mechanism is basically that of electrophilic aromatic substitutions where the diazonium ion is electrophile.

N NCl + H OH OH
$$(pH 9-10)$$

N NCl + H OH $(pH 9-10)$

N = N OH + Cl H_2 C

P-Hydroxy-azobenzene $(orange dye)$


N NCl + H NH₂ $(pH 4-5)$

N = N NH₂ + Cl H_2 C

N = N NH₂ + Cl H_2 C

(iii) Hoffman's bromamide reaction:

When a primary acid amide is heated with an aqueous or ethanolic solution of NaOH or KOH and bromine (i.e., NaOBr or KOBr), it gives a primary amine with one carbon atom less.



(iii)
$$CH_3 - CH = CH_2$$
 H_2O

Propene

$$CH_3 - CH - CH_3$$

OH

Propan -2-ol

OMgBr

OH

OH

Propanone

$$CH_3 - C - CH_3$$

OH

Propanone

$$CH_3 - C - CH_3$$

OH

Propanone

$$CH_3 - C - CH_3$$

$$CH_3 - C$$

- (i) It is used for making combs, fountain pen barrels, phonograph records. 27. It is used widely in making electrical goods such as switches, plugs, handles of various utensils.
 - (ii) It is used in making bristles for brushes, ropes. It is used for making carpets and fabrics in textile industry.
 - (iii) It is used in the manufacture of rain coats, hand bags, water pipes, vinyl flooring. It is used for insulating electric wires.
- 28. (a) An experimentally determined expression which relates the rate of reaction with the concentration of reactants is called rate law while the rate of reaction when concentration of each reactant is unity in a rate law expression is called rate constant.
 - (i) Comparing power of mole in L^{-1} mol s^{-1} and (mol L^{-1})¹⁻ⁿ s^{-1} , We get

$$1 = 1 - n$$
 $n = 0$ *i.e.*, zero order reaction

(ii) Again comparing power of mole in L $\text{mol}^{-1}\text{s}^{-1}$ and $(\text{mol L}^{-1})^{1-n}\text{s}^{-1}$, we get

$$-1 = 1 - n$$
 $n = 2$, *i.e.*, second order reaction

(i) When t = 20 min, [A] = 0.289 mol L⁻¹ (b)

Also,
$$[A]_0 = 0.400 \text{ mol } L^{-1}$$

For a first order reaction

$$k = \frac{2.303}{t} \log \frac{[A]}{[A]}$$

$$k = \frac{2.303}{20} \log \frac{0.400}{0.289}$$

$$k = \frac{2.303}{20} \log \frac{4.00}{2.89}$$

$$k = \frac{2.303}{20} [\log 4.00 - \log 2.89]$$

$$k = \frac{2.303}{20} [0.6021 - 0.4609]$$

$$k = \frac{2.303}{20} 0.1412$$

$$k = 2.303 \times 0.00706 = 0.016259 \text{ min}^{-1}$$

$$k = 2.303 \times 0.00706 = 0.016259 \text{ min}^{-1}$$

$$k = 1.6259 \times 10^{-2} \,\mathrm{min}^{-1}$$

(ii) Initial rate, i.e., rate of reaction when
$$t = 0$$

When,
$$t = 0.00 \text{ min}$$
, [A] = 0.400 mol L⁻¹

Also,
$$k = 1.626 \times 10^{-2} \text{ min}^{-1}$$

Initial rate =
$$k$$
 [A] = $1.626 \times 10^{-2} \text{ min}^{-1} \times 0.400 \text{ mol L}^{-1}$
= $6.504 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$.

(iii)
$$t = \frac{2.303}{k} \log \frac{[A]_o}{[A]}$$

Here, $k = 1.6259 \times 10^{-2} \text{ min}^{-1}$, $[A]_0 = 0.400 \text{ M}$, [A] = 0.350 MSubstituting these values in the above equation, we get

$$t = \frac{2.303}{1.6259 \quad 10^{-2}} \log \frac{0.400}{0.350} = \frac{2.303}{1.6259 \quad 10^{-2}} [\log 40 - \log 35]$$

$$= 141.583 [1.6021 - 1.5441]$$

$$= 141.583 \times 0.0580$$

$$= 141.583 \times 0.0580 = 8.21 \text{ min.}$$

- OR
- (i) Order of a reaction may be defined as the sum of the powers of the concentration terms of (*a*) the reactants in the rate law expression.
 - (ii) Elementary step: Each step of a complex reaction is called an elementary step.

(b)
$$t = \frac{2.303}{K} \log \frac{[A]_o}{[A]}$$

Here, K =
$$5.10 \times 10^{-3} \text{ min}^{-1}$$
, t = $3 \times 60 \times 60 \text{ min} = 10800 \text{ min}$
[A]_o = 0.1 M

Substituting these values in the above equation,

We get

$$10800 \min = \frac{2.303}{5.1 \cdot 10^{-3} \min^{-1}} \log \frac{0.1}{[A]}$$
$$\log \frac{0.1}{[A]} = \frac{10800 \cdot 51 \cdot 10^{-3}}{2.303} = 23.9166$$
$$\frac{0.1}{[A]} = \text{Antilog } 23.9166 = 8.255 \times 10^{23}$$
$$[A] = \frac{0.1}{8.255 \cdot 10^{23}} = 1.21 \times 10^{-25} \text{M}$$

29. (a) (i)
$$MnO_4^- + 8H^+ + 5\bar{e}$$

$$Mn^{2+} + 4H_2O] \times 2$$

$$C_2O_4^{2}$$

$$2CO_2 + 2\overline{e}] \times 5$$

(i)
$$MnO_4^- + 8H^+ + 5e$$
 $Mn^{2+} + 4H_2O] \times 2$
 $C_2O_4^{2-}$ $2CO_2 + 2\overline{e}] \times 5$
 $2MnO_4^- + 5C_2O_4^{2-} + 16H^+$ $2Mn^{2+} + 10CO_2 + 8H_2O$
(ii) $Cr_2O_7^{2-} + 14H^+ + 6\overline{e}$ $2Cr^{3+} + 7H_2O$

(ii)
$$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\overline{\text{e}}$$
 $2\text{Cr}^{3+} + 7\text{H}_2\text{O}$

$$H_2S$$
 $2H^+ + S + 2e^-] \times 3$

$$Cr_2O_7^{2-} + 3H_2S + 8H^+$$
 $2Cr^{3+} + 3S + 7H_2O$

(b) (i) The catalytic activity of transition metals and their compounds is attributed to the following reasons:

Because of their variable oxidation states transition metals form unstable intermediate compounds and provide a new path with lower activation for the reaction.

In some cases, the transition metal provides a suitable large surface area with free valencies on which reactants are adsorbed.

In general in the same group of d block elements, the 4d and 5d transition element has larger size than that of 3d elements. Thus, the valence electrons are less lightly held and hence can form metal-metal bond more frequently.

- (ii) Mn³⁺ is less stable and changes to Mn²⁺ which is more stable due to half filled d-orbital configuration. That is why, Mn³⁺ undergoes disproportionation reaction.
- (iii) Co (III) has electric configuration 3d, 4s, i.e., it has three unpaired electron. In the presence of strong ligands, two unpaired electrons in 3d subshell pair up and third unpaired electron shift to higher energy subshell from where it can be easily lost and hence oxidised to Co(III).

OR

(a) (i)
$$2Fe^{3+} + 2I^{-}$$
 $2Fe^{2+} + I_{2}$

(ii)
$$2CrO_4^{2-}$$
 2H $Cr_2O_7^{2-}$ H₂O

(b) (i) In aqueous solution Cu⁺ undergoes disproportionation to form a more stable Cu²⁺ ion.

$$2 \text{ Cu}^+(\text{aq})$$
 $2 \text{ Cu}^{2+}(\text{aq}) + \text{Cu}(\text{s})$

The higher stability of Cu^{2+} in aqueous solution may be attributed to its greater negative $_{hyd}H^{\circ}$ than that of Cu^{+} . It compensates the second ionization enthalpy of Cu involved in the formation of Cu^{2+} ions.

- (ii) Cr^{2+} is reducing as its configuration changes from d^4 to d^3 , a more stable half filled t_{2g} configuration while Mn^{3+} is oxidising as Mn^{3+} to Mn^{2+} results a more stable half filled d^5 configuration.
- (iii) The catalytic activity of transition metals and their compounds is attributed to the following reasons:

Because of their variable oxidation states transition metals form unstable intermediate compounds and provide a new path with lower activation for the reaction.

In some cases, the transition metal provides a suitable large surface area with free valencies on which reactants are adsorbed.

30. (a) (i) Propanal and propanone

Tollen's reagent test: Propanal being an aldehyde reduces Tollen's reagent to silver mirror but propanone being a ketone does not.

(ii) Benzaldehyde and acetophenone

Iodoform test: Acetophenone being a methyl ketone on treatment with I₂/NaOH undergoes Iodoform reaction to give yellow ppt. of iodoform but benzaldehyde does not.

(a) (i) Cannizzaro reaction: Aldehydes which do not have an -hydrogen atom undergoes disproportionation reactions on treatment with concentrated alkali to give a mixture of carboxylic acid salt and alcohol.

2 HCHO
Formaldehyde

Conc. NaOH
Sod. formate

CH₃ OH
Methyl alcohol

(ii) **Decarboxylation:** Carboxylic acids lose carbon dioxide to form hydrocarbons when their sodium salts are heated with sodalime.

$$(b) \quad (i) \qquad \begin{array}{c} \text{CH}_2\text{CH}_3 \\ \text{KMnO}_4 \\ \text{KOH, heat} \end{array} \qquad \begin{array}{c} \text{COOK} \\ \text{KOH, heat} \end{array}$$

$$(ii) \qquad \begin{array}{c} \text{COOH} \\ \text{COOH} \\ \text{COOH} \\ \text{Phthalic acid} \end{array} \qquad \begin{array}{c} \text{COCl}_2 \\ \text{heat} \end{array} \qquad \begin{array}{c} \text{COCl}_2 \\ \text{Phthaloyl chloride} \end{array}$$

$$(iii) \quad \text{C}_6\text{H}_5\text{CONH}_2 \qquad \begin{array}{c} \text{H}_3\text{O} \\ \text{heat} \end{array} \qquad \text{C}_6\text{H}_5\text{COOH} \qquad + \quad \text{NH}_3 \end{array}$$

CBSE (Foreign) SET-II

1. Crystalline solids are anisotropic in nature means some of their physical properties like electrical conductivity, refractive index, etc., are different in different directions.

2. Henry's law: It states that "the partial pressure of the gas in vapour phase (p) is proportional to the mole fraction of the gas (x) in the solution" and is expressed as

$$p \quad K_H x$$

where, K_H is the Henry's law constant.

Significance of K_H. Higher the value of Henry's law constant K_H , the lower is the solubility of the gas in the liquid.

- 3. Because CO has bonds.
- **8.** Chemical substances which prevent the growth of microorganisms or kill them but are not harmful to living tissues are called antiseptics.

For example, Dettol, Savlon, etc.


12.

Chemisorption	Physisorption
It arises due to chemical bond formation.	It arises because of van der waals' forces.
It is irreversible in nature.	It is reversible.
It is highly specific in nature.	It is not specific in nature.
It results into unimolecular layers on adsorbent surface.	It results into multimolecular layers on adsorbent surface under high pressure.


16. Primary structure refers to the sequence in which various amino acids are arranged in a protein while secondary structure refers to the shape in which a long polypeptide chain can exist as a result of

regular folding of the backbone of the polypeptide chain due to hydrogen bonding between — C— and —NH—groups of the peptide bond.

17. (i) Geometrical isomers of $[Cr(NH_3)_4Cl_2]^+$




cis-Tetraamminedichlorido chromium (III) ion




trans-Tetraamminedichlorido chromium (III) ion

(ii) Optical isomers of [Co(en)₃]³⁺



dextro-Tris-(ethane-1,2,-diamine) cobalt (III) ion



laevo-Tris-(ethane-1,2-diamine) cobalt (III) ion

- **18.** (*i*) **Antacids:** These are the chemical substances which neutralize the excess acid and raise the pH to an appropriate level in the stomach. Sodium hydrogen carbonate or a mixture of aluminium and magnesium hydroxide are some common antacids.
 - (ii) **Detergents:** Detergents are cleansing agents which have all the properties of soaps, but actually do not contain any soap, *e.g.*, sodium dodecylbenzene sulphonate. These can be used both in soft and hard water as they give foam even in hard water. Detergents are mainly classified into three categories:
 - (a) Anionic detergents
 - (b) Cationic detergents
 - (c) Non-ionic detergents
- **21.** (i) Leaching of bauxite ore to prepare pure alumina:

$$Al_2O_3(s) + 2NaOH(aq) + 3H_2O(l)$$
 $2Na[Al (OH)_4] (aq)$ $2Na[Al(OH)_4](aq) + CO_2(g)$ $Al_2O_3.xH_2O(s) + 2NaHCO_3 (aq)$ $Al_2O_3.xH_2O(s)$ $Al_2O_3.xH_2O(g)$

(ii) Refining of zirconium by van Arkel method.

$$\operatorname{Zr}(s) + 2\operatorname{I}_{2}(g)$$
 870K $\operatorname{ZrI}_{4}(g)$

$$ZrI_4$$
 (g) $\frac{2075 \text{ K}}{Tungsten \text{ filament}}$ $Zr(s)$ $2I_2$

(iii) Extraction of gold:

$$4\text{Au}(s) + 8\text{CN}^{-}(aq) + 2\text{H}_{2}\text{O}(l) + \text{O}_{2}(g)$$
 $4[\text{Au}(\text{CN})_{2}]^{-}(aq) + 4\text{OH}(aq)$

Gold is recovered from [Au (CN₂)] Complex by displacement method using a more eletropositive metal zinc.


$$2[Au (CN)_2]^- (aq) + Zn (s)$$
 2 Au (s)+ $[Zn (CN^{4+})]^{2-} (aq)$

22. (i)
$$P_4 + 10SO_2Cl_2$$
 4 $PCl_5 + 10SO_2$

(ii)
$$2Fe^{3+} + XeO_3 + 6HFSO_2 + 2H_2O$$
 $2Fe^{2+} + 4H^+ + SO_4^{2-}$

(iii)
$$XeF_6 + 3H_2O$$
 (excess)

- 27. (a) In nylon-6, 6 both the monomers hexamethylene diamine and adipic acid contain six carbon atoms each.
 - (b) Neoprene is obtained by free radical polymerisation of chloroprene.



CBSE (Foreign) SET-III

- 1. 2
- 2. Raoult's law: It states that for any solution the partial pressure of each volatile component in the solution is directly proportional to its mole fraction.
- 3. Potassium fluorido chromate (III).
- 4. 2-bromobutane as secondary carbocation formed is more stable as compared to the primary carbocation in case of 1-bromobutane.
- 12. (i) Aerosols: These are the colloidal systems in which dispersion medium is gas and dispsed, phase is either solid or liquid e.g., smoke, insecticide sprays.
 - (ii) Coagulation: The process of aggregating together the charged colloidal particles so as to change them into large sized particle which ultimately settle as down under the force of gravity as a precipitate is called coagulation.
- 14. (i) Because bond dissociation enthalpy of H—Cl is lower than H—F.
 - (ii) White phosphorus is more reactive than red phosphorus due to its discrete tetrahedral structure and angular strain. Red phosphorus is less reactive due its polymeric structure.
- (i) [Co(NH₃)₅Cl₁]Cl₂: Pentaamminechlorido cobalt (III) chloride

Co(III): [Ar] 3d⁶ 4s^o —d²sp³ hybridisation leads to octahedral shape.

Magnetic behaviour: Diamagnetic.

(ii) K₂[Ni(CN)₄]: Potassium tetracyano nickelate (II)

Ni(II): [Ar] 3d⁸ 4s° —dsp² hybridisation leads to square planar shape.

Magnetic behaviour: Diamagnetic.

19. As the atoms just touch each other on the diagonal across the face of unit cell, therefore

$$b^{2} = a^{2} + a^{2} = 2a^{2}$$

 $b = \sqrt{2}a$ (i)

....(ii) Also, b = r + 2r + r = 4r

From (i) and (ii), we get

$$4r = \sqrt{2}a$$
 $r = \frac{\sqrt{2}}{4}a$ $r = \frac{1.414 \quad 409 \text{ pm}}{4} = 144.58 \text{ pm}$



- **22.** (i) C + $2H_2SO_4$ (conc.) $CO_2(g) + 2H_2O$ (l) + $2SO_2(g)$ (ii) P_4 + 3NaOH + $3H_2O$ PH_3 + $3NaH_2$ PO_2

228 Xam idea Chemistry—XII

27. Polymers which are formed by the repeated addition of monomers molecules possessing double or triple bonds are called the addition polymers.

The two types of addition polymers are:

(i) **Homopolymers:** The addition polymers formed by the polymerisation of a single monomeric species are called homopolymers, *e.g.*, polyethene.

$$\begin{array}{ccc} \text{n CH}_2 = \text{CH}_2 & & -(\text{CH}_2 - \text{CH}_2)_{\overline{n}} \\ & & \text{Ethene} & & \text{Polyethene} \end{array}$$

(ii) Copolymers: The polymers made by addition polymerisation from two different monomers are known as copolymers. *e.g.*, Buna-s.

$$nCH_2 = CH - CH_2 + CH_2 + CH_2 - C$$

CBSE EXAMINATION PAPERS

DELHI-2012

Time allowed: 3 hours Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Delhi) SET-I

- 1. What is meant by 'doping' in a semiconductor?
- 2. What is the role of graphite in the electrometallurgy of aluminium?
- **3.** Which one of PCl₄⁺ and PCl₄⁻ is not likely to exist and why?
- **4.** Give the IUPAC name of the following compound:

- **5.** Draw the structural formula of 2-methylpropan-2-ol molecule.
- **6.** Arrange the following compounds in increasing order of their reactivity in nucleophilic addition reactions: ethanal, propanal, propanone, butanone.
- 7. Arrange the following in decreasing order of their basic strength in aqueous solutions:

- **8.** Define the term 'homopolymerisation' giving an example.
- **9.** A 1.00 molal aqueous solution of trichloroacetic acid (CCl₃COOH) is heated to its boiling point. The solution has the boiling point of 100.18°C. Determine the van't Hoff factor for trichloroacetic acid. (K_b for water = 0.512 K kg mol⁻¹)

OR

Define the following terms:

- (i) Mole fraction
- (ii) Isotonic solutions
- (iii) Van't Hoff factor
- (iv) Ideal solution
- **10.** What do you understand by the 'order of a reaction'? Identify the reaction order from each of the following units of reaction rate constant:
 - (i) L^{-1} mol s^{-1}
 - (*ii*) L mol^{-1} s^{-1} .

- 11. Name the two groups into which phenomenon of catalysis can be divided. Give an example of each group with the chemical equation involved.
- 12. What is meant by coagulation of a colloidal solution? Describe briefly any three methods by which coagulation of lyophobic sols can be carried out.
- **13.** Describe the principle involved in each of the following processes.
 - (i) Mond process for refining of Nickel.
 - (ii) Column chromatography for purification of rare elements.
- 14. Explain the following giving an appropriate reason in each case.
 - (i) O₂ and F₂ both stabilise higher oxidation states of metals but O₂ exceeds F₂ in doing so.
 - (ii) Structures of Xenon fluorides cannot be explained by Valence Bond approach.
- **15.** Complete the following chemical equations:
 - (i) $Cr_2O_7^{2-} + H^+ + I^-$
 - $(ii) \text{ MnO}_4^- + \text{NO}_2^- + \text{H}^+$
- **16.** What is meant by (*i*) peptide linkage (*ii*) biocatalysts?
- 17. Write any two reactions of glucose which cannot be explained by the open chain structure of glucose molecules.
- **18.** Draw the structure of the monomer for each of the following polymers:
 - (i) Nylon-6
 - (ii) Polypropene
- **19.** Tungsten crystallises in body centred cubic unit cell. If the edge of the unit cell is 316.5 pm, what is the radius of tungsten atom?

OR

Iron has a body centred cubic unit cell with a cell dimension of 286.65 pm. The density of iron is 7.874 g cm^{-3} . Use this information to calculate Avogadro's number. (At. mass of Fe = 55.845 u)

- **20.** Calculate the amount of KCl which must be added to 1 kg of water so that the freezing point is depressed by 2 K. (K_f for water = 1.86 K kg mol⁻¹)
- **21.** For the reaction

$$2NO(g) + Cl_2(g)$$
 $2NOCl(g)$,

the following data were collected. All the measurements were taken at 263 K:

Experiment No.	Initial [NO] (M)	Initial [Cl ₂] (M)	Initial rate of disappearance of Cl ₂ (M/min)
1	0.15	0.15	0.60
2	0.15	0.30	1.20
3	0.30	0.15	2.40
4	0.25	0.25	?

- (a) Write the expression for rate law.
- (b) Calculate the value of rate constant and specify its units.
- (c) What is the initial rate of disappearance of Cl₂ in experiment 4?

- 22. How would you account for the following?
 - (i) Many of the transition elements are known to form interstitial compounds.
 - (ii) The metallic radii of the third (5d) series of transition metals are virtually the same as those of the corresponding group members of the second (4d) series.
 - (iii) Lanthanoids form primarily +3 ions, while the actinoids usually have higher oxidation states in their compounds, +4 or even +6 being typical.
- 23. Give the formula of each of the following coordination entities:
 - (i) Co³⁺ ion is bound to one Cl⁻, one NH₃ molecule and two bidentate ethylene diamine (en) molecules.
 - (ii) Ni²⁺ ion is bound to two water molecules and two oxalate ions.

Write the name and magnetic behaviour of each of the above coordination entities.

(At. nos.
$$Co = 27$$
, $Ni = 28$)

- **24.** Although chlorine is an electron withdrawing group, yet it is ortho-, para-directing in electrophilic aromatic substitution reactions. Explain why it is so?
- **25.** Draw the structure and name the product formed if the following alcohols are oxidized. Assume that an excess of oxidizing agent is used.
 - (i) CH₃CH₂CH₂CH₂OH
 - (ii) 2-butenol
 - (iii) 2-methyl-1-propanol
- **26.** Write chemical equations for the following conversions:
 - (i) Nitrobenzene to benzoic acid
 - (ii) Benzyl chloride to 2-phenylethanamine
 - (iii) Aniline to benzyl alcohol
- **27.** What are the following substances? Give one example of each one of them:
 - (i) Tranquillisers
 - (ii) Food preservatives
 - (iii) Synthetic detergents
- **28.** (a) What type of a battery is the lead storage battery? Write the anode and the cathode reactions and the overall reaction occurring in a lead storage battery when current is drawn from it.
 - (b) In the button cell, widely used in watches, the following reaction takes place

$$Zn(s) + Ag_2O(s) + H_2O(l)$$
 $Zn^{2+}(aq) + 2Ag(s) + 2OH^{-}(aq)$

Determine E^o and G^o for the reaction.

(Given:
$$E_{Ag/Ag}^{o} = + 0.80 \text{ V}, E_{Zn^2/Zn}^{o} = - 0.76 \text{ V}$$
)

OR

- (a) Define molar conductivity of a solution and explain how molar conductivity changes with change in concentration of solution for a weak and a strong electrolyte.
- (b) The resistance of a conductivity cell containing 0.001 M KCl solution at 298 K is 1500 . What is the cell constant if the conductivity of 0.001 M KCl solution at 298 K is 0.146×10^{-3} S cm⁻¹?

- **29.** (a) Complete the following chemical reaction equations:
 - $(i) P_4 + SO_2Cl_2$
 - (ii) XeF₆ + H₂O
 - (b) Predict the shape and the asked angle (90° or more or less) in each of the following cases:
 - (i) SO_3^{2-} and the angle O—S—O
 - (ii) ClF₃ and the angle F—Cl—F
 - (iii) XeF₂ and the angle F—Xe—F

OR

- (a) Complete the following chemical equations:
 - (i) NaOH + Cl₂ (hot and conc.)
 - (ii) $XeF_4 + O_2F_2$
- (b) Draw the structures of the following molecules:
 - (i) H₃PO₂
 - (ii) H₂S₂O₇
 - (iii) XeOF₄
- **30.** (a) Illustrate the following name reactions giving suitable example in each case:
 - (i) Clemmensen reduction
 - (ii) Hell-Volhard-Zelinsky reaction
 - (b) How are the following conversions carried out?
 - (i) Ethylcyanide to ethanoic acid
 - (ii) Butan-1-ol to butanoic acid
 - (iii) Benzoic acid to m-bromobenzoic acid

OR

- (a) Illustrate the following reactions giving a suitable example for each:
 - (i) Cross aldol condensation
 - (ii) Decarboxylation
- (b) Give simple tests to distinguish between the following pairs of compounds
 - (i) Pentan-2-one and Pentan-3-one
 - (ii) Benzaldehyde and Acetophenone
 - (iii) Phenol and Benzoic acid

CBSE (Delhi) SET-II

Questions Uncommon to Set-I

- 1. Write a point of distinction between a metallic solid and an ionic solid other than metallic lustre.
- 11. Describe a conspicuous change observed when
 - (i) a solution of NaCl is added to a sol of hydrated ferric oxide.
 - (ii) a beam of light is passed through a solution of NaCl and then through a sol.

- **13.** Describe the following:
 - (i) The role of cryolite in electrometallurgy of aluminium.
 - (ii) The role of carbon monoxide in the refining of crude nickel.
- **18.** Write the main structural difference between DNA and RNA. Of the two bases, thymine and uracil, which one is present in DNA?
- **20.** A solution of glycerol (C₃H₈O₃) in water was prepared by dissolving some glycerol in 500 g of water. This solution has a boiling point of 100.42 °C while pure water boils at 100 °C. What mass of glycerol was dissolved to make the solution?
 - $(K_b \text{ for water} = 0.512 \text{ K kg mol}^{-1})$
- 22. State a reason for each of the following situations:
 - (i) Co²⁺ is easily oxidized to Co³⁺ in the presence of a strong ligand.
 - (ii) CO is a stronger complexing reagent than NH₃.
 - (iii) The molecular shape of Ni(CO)₄ is not the same as that of $[Ni(CN)_4]^{2-}$.
- 23. How would you account for the following?
 - (i) With the same d-orbital configuration (d^4) Cr^{2+} is a reducing agent while Mn^{3+} is an oxidizing agent.
 - (ii) The actinoids exhibit a larger number of oxidation states than the corresponding members in the lanthanoid series.
 - (iii) Most of the transition metal ions exhibit characteristic colours in aqueous solutions.
- **30.** (a) Give a possible explanation for each one of the following:
 - (i) There are two —NH₂ groups in semicarbazide. However, only one such group is involved in the formation of semicarbazones.
 - (ii) Cyclohexanone forms cyanohydrin in good yield but 2, 4, 6-trimethylcyclohexanone does not.
 - (b) An organic compound with molecular formula C₉H₁₀O forms 2, 4-DNP derivative, reduces Tollen's reagent and undergoes Cannizzaro's reaction. On vigorous oxidation it gives 1,2-benzene-di-carboxylic acid. Identify the compound.

OR

- (a) Give chemical tests to distinguish between
 - (i) Phenol and Benzoic acid
 - (ii) Benzophenone and Acetophenone
- (b) Write the structures of the main products of following reactions:

(i)
$$+ C_6H_5COCl$$
 Anhydrous AICl₃ CS_2 (ii) $+ H_3CCC$ $+ CCC$ $+ CC$ $+ C$ $+ C$

CBSE (Delhi) SET-III

Questions Uncommon to Set-I and II

- **5.** Draw the structure of hex-1-en-3-ol compound.
- **20.** 15.0 g of an unknown molecular material was dissolved in 450 g of water. The resulting solution was found to freeze at -0.34 °C. What is the molar mass of this material? (K_f for water = 1.86 K kg mol⁻¹)
- 22. Explain the following observations giving an appropriate reason for each.
 - (i) The enthalpies of atomization of transition elements are quite high.
 - (ii) There occurs much more frequent metal-metal bonding in compounds of heavy transition metals (i.e., 3rd series).
 - (iii) Mn²⁺ is much more resistant than Fe²⁺ towards oxidation.
- 23. Write the name, the structure and the magnetic behaviour of each one of the following complexes:
 - (i) $[Pt(NH_3)Cl(NO_2)]$
 - (ii) [Co(NH₃)₄Cl₂]Cl
 - (iii) Ni(CO)₄
 - (At. nos. Co = 27, Ni = 28, Pt = 78)
- **27.** Explain the following terms giving one example of each type:
 - (i) Antacids
 - (ii) Disinfectants
 - (iii) Enzymes
- **30.** (a) Draw the molecular structures of following compounds:
 - (i) XeF₆
 - (ii) H₂S₂O₈
 - (b) Explain the following observations:
 - (i) The molecules NH₃ and NF₃ have dipole moments which are of opposite directions.
 - (ii) All the bonds in PCl₅ molecule are not equivalent.
 - (iii) Sulphur in vapour state exhibits paramagnetism.

OF

- (a) Complete the following chemical equations:
 - (i) $XeF_4 + SbF_5$
 - (ii) $Cl_2 + F_2(excess)$
- (b) Explain each of the following:
 - (i) Nitrogen is much less reactive than phosphorus.
 - (ii) The stability of +5 oxidation state decreases down group 15.
 - (iii) The bond angles (O—N—O) are not of the same value in NO₂⁻ and NO₂⁺.



CBSE (Delhi) SET-I

- 1. The process of addition of an appropriate amount of suitable impurity to an intrinsic semiconductor to enhance its conductivity is called doping.
- 2. Graphite is used as electrodes in the electrometallurgy of aluminium. The electrode reactions are:

- 3. PCl_4^- , as phosphorus has 10 electrons which can't be accommodated in sp^3 hybrid orbitals.
- **4.** 3-Bromo-2-methylpropene

- **6.** Butanone < Propanone < Propanal < Ethanal
- 7. $(CH_3)_2NH > CH_3NH_2 > (CH_3)_3N > NH_3$
- **8.** Homopolymerisation is a polymerisation reaction in which monomers of one kind are allowed to polymerise and form a homopolymer.

$$n \text{ CH}_2$$
—CH $_2$ Polymerization —CH $_2$ —CH $_2$ — $_n$ Ethene Polyethene (Homopolymer)

9.
$$T_b = 100.18$$
°C $- 100$ °C $= 0.18$ °C or 0.18 K; $K_b = 0.512$ K kg mol⁻¹; $m = 1$ mol kg⁻¹
$$T_b = i \ K_b \ m$$
$$0.18 \ \text{K} = i \times 0.512 \ \text{K kg mol}^{-1} \times 1 \ \text{mol kg}^{-1}$$
$$i = \frac{0.18}{0.512} = 0.35$$

OR

- (i) Mole fraction of a component in a solution may be defined as the ratio of moles of that component to the total number of moles of all the components present in the solution.
- (ii) The solutions of the same osmotic pressure at a given temperature are called isotonic solutions.
- (iii) van't Hoff's factor (i) may be defined as the ratio of normal molecular mass to observed molecular mass or the ratio of observed colligative property to normal colligative property.
- (iv) A solution which obeys Raoult's law over the entire range of concentration is called an ideal solution.
- **10.** The sum of powers of the concentration of the reactants in the rate law expression is called order of reaction.

For a general reaction: aA + bB Products

If rate = $k[A]^m [B]^n$; order of reaction = m + n

(i) General unit of rate constant, $k = (\text{mol } L^{-1})^{1-n} \text{ s}^{-1}$

$$L^{-1} \text{ mol } s^{-1} = (\text{mol } L^{-1})^{1-n} s^{-1}$$

$$-1 = -1 + n$$
 $n = 0$ or $1 = 1 - n$ $n = 0$ Reaction order $= 0$ L $\text{mol}^{-1} \text{ s}^{-1} = (\text{mol } \text{L}^{-1})^{1-n} \text{ s}^{-1}$

(ii)
$$L \text{ mol}^{-1} \text{ s}^{-1} = (\text{mol } L^{-1})^{1-n} \text{ s}^{-1}$$

$$1 = -1 + n$$
 $n = 2$ or $-1 = 1 - n$ $n = 2$ Reaction order $= 2$

- 11. The two groups into which phenomenon of catalysis can be divided are:
 - (i) Homogeneous catalysis: When the reactants and the catalyst are in the same phase, the catalysis is said to be homogeneous catalysis. For example, SO2 is oxidized to SO3 in the presence of nitric oxide, NO as catalyst.

$$2SO_2(g) + O_2(g)$$
 $^{NO(g)}$ $2SO_3(g)$

(ii) Heterogeneous catalysis: When the reactants are in a different phase than the catalyst, the catalysis is said to be heterogeneous. For example, the combination of dinitrogen and dihydrogen to form ammonia in the presence of finally divided iron as a catalyst.

$$N_2(g) + 3H_2(g)$$
 Fe(s) $2NH_3(g)$

- 12. The process of settling of colloidal particles through induced aggregation by the addition of some suitable electrolyte is known as coagulation. Three methods by which coagulation of lyophobic sols can be carried out are:
 - (i) Electrophoresis: During electrophoresis the colloidal particles move towards oppositely charged electrodes, get discharged and coagulated.
 - (ii) Boiling: On boiling a sol, the adsorbed layer is disturbed due to increased collision with the molecules of dispersion medium. This reduces the charge on the particles which ultimately settle down in the form of a precipitate.
 - (iii) Addition of Electrolytes: When excess of an electrolyte is added to a colloidal solution, the colloids interact with ions carrying charge opposite to that present on themselves. This causes neutralisation leading to their coagulation.
- 13. (i) In Mond process, nickel is converted into its volatile complex, teracarbonyl nickel and gets collected elsewhere. It is then decomposed to give pure nickel.

- (ii) Column chromatography is based on the principle that different components of a mixture are differently adsorbed on an adsorbent. In purification of rare earth elements ion-exchange is used as an adsorbent.
- **14.** (i) It is due to the ability of oxygen to form multiple bonds with metals.
 - (ii) This is because the energy required for the promotion of electrons in xenon is very high.

15. (i)
$$Cr_2O_7^{2^-} + 14H^+ + 6e^ 2Cr^{3^+} + 7H_2O$$

$$2I^- I_2 + 2e^-] \times 3$$

$$Cr_2O_7^{2^-} + 6I^- + 14H^+ 2Cr^{3^+} + 3I_2 + 7H_2O$$

(ii)
$$MnO_4^- + 8H^+ + 5e^ Mn^{2+} + 4H_2O] \times 2$$

 $NO_2^- + H_2O$ $NO_3^- + 2H^+ + 2e^-] \times 5$
 $2MnO_4^- + 5NO_2^- + 6H^+$ $2Mn^{2+} + 5NO_3^- + 3H_2O$

- **16.** (*i*) Peptide linkage is an amide (—CONH₂) linkage formed between —COOH group of one -amino acid and —NH₂ group of other -amino acid by the loss of a water molecule in a peptide.
 - (ii) A number of reactions that occur in the bodies of animals and plants to maintain the life process are catalysed by enzymes. The enzymes are thus termed as biocatalysts. Almost all the enzymes are globular proteins. An enzyme catalyses a biochemical reaction by providing alternate lower activation path thereby increasing the rate of the biochemical reaction. For example, activation energy for acid hydrolysis of sucrose is 6.22 kJ mol⁻¹ while the activation energy is only 2.15 kJ mol⁻¹ when hydrolyzed in the presence of enzyme sucrose.
- **17.** (*i*) The pentaacetate of glucose does not react with hydroxylamine indicating the absence of free —CHO group.
 - (ii) D—(+)—glucose when heated with methly alcohol in the presence of dry HCl gas forms a mixture containing methyl -D-glucoside and methyl -D-glucoside.

18.
$$\begin{array}{c} H_2C \xrightarrow{N} C = O \\ (i) \ H_2C \xrightarrow{CH_2} CH_2 \\ H_2C - CH_2 \end{array}$$

$$\begin{array}{c} (ii) \ CH_3 - CH = CH_2 \\ Propene \end{array}$$

19. a = 316.5 pm

For *bcc* unit cell,
$$r = \frac{\sqrt{3}}{4}a$$

$$r = \frac{\sqrt{3}}{4} \times 316.5 \text{ pm} = 137.04 \text{ pm}$$

OR

 $a = 286.65 \text{ pm} = 286.65 \times 10^{-10} \text{ cm}; M = 55.845 \text{ g mol}^{-1}; d = 7.874 \text{ g cm}^{-3}$

For bcc unit cell, z = 2

Substituting the values in the expression, $N_A = \frac{z}{a^3} \frac{M}{d}$, we get

$$N_A = \frac{2 \cdot 55.845 \text{ g mol}^{-1}}{(286.65 \cdot 10^{-10} \text{ cm})^3 \cdot 7.874 \text{ g cm}^{-3}}$$
$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$

20. i = 2, $W_A = 1$ kg = 1000 g; $T_f = 2$ K; $K_f = 1.86$ K kg mol⁻¹; $M_B = 74.5$ g mol⁻¹ Substituting these values in the expression,

$$W_B = \frac{T_f \quad M_B \quad W_A}{i \quad K_f \quad 1000}$$

$$W_B = \frac{2 \text{ K} \quad 74.5 \text{ g mol}^{-1} \quad 1000 \text{ g}}{2 \quad 1.86 \text{ K kg mol}^{-1} \quad 1000 \text{ g kg}^{-1}} = 40.05 \text{ g}$$

21. Suppose order w.r.t. NO is m and order w.r.t. Cl_2 is n. Then the rate will be

Rate =
$$k[NO]^m[Cl_2]^n$$

Substituting the values of experiment 1 to 3 in the rate expression, we get

$$0.60 = k(0.15)^m (0.15)^n \qquad \dots (i)$$

$$1.20 = k(0.15)^m (0.30)^n \qquad ...(ii)$$

$$2.40 = k(0.30)^{m} (0.15)^{n} \qquad ...(iii)$$

Dividing equation (iii) by (i), we get

$$\frac{2.40}{0.60} = \frac{k(0.30)^m (0.15)^n}{k(0.15)^m (0.15)^n}$$

$$4 = 2^m \quad \text{or} \quad 2^2 = 2^m \quad \text{or} \quad m = 2$$

Dividing equation (ii) by (i), we get

$$\frac{1.20}{0.60} = \frac{k(0.15)^m (0.30)^n}{k(0.15)^m (0.15)^n}$$

$$2 = 2^n$$
 or $n = 1$

- $2 = 2^n$ or n = 1(a) Rate law expression is, Rate = $k[NO]^2$ [Cl₂]
- (b) $0.60 \text{ mol } L^{-1} \text{ min}^{-1} = k(0.15 \text{ mol } L^{-1})^2 (0.15 \text{ mol } L^{-1})$

$$k = 177.77 \text{ mol}^{-2} \text{ L}^2 \text{ min}^{-1}$$

- (i) Interstitial compounds are well known for many of the transition elements because the 22. transition elements are capable of entrapping small sized atoms such as H, C and N in the interstitial sites in their crystal lattices. These trapped atoms get bonded to the atoms of transition elements. TiC, Fe₃H, Mn₄N, etc. are some interstitial compounds
 - (ii) This is due to filling of 4f orbitals which have poor shielding effect.
 - (iii) This is due to very small energy gaps between 5f, 6d and 7s subshells in actinoids.
- 23. (i) [Co(NH₃)Cl(en)₂]²⁺: Amminechloridobis (ethane-1, 2-diamine)-cobalt(III) ion Magnetic behaviour : Diamagnetic
 - (ii) $[Ni(H_2O)_2(OX)_2]^{2-}$: Diaquadioxalatonickelate (II) ion Magnetic behaviour: Paramagnetic
- 24. Chlorine withdraws electrons through negative inductive effect and releases electron through resonance. Through negative inductive effect, chlorine destabilizes the intermediate carbocation formed during the electrophilic substitution. Through resonance effect, chlorine tends to stabilize the carbocation and the effect is more pronounced at ortho and para-positions

Benzyl alcohol

Resonance effect stabilizes the intermediate carbocation

The negative inductive effect is stronger than resonance effect and causes net electron withdrawal and thus causes net deactivation. The resonance effect tends to oppose the inductive effect for the attack at ortho and para positions and hence makes the deactivation less for ortho and para attack. Thus the electrophilic substitution takes place at ortho and para positions..

Alternatively,

NH₂

$$\begin{array}{c}
NaNO_2/HCl \\
\hline
NaNO_2/HCl \\
\hline
273-278 \text{ K}
\end{array}$$

$$\begin{array}{c}
H_3PO_2/H_2O \\
\hline
CH_2OH
\end{array}$$

$$\begin{array}{c}
CHCl_3/\\
\hline
anhyd. AlCl_3
\end{array}$$

$$\begin{array}{c}
CH_2/h \\
\hline
CH_2-Cl
\end{array}$$

$$\begin{array}{c}
NaOH(aq) \\
\hline
Benzyl alchol
\end{array}$$
Benzyl alchol

- **27.** (*i*) Tranquilizers are a class of chemical compounds used for the treatment of stress, fatigue, and mild or even severe mental diseases. These relieve anxiety, stress, irritability or excitement by inducing a sense of well-being, *e.g.*, iproniazid, chlordiazopoxide, equanil, luminal, etc.
 - (ii) Food preservatives are the chemical substances which are added to food materials to prevent their spoilage due to microbial growth and to retain their nutritive value for long periods. Preservatives prevent the rancidity of food and inhibit growth or kill the microorganisms. The most-common preservatives used are sugar, vegetable oil, sodium benzoate, salts of ascorbic acid and propanoic acid.
 - (iii) Synthetic detergents are cleansing agents, which have all the properties of soaps but actually do not contain any soap. These can be used in both soft water and hard water as they produce foam even in hard water. These are mainly classified into three categories:
 - Anionic detergents, e.g., sodium dodecylbenzene sulphonate
 - Cationic detergents, e.g., cetyltrimethyl ammonium bromide
 - Non-ionic detergents, e.g., polyethylene glycol stearate

 $G^{\circ} = -301.08 \text{ kJ mol}^{-1}$

28. (a) Lead storage battery is a secondary battery. The reactions occurring in lead storage battery when current is drawn from it are:

Anode:
$$Pb(s) + SO_4^{2-}(aq)$$
 $PbSO_4(s) + 2e^ PbSO_4(s) + 2e^ PbSO_4(s) + SO_4^{2-}(aq) + 4H^+(aq) + 2e^ PbSO_4(s) + 2H_2O(l)$ $Overall \ reaction:$ $Pb(s) + PbO_2(s) + 2H_2SO_4(aq)$ $2PbSO_4(s) + 2H_2O(l)$ $Zn = Zn^{2+} + 2e^ Zn^{2+} + 2e^ Zn^{2+} + 2e^ Zn^{2+} + 2e^ Zn^{2+} + 2ag + 2OH^ Zn^{2+} + 2ag + 2OH^ Zn^{2+} + 2ag + 2OH^-$; $Zn^{2+} + 2ag + 2OH^-$; Zn^{2

(a) Molar conductivity of a solution at a given concentration is the conductance of the volume V of the solution containing one mole of electrolyte kept between two electrodes with area of cross section A and distance of unit length.


$$V = \frac{1000}{c}$$

where $\,$ is the conductivity and V is the volume of solution containing one mole of the electrolyte and c is the molar concentration.

Molar conductivity increases with decrease in concentration or increase in dilution as the number of ions as well as the mobility of ions increases with increase in dilution.

For strong electrolytes, the number of ions does not increase appreciably on dilution and only mobility of ions increases due to decrease in interionic attractions. Therefore,

 $_m$ increases a little as shown in figure by a straight line. For weak electrolytes, number of ions as well as mobility of ions increases on dilution which results in a very large increase in molar conductivity, especially at infinite dilution (*i.e.*, concentration, c 0) as shown by curve in figure.



(b) R = 1500 ; $= 0.146 \times 10^{-3} \text{ S cm}^{-1}$


Substituting the values in the expression,


Cell constant = $\times R$

Cell constant = $0.146 \times 10^{-3} \text{ S cm}^{-1} \times 1500$

$$= 0.219 \text{ cm}^{-1}$$

- **29.** (a) (i) $P_4 + 10SO_2Cl_2$ 4PCl₅ + $10SO_2$
 - (ii) XeF₆ + 3H₂O XeO₃ + 6HF
 - (b) (i) There are three bond pairs and one lone pair of electrons around S atoms in SO_3^{2-} . Therefore according to VSEPR theory, SO_3^{2-} should be pyramidal. The angle O—S—O is greater than 90° .






CIF₃

(ii) There are three bond pairs and two lone pairs of electrons around Cl atom in ClF₃. Therefore according to VSEPR theory, ClF₃ should be bent T-shaped. The angle F—Cl—F is less than 90°.

(iii) There are two bond pairs and three lone pairs of electrons around Xe atoms in XeF₂. Therefore according to VSEPR theory, XeF₂ should be linear. The angle F—Xe—F is greater than 90°.



OR


(a) (i)
$$6$$
NaOH + 3 Cl₂ 5 NaCl + NaClO₃ + 3 H₂O (hot and conc)

XeF₂

(ii)
$$XeF_4 + O_2F_2$$

 $XeF_6 + O_2$

H₃PO₂



XeOF₄

(i) The carbonyl group of aldehydes and ketones is reduced to CH₂ group on treatment with zinc-amalgam and concentrated hydrochloric acid. The reaction is known as Clemmensen reduction

$$C=O$$
 $\xrightarrow{Zn-Hg}$ $>CH_2 + H_2O$

$$CH_3COCH_3 + 4(H)$$
 $\xrightarrow{Zn-Hg}$ $CH_3CH_2CH_3 + H_2O$
Acetone Propane

(ii) Carboxylic acids having an -hydrogen are halogenated at the -position on treatment with chlorine or bromine in the presence of small amount of red phosphorus to give -halocarboxylic acids. This reaction is called Hell-Volhard-Zelinsky reaction.

(b) (i)
$$CH_3CH_2CN$$
 $\xrightarrow{H_3O^+}$ CH_3CH_2COOH $\xrightarrow{+NH_3}$ $CH_3CH_2CONH_2$ $\xrightarrow{Br_2/KOH}$ CH_3COOH $\xleftarrow{K_2Cr_2O_7/H_2SO_4}$ CH_3CH_2OH $\xleftarrow{HNO_2}$ $CH_3CH_2NH_2$ $CH_3CH_2NH_2$

(iii)
$$CH_3$$
— CH_2 — CH_2 — CH_2 — OH

Butan-1-ol

COOH

Butanoic acid

COOH

Br₂/FeBr₃

Benzoic acid

 m -Bromobenzoic acid

(a) (i) When aldol condensation is carried out between two different aldehydes and/or ketones, it is called cross aldol condensation.

(ii) Sodium salt of carboxylic acids on heating with soda lime lose carbon dioxide and form hydrocarbons. This reaction is called decarboxylation.

$$\begin{array}{ccc} R - COONa & {}^{\rm NaOH \ and \ CaO} & R - H & {}^{\rm Na}_2{\rm CO}_3 \\ {}^{\rm Sodium \ carboxylate} & {}^{\rm Na}_2{\rm CO}_3 \end{array}$$

(b) (i) Pentan-2-one when treated with NaOI (I₂/NaOH) gives yellow precipitate of iodoform but pentan-3-one does not give this test.

(ii) Benzaldehyde being an aldehyde gives silver mirror with Tollen's reagent but acetophenone being a ketone does not give this test.

$$C_6H_6CHO + 2[Ag(NH_3)_2]^+ + 3OH^ C_6H_5COO^- + 2Ag + 4NH_3 + 2H_2O$$

Benzaldehyde Tollen's reagent Silver mirror

(iii) Benzoic acid produces brisk effervescence with sodium bicarbonate solution while phenol does not.

$$C_6H_5COOH + NaHCO_3$$
 $C_6H_5COONa + CO_2 + H_2O$ Benzoic acid Sodium bicarbonate

CBSE (Delhi) SET-II

- 1. Metallic solids conduct electricity in solid state while ionic solids are insulators in solid state.
- 11. (i) The positively charged colloidal particles of Fe(OH)₃ get coagulated by negatively charged Cl⁻ ions provided by electrolyte sodium chloride.
 - (ii) The path of light is not visible when light is passed through NaCl solution but visible when light is passed through a sol due to scattering of light by colloidal particles.
- **13.** (*i*) The role of electrolyte is two-fold:
 - It covers the melting point of the mixture to about 1250 K.
 - It increases the electrical conductivity of the mixture.
 - (ii) When nickel is heated with carbon monoxide it forms a volatile compound tetracarbonyl nickel, Ni(CO)₄ which on further heating at higher temperature decomposes to give pure nickel.

Ni + 4CO
$$^{330-350K}$$
 Ni(CO) $_4$; Ni(CO) $_4$ $^{450-470K}$ Ni + 4CO Impure nickel

18.

Structural difference

DNA	RNA	
1. The sugar present in DNA is 2-deoxy-D-(-)-ribose.	1. The sugar present in RNA is D-(–)-ribose.	
2. DNA has double stranded -helix structure.	2. RNA has single stranded -helix structure.	
3. DNA contains cytosine and thymine as pyrimidine bases.	3. RNA contains cytosine and uracil as pyrimidine bases.	
4. DNA molecules are very large and their molecular mass may vary from $6 \times 10^6 - 16 \times 10^6$ u.	4. RNA molecules are much smaller and their molecular may vary from 20,000–40,000 u.	

Thymine is present in DNA.


20.
$$T_b = 100.42$$
°C $- 100.00$ °C $= 0.42$ °C or 0.42 K; $W_A = 500$ g $K_b = 0.52$ K kg mol⁻¹; $M_B = 92$ g mol⁻¹

Substituting these values in the expression,

$$W_B = \frac{T_b \quad M_B \quad W_A}{K_b \quad 1000}, \text{ we get}$$

$$W_B = \frac{0.42 \text{ K} \quad 92 \text{ g mol}^{-1} \quad 500 \text{ g}}{0.512 \text{ K kg mol}^{-1} \quad 1000 \text{ g kg}^{-1}} = 37.73 \text{ g}$$

- **22.** (*i*) Because strong ligand causes spin pairing giving rise to diamagnetic octahedral complex which are very stable and have very large crystal field stabilization energy. This splitting energy overcomes the ionization enthalpy.
 - (ii) CO is capable of accepting an appreciable amount of electron density from a filled metal d-orbital into vacant antibonding *-orbital of its own. This type of interaction increases the value of crystal field stabilization energy. This explain why CO is a stronger complexing reagent than NH₃.
 - (iii) Ni in $[Ni(CO)_4]$ is sp^3 hybridized. Hence, it is tetrahedral. In $[Ni(CN)_4]^{2-}$, the Ni^{2+} is dsp^2 hybridized. Hence, it has square planar geometry.



- **23.** (*i*) Cr^{2+} is a reducing agent as its configuration changes from d^4 to d^3 , the later having a half-filled t_{2g} level. On the other hand, the change from Mn^{3+} to Mn^{2+} results in the extra stable half-filled (d^5) configuration and hence Mn^{3+} is an oxidizing agent.
 - (ii) This is due to very small energy gaps between 5f, 6d and 7s subshells in actinoids.
 - (iii) In aqueous solutions, the transition metal ions which have partially filled *d*-orbitals undergo *d*-*d* transition by absorbing light from visible region and radiate complementary colour.

$$\begin{array}{c|c} \vdots \ddot{O} \\ \vdots \ddot{O} \\ \vdots \\ H_2 \ddot{N} - C - \ddot{N}H - \ddot{N}H_2 \longrightarrow H_2 \ddot{N} - C = \ddot{N}H - \ddot{N}H_2 \\ \vdots \ddot{O} \vdots \\ \vdots \ddot{O} \vdots \\ \vdots \ddot{O} \vdots \ddot{O} \vdots \\ \ddot{O} \vdots \\ \vdots \ddot{O} \vdots \\ \ddot{O} \vdots \\ \vdots \ddot{O} \vdots \\ \ddot{O}$$

Although semicarbazide has two — NH_2 group but one which is directly attached to C=O is involved in the resonance as shown above. Consequently, electron density on this — NH_2 group decreases and hence it does not act as a nucleophile. On the other hand, the lone pair of electrons on the other — NH_2 group is not involved in resonance and hence is available for nucleophilic attack on the C=O group of aldehydes and ketones.

(ii) Due to presence of three methyl groups, the nucleophilic attack by CN⁻ ion does not occur due to steric hindrance in 2, 4, 6-trimethyl cylcohexanone. As there is no such steric hindrance in cyclohexanone so nucleophilic attack by the CN⁻ ion occurs readily and hence cyclohexanone cyanohydrin is obtained in good yield.

The reactions involved in the question are:

$$CHO + 2[Ag(NH_3)_2]^+ + 3OH$$
 $COO^- + 2Ag^- + 2H_2O + 4NH_3$ CH_2CH_3

Tollen's reagent Silver mirror

2 CHO
$$OH$$
 CH₂CH₃ CH₂CH₃ CH₂CH₃

$$\begin{array}{c|c} CHO & \xrightarrow{KMnO_4/KOH} & \\ \hline COO^- & \xrightarrow{H_3O^+} & \\ \hline COOH & \\ \hline \\ COOH & \\ \hline \end{array}$$

2-Ethyl benzaldehyde

1, 2-Benzenedicarboxylic acid

OR

(a) (i) Phenol gives a violet colouration with neutral FeCl₃ solution while benzoic acid gives buff coloured precipitate of ferric benzoate.

$$\begin{aligned} & 6C_6H_5OH \ + \ FeCl_3 & & [Fe(OC_6H_5)_6]^{3-} \ + \ 3H^+ \ + \ 3HCl \\ & & Violet\ complex \\ & 3C_6H_5COOH \ + \ FeCl_3 & & (C_6H_5COO)_3Fe \ + \ 3HCl \\ & Benzoic\ acid & & Buff\ coloured\ ppt. \end{aligned}$$

(ii) Acetophenone being a methyl ketone when treated with NaOI(I₂/NaOH) gives yellow precipitate of iodoform while benzophenone does not give this test.

Bezophenone p-Nitro benzaldehyde

CBSE (Delhi) SET-III

5.
$$CH_3$$
— CH_2 — CH_2 — CH — CH — CH_2
OH


20.
$$f = 0$$
°C $- (-0.34$ °C) $= 0.34$ °C or 0.34 K
 $W_B = 15.0$ g; $W_A = 450$ g; $K_f = 1.86$ K kg mol⁻¹

Substituting these values in the expression,

$$M_B = \frac{K_f \quad W_B \quad 1000}{T_f \quad W_A}$$
, we get


$$M_B = \frac{1.86 \text{ K kg mol}^{-1} \quad 15.0 \text{ g} \quad 1000 \text{ g kg}^{-1}}{0.34 \text{ K} \quad 450 \text{ g}} = 182.35 \text{ g mol}^{-1}$$

- 22. (i) Because of large number of unpaired electrons in their atoms, the transition metals have strong interatomic interactions and hence stronger bonding between atoms, resulting in higher enthalpies of atomization.
 - (ii) In the same group of d-block elements, the 4d and 5d transition elements are larger in size than than those of 3d elements. Hence, the valence electrons are less tightly held and form metal-metal bond more frequently.
 - (iii) $Mn^{3+}(d^4)$ is less stable than Mn^{2+} (d^5 , half filled) while Fe^{3+} (d^5 , half filled) is more stable than Fe²⁺ (d^4). That is why Mn²⁺ is more resistance than Fe²⁺ towards oxidation.
- (i) $[Pt(NH_3)_2CINO_2]$: Diamminechloridonitrito-N-platinum (II); $Pt^{2+} = 5d^8$




Structure = Square planar; Magnetic behaviour : Diamagnetic

(ii) $[CO(NH_3)_4Cl_2]Cl$: Tetraamminedichloridocobalt (III) chloride; $Co^{3+} = 3d^6$



Structure = Octahedral; Magnetic behaviour : Diamagnetic


(iii) $Ni(CO)_4$: Tetracrabonyl nickel(o); $Ni = 3d^8 4s^2$




Structure = Tetrahedral; Magnetic behaviour : Diamagnetic

- **27.** (*i*) **Antacids:** Chemical substances which remove excess acid in the stomach and raise the pH to appropriate level, *e.g.*, sodium hydrogencarbonate, a mixture of aluminium and magnesium hydroxide, ranitidine, etc.
 - (ii) **Disinfectants:** These are the chemical substances which kill microorganisms or stop their growth but are harmful to human tissues, e.g., phenol (1%), chlorine in concentration of 0.2 to 0.4 pm in aqueous solution, SO_2 , etc.
 - (iii) Enzymes: Enzymes are globular proteins with high molecular mass ranging from 15,000 to 1,000,000 g mol⁻¹, which form colloidal solution in water. A number of reactions that occur in the body of animals and plants to maintain the life processes are catalyzed by enzymes, therefore enzymes are termed as biochemical catalysts.

30.



- (b) (i) This is because in NH₃ nitrogen is more electronegative than hydrogen while in NF₃, nitrogen is less electronegative than fluorine.
 - (ii) In gaseous and liquid phases PCl₅ has a trigonal bipyramidal structure. The three equatorial P—Cl bonds are equivalent, while the two axial bonds are longer then equitorial bonds. This is due to the fact that the axial bond pairs suffer greater repulsion as compared to equatorial bond pairs.



(iii) In vapour state sulphur partly exists as S_2 molecule which has two unpaired electrons in the antibonding * orbitals like O_2 and hence it exhibits paramagnetism.

OR

- (a) (i) $XeF_4 + SbF_5 = [XeF_3]^+[SbF_6]^-$ (ii) $Cl_2 + 3F_2(excess) = {}^{573}K = 2ClF_3$
- (b) (i) As $N \equiv N$ triple bond (941.4 kJ mol⁻¹) is much stronger than P—P single bond (213 k J mol⁻¹), therefore nitrogen is much less reactive than phosphorus.
 - (ii) Due to inert pair effect stability of +5 oxidation decreases down the group 15.
 - (iii) In NO_2^- , nitrogen has a lone pair of electrons. As lone pair-bond pair repulsion is greater than bond pair-bond repulsion, thus O-N-O bond angle in NO_2^- is less than NO_2^+ .



CBSE EXAMINATION PAPERS

ALL INDIA-2012

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question numbers 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question numbers 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question numbers 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question numbers 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables, if necessary. Use of calculators is not allowed.

CBSE (All India) SET-I

- 1. How may the conductivity of an intrinsic semiconductor be increased?
- **2.** Define 'peptization'.
- 3. How is copper extracted from a low grade ore of it?
- **4.** Which is a stronger reducing agent, SbH₃ or BiH₃, and why?
- **5.** What happens when bromine attacks $CH_2 = CH CH_2 C = CH$?
- **6.** Write the IUPAC name of the following:

$$CH_3$$
— CH_2 — CH = CH — C — H

- 7. Write the structure of the product obtained when glucose is oxidised with nitric acid.
- **8.** Differentiate between disinfectants and antiseptics.
- **9.** Express the relation among cell constant, resistance of the solution in the cell and conductivity of the solution. How is molar conductivity of a solution related to its conductivity?

OR

The molar conductivity of a 1.5 M solution of an electrolyte is found to be 138.9 S cm² mol⁻¹. Calculate the conductivity of this solution.

- **10.** A reaction is of second order with respect to a reactant. How is its rate affected if the concentration of the reactant is (i) doubled (ii) reduced to half?
- 11. Which methods are usually employed for purifying the following metals
 - (i) Nickel
 - (ii) Germanium

Mention the principle behind each of them.

- **12.** Explain the following facts giving appropriate reason in each case:
 - (i) NF₃ is an exothermic compound whereas NCl₃ is not.
 - (ii) All the bonds in SF_4 are not equivalent.

- **13.** Complete the following chemical reaction equations:
 - (*i*) $Cr_2O_7^{2-}$ I^- H
 - (ii) MnO $_{4}^{-}$ NO $_{2}^{-}$ H
- 14. Explain the mechanism of acid catalysed hydration of an alkene to form corresponding alcohol.
- **15.** Explain the following behaviours:
 - (i) Alcohols are more soluble in water than the hydrocarbons of comparable molecular masses.
 - (ii) Ortho-nitrophenol is more acidic than ortho-methoxyphenol.
- **16.** Describe the following, giving the relevant chemical equation in each case:
 - (i) Carbylamine reaction
 - (ii) Hoffmann's bromamide reaction
- **17.** Complete the following reaction equations:
 - (i) $C_6H_5N_2Cl + H_3PO_2 + H_2O$
 - (ii) $C_6H_5NH_2 + Br_2(aq)$
- **18.** What are food preservatives? Name two such substances.
- **19.** Copper crystallises with face centred cubic unit cell. If the radius of copper atom is 127.8 pm, calculate the density of copper metal.

(Atomic mass of Cu = 63.55 u and Avogadro's number $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

OR

Iron has a body centred cubic unit cell with a cell dimension of 286.65 pm. The density of iron is 7.87 g cm^{-3} . Use this information to calculate Avogadro's number. (At. mass of Fe = 56.0 u)

- **20.** The electrical resistance of a column of 0.05 M NaOH solution of diameter 1 cm and length 50 cm is 5.55×10^3 ohm. Calculate its resistivity, conductivity and molar conductivity.
- 21. The reaction, $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ contributes to air pollution whenever a fuel is burnt in air at a high temperature. At 1500 K, equilibrium constant K for it is 1.0×10^{-5} . Suppose in a case $[N_2]$ = 0.80 mol L^{-1} and $[O_2] = 0.20$ mol L^{-1} before any reaction occurs. Calculate the equilibrium concentrations of the reactants and the product after the mixture has been heated to 1500 K.
- **22.** Explain the following terms giving a suitable example for each:
 - (i) Aerosol

(ii) Emulsion

- (iii) Micelle
- 23. How would you account for the following:
 - (i) Among lanthanoids, Ln (III) compounds are predominant. However, occasionally in solutions or in solid compounds, +2 and +4 ions are also obtained.
 - (ii) The $\rm E_{M^2/M}^o$ for copper is positive (0.34 V). Copper is the only metal in the first series of transition elements showing this behaviour.
 - (iii) The metallic radii of the third (5d) series of transition metals are nearly the same as those of the corresponding members of the second series.
- 24. Name the following coordination entities and draw the structures of their stereoisomers:
 - (i) $[Co(en)_2 Cl_2]^+$ (en = ethan-1, 2-diamine)
 - (ii) $[Cr(C_2O_4)_3]^{3-}$


(iii) [Co(NH₃)₃ Cl₃]

(Atomic numbers: Cr = 24, Co = 27)

- **25.** Answer the following questions:
 - (i) What is meant by chirality of a compound? Give an example.
 - (ii) Which one of the following compounds is more easily hydrolysed by KOH and why?

CH₃CHClCH₂CH₃ or CH₃CH₂CH₂Cl

(iii) Which one undergoes S_N2 substitution reaction faster and why?



- **26.** What is essentially the difference between -glucose and -glucose? What is meant by pyranose structure of glucose?
- 27. Differentiate between thermoplastic and thermosetting polymers. Give one example of each.
- **28.** (a) Define the following terms:
 - (i) Mole fraction

- (ii) Ideal solution
- (b) 15.0 g of an unknown molecular material is dissolved in 450 g of water. The resulting solution freezes at -0.34°C. What is the molar mass of the material? (K_f for water = 1.86 K kg mol⁻¹)

OR

- (a) Explain the following:
 - (i) Henry's law about dissolution of a gas in a liquid.
 - (ii) Boiling point elevation constant for a solvent.
- (b) A solution of glycerol ($C_3H_8O_3$) in water was prepared by dissolving some glycerol in 500 g of water. This solution has a boiling point of 100.42°C. What mass of glycerol was dissolved to make this solution? (K_b for water = 0.512 K kg mol⁻¹)
- 29. (a) Draw the molecular structures of the following compounds
 - (i) N₂O₅

- (ii) XeOF₄
- (b) Explain the following observations:
 - (i) Sulphur has a greater tendency for catenation than oxygen.
 - (ii) ICl is more reactive than I₂.
 - (iii) Despite lower value of its electron gain enthalpy with negative sign, fluorine (F_2) is a stronger oxidising agent than Cl_2 .

OR

- (a) Complete the following chemical equations:
 - (i) $Cu + HNO_3$ (dilute)
 - (ii) $XeF_4 + O_2F_2$
- (b) Explain the following observations:
 - (i) Phosphorus has a greater tendency for catenation than nitrogen.
 - (ii) Oxygen is a gas but sulphur a solid.
 - (iii) The halogens are coloured.
- **30.** (a) Write a suitable chemical equation to complete each of the following transformations:
 - (i) Butan-1-ol to butanoic acid
 - (ii) 4-Methylacetophenone to benzene-1, 4-dicarboxlylic acid.

(b) An organic compound with molecular formula C₉H₁₀O forms 2,4-DNP derivative, reduces Tollen's reagent and undergoes Cannizzaro's reaction. On vigorous oxidation it gives 1,2-benzenedicarboxylic acid. Identify the compound.

OR

- (a) Give chemical tests to distinguish between
 - (i) Propanol and propanone
 - (ii) Benzaldehyde and acetophenone
- (b) Arrange the following compounds in increasing order of their property as indicated:
 - (i) Acetaldehyde, Acetone, Methyl tert-butyl ketone (reactivity towards HCN)
 - (ii) Benzoic acid, 3,4-Dinitrobenzoic acid, 4-Methoxybenzoic acid (acid strength)
 - (iii) CH₃CH₂CH(Br)COOH, CH₃CH(Br)CH₂COOH, (CH₃)₂CHCOOH (acid strength)

CBSE (All India) SET-II

Questions Uncommon to Set-I

- 1. Which stoichiometric defect increases the density of a solid?
- 2. What is meant by 'shape selective catalysis'?
- 3. What is the role of collectors in Froth Floatation process?
- **6.** Write the IUPAC name of Ph—CH=CH—CHO.
- 17. Explain the cleaning action of soap. Why do soaps not work in hard water?
- **20.** A voltaic cell is set up at 25°C with the following half cells:

Write an equation for the reaction that occurs when the cell generates an electric current and determine the cell potential.

$$E_{\text{Ni}^{2+}/\text{Ni}}^{\text{o}} = -0.25 \text{ V}; E_{\text{Al}^{3+}/\text{Al}}^{\text{o}} = -1.66 \text{ V}. (\log 8 \times 10^{-6} = -0.54)$$

- **23.** Explain the following observations:
 - (i) Many of the transition elements are known to form interstitial compounds.
 - (ii) There is a general increase in density from titanium (Z = 22) to copper (Z = 29).
 - (iii) The members of the actinoid series exhibit a larger number of oxidation states than the corresponding members of the lanthanoid series.
- **27.** Explain the following terms giving a suitable example for each:
 - (i) Elastomers

(ii) Condensation polymers

- (iii) Addition polymers
- **30.** (a) Draw the structures of the following molecules:
 - (i) H₃PO₂

- (ii) ClF₃
- (b) Explain the following observations:
 - (i) Nitrogen is much less reactive than phosphorus.
 - (ii) Despite having greater polarity, hydrogen fluoride boils at a lower temperature than water.
 - (iii) Sulphur has a greater tendency for catenation than oxygen in the same group.

OR

- (a) Draw the structures of the following molecules:
 - (i) N₂O₅

- (ii) HClO₄
- (b) Explain the following observations:
 - (i) H₂S is more acidic than H₂O.
 - (ii) Fluorine does not exhibit any positive oxidation state.
 - (iii) Helium forms no real chemical compound.

CBSE (All India) SET-III

Questions Uncommon to Set-I and II

- **1.** What are *n*-type semiconductors?
- **4.** What is the basicity of H₃PO₂ acid and why?
- 5. Write the IUPAC name of the following:



- 7. Write a reaction which shows that all the carbon atoms in glucose are linked in a straight chain.
- **8.** What is the cause of a feeling of depression in human beings? Name a drug which can be useful in treating this depression.
- 11. Explain the role of each of the following:
 - (i) NaCN in the extraction of silver.
 - (ii) SiO_2 in the extraction of copper.
- **18.** Differentiate between disinfectants and antiseptics. Give one example of each group.
- 22. Write three distinct features of chemisorptions which are not found in physisorptions.
- **23.** Explain each of the following observations:
 - (i) With the same d-orbital configuration (d^4), Cr^{2+} is a reducing agent while Mn^{3+} is an oxidising agent.
 - (ii) Actinoids exhibit a much larger number of oxidation states than the lanthanoids.
 - (iii) There is hardly any increase in atomic size with increasing atomic numbers in a series of transition metals.
- 24. Name the following coordination entities and describe their structures:
 - (*i*) $[Fe(CN)_6]^{4-}$

(ii) $[Cr(NH_3)_4Cl_2]^+$

(iii) $[Ni(CN)_4]^{2-}$

(Atomic Numbers Fe =
$$26$$
, Cr = 24 , Ni = 28)

- **26.** Define the following as related to proteins:
 - (i) Peptide linkage
 - (ii) Primary structure
 - (iii) Denaturation

Solutions

CBSE (All India) SET-I

- 1. The conductivity is increased by adding an appropriate amount of impurity which is electron rich or election deficient as compared to intrinsic semiconductor.
- 2. The process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of small amount of electrolyte is called peptisation.
- **3.** Copper is extracted by hydrometallurgy from low grade copper ores. It is leached out using acid or bacteria. The solution containing Cu²⁺ ion is treated with scrap iron or H₂.

$$Cu^{2+}(aq) + Fe(s)$$
 $Fe^{2+}(aq) + Cu(s)$
OR
 $Cu^{2+}(aq) + H_2(g)$ $Cu(s) + 2H^+(aq)$

- 4. BiH₃, as BiH₃ has lower bond dissociation enthalpy than SbH₃.
- 5. The reddish brown colour of bromine is discharged.
- **6.** Pent-2-enal

7. CHO
$$(CHOH)_4$$
 $(CHOH)_4$ $(C$

8. Disinfectants are applied to non-living objects whereas antiseptics are applied to living tissues.

9. Conductivity () =
$$\frac{1}{\text{Resistance }(R)} \times \text{Cell constant}$$

Molar Conductivity (
$$_{m}$$
) = $\frac{\text{Conductivity () } 1000}{\text{Molarity (} M\text{)}}$

OR
$$_{m} = \frac{1000}{M} = \frac{\frac{m}{1000}}{1000}$$

$$= \frac{138.95 \text{ cm}^{2} \text{ mol}^{-1} \quad 1.5 \text{ mol L}^{-1}}{1000 \text{ cm}^{3} \text{ L}^{-1}} = 0.208355 \text{ cm}^{-1}$$

- **10.** Rate = $k[R]^2$
 - (i) If [R] is doubled, Rate = $k[2R]^2 = 4k[R]^2 = 4$ times

(ii) If [R] is reduced to
$$\frac{[R]}{2}$$
, Rate = $k \left(\frac{R}{2} \right)^2 = \frac{1}{4}k \left(R \right)^2 = \frac{1}{4}$ th

11. (i) Nickel is purified by vapour phase refining. In this method nickel is heated in a stream of carbon monoxide forming a volatile complex, nickel tetracarbonyl (Ni(CO)₄), which on further heating decomposes to give pure nickel.

- (ii) Germanium is purified by zone refining. This method is based on the principle that the impurities are more soluble in molten state than in the solid state of the metal.
- **12.** (*i*) This is due to lower bond dissociation enthalpy of fluorine than chlorine and small and strong bond formed by fluorine with nitrogen.
 - (ii) SF₄ has trigonal bipyramidal structure in which one of the equatorial positions is occupied by a lone pair of electrons. The two equatorial S–F bonds are equivalent, while the two axial S–F bonds are longer than equatorial bonds. This is because the axial bond pairs suffer more repulsion as compared to equatorial bond pairs.

13. (i)
$$Cr_2O_7^{2-} + 14H^+ + 6e^ 2Cr^{3+} + 7H_2O$$
 $2I^ I_2 + 2e^-] \times 3$ $Cr_2O_7^{2-} + 6I^- + 14H^+$ $2Cr^{3+} + 3I_2 + 7H_2O$ (ii) $MnO_4^- + 8H^+ + 5e^ Mn^{2+} + 4H_2O] \times 2$ $NO_2^- + H_2O$ $NO_3^- + 2H^+ + 2e^-] \times 5$ $2MnO_4^- + 5NO_2^- + 6H^+$ $2Mn^{2+} + 5NO_3^- + 3H_2O$

The mechanism of the reaction involves the following three steps:

(i) Protonation of alkene to form carbocation by electrophilic attack of H₃O⁺.

(ii) Nucleophilic attack of water on carbocation.

(iii) Deprotonation to form an alcohol.

- **15.** (i) This is due to their ability to form hydrogen bonds with water molecules.
 - (ii) Due to —R and —I effect of —NO₂ group, the electron density in the O—H bond decreases in ortho-nitrophenol and hence the release of H⁺ ion becomes easy. On the other hand, due to +R

effect of the —OCH₃ group, the electron density in the O—H bond in ortho-methoxyphenol increases, thereby making the release of H⁺ ion difficult.

(i) Carbylamine reaction: Aliphatic and aromatic primary amines when heated with chloroform and 16. ethanolic potassium hydroxide form carbylamines or isocyanides which are foul smelling substances. Secondary and tertiary amines do not show this reaction.

(ii) Hoffmann's bromamide reaction: When a primary acid amide is heated with bromine in an aqueous or ethanolic solution of NaOH, a primary amine is obtained. The amine so obtained contains one carbon less than that present in the amide.

diazonium chloride

- 18. Food preservatives are the chemical substances which are added to food materials to prevent their spoilage and to retain their nutritive value for long periods. Preservatives prevent the rancidity of food and inhibit the growth of or kill the microorganisms. Two such substances are sodium benzoate and salts of ascorbic acid.
- $a = 2\sqrt{2}r$ **19.** For *fcc*. $a = 2\sqrt{2} \times 127.8 \text{ pm} = 361.4 \text{ pm} = 361.4 \times 10^{-10} \text{ cm}$ z = 4; $M = 63.55 \text{ g mol}^{-1}$; $a = 3.614 \times 10^{-8} \text{ cm}$; $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$ Substituting the values in the expression,

$$d = \frac{z}{a^3} \frac{M}{N_A}, \text{ we get}$$

$$d = \frac{4 + 63.55 \text{ g mol}^{-1}}{(3.614 + 10^{-8} \text{ cm})^3 + 6.02 + 10^{23} \text{ mol}^{-1}} = 8.95 \text{ g cm}^{-3}$$
OR

Here,
$$z = 2$$
, M = 56 g mol⁻¹, $d = 7.87$ g cm⁻³
 $a = 286.65$ pm = 286 .65 × 10⁻¹⁰cm = 2.8

Substituting the values in the expression,

$$N_A = \frac{z \quad M}{a^3 \quad d}, \text{ we get}$$

$$N_A = \frac{2 \quad 56 \text{ g mol}^{-1}}{(286.65 \quad 10^{-10} \text{ cm})^3 \quad 7.87 \text{ g cm}^{-3}}$$

$$= 6.042 \times 10^{23} \text{ mol}^{-1}$$

$$= 6.042 \times 10^{23} \text{ mol}^{-1}$$

$$= 0.785 \text{ cm}^2; l = 50 \text{ cm}$$

$$\text{Resistivity} = \frac{R}{l} = \frac{A}{l} = \frac{5.55 \quad 10^3 \text{ ohm} \quad 0.785 \text{cm}^2}{50 \text{ cm}} = 87.135 \text{ ohm cm}$$

$$\text{Conductivity} = \frac{1}{87.135 \text{ ohm cm}} = 0.01148 \text{ s cm}^{-1}$$

$$\text{Molar Conductivity} = \frac{1000}{M} = \frac{0.01148 \text{ s cm}^{-1} \quad 1000 \text{ cm}^3 \text{ L}^{-1}}{0.05 \text{ mol L}^{-1}}$$

$$= 229. 6 \text{ s cm}^2 \text{ mol}^{-1}$$

$$\text{Initial conc. in mol L}^{-1} \qquad 0.8 \qquad 0.2 \qquad 0$$

-x +2xChange in conc. in mol L^{-1} -xEquilibrium conc. in mol L⁻¹ 0.8-x 0.2-x 2x

$$K_C = \frac{[\text{NO}]^2}{[\text{N}_2][\text{O}_2]}$$
 $1 \times 10^{-5} = \frac{(2x)^2}{(0.8 + x)(0.2 + x)}$

As x << 0.2, therefore 0.8 - x = 0.8 and 0.2 - x = 0.2

$$1 \times 10^{-5} = \frac{4x^2}{0.16}$$

$$4x^2 = 16 \times 10^{-7}$$

$$x = 6.324 \times 10^{-4} \text{ mol } L^{-1},$$


$$2x = 2 \times 6.324 \times 10^{-4} = 12.648 \times 10^{-4} \text{ mol } L^{-1}$$

Thus at equilibrium

$$[N_2] = 0.8 \text{ mol L}^{-1}$$
, $[O_2] = 0.2 \text{ mol L}^{-1}$ and $[NO] = 1.265 \times 10^{-3} \text{ mol L}^{-1}$


- 22. (i) A colloidal system in which dispersion medium is gas and dispersed phase is either solid or liquid is called aerosol, e.g., smoke, fog, etc.
 - (ii) An emulsion is a colloidal system in which both the dispersed phase and the dispersion medium are liquids, e.g., milk, cod liver oil, etc.
 - (iii) There are some substances such as soap which at low concentration behave as normal electrolytes, but at higher concentration exhibit colloidal behaviour due to the formation of aggregates. The aggregated particles thus formed are known as micelles or associated colloids.
- (i) Lanthanoid metals show +2 and +4 oxidation states to attain extra stable f^0 and f^7 configurations. 23.
 - (ii) The main reason for positive $E^{\circ}(0.34 \text{ V})$ value for copper is that the sum of enthalpies for sublimation and ionisation is not balanced by hydration enthalpy.

- (iii) This is due to lanthanoid contraction which arises due to filling of 4 f-orbitals which have poor shielding effect.
- 24. (i) [CO(en)₂Cl₂]⁺: Dichloride bis (ethane-1,2-diamine) cobalt (III) ion




Geometrical isomers

(ii) [Cr(C₂O₄)₃]³⁻: Trioxalato chromium (III) ion



(iii) [CO(NH₃)₃Cl₃]: Triammine trichloride cobalt (III)




- **25.** (*i*) Chiral molecules (or compounds) are those molecules which are non-superimposible on their mirror images and this property is known as chirality. Butan-2-ol is an example of chiral molecule.
 - (ii) Due to +I effect of alkyl groups the 2° carbonium ion CH₃—CH—CH₂—CH₃ derived from sec. butyl chloride is more stable than the 1° carbonium ion CH₃—CH₂—CH₂ derived from n-propyl chloride. Therefore sec. butyl chloride gets hydrolyzed more easily than n-propyl chloride under S_N1 conditions.



As Iodine is a better leaving group due to its larger size, it will be released at a faster rate in the presence of incoming nucleophile.

26. -Glucose and -Glucose differ only in the configuration of hydroxy group at C₁ and are called anomers and the C₁ carbon is called anomeric carbon. The six membered cyclic structure of glucose is called pyranose structural (-or -), in analogy with pyrane. The cyclic structure of glucose is more correctly represented by Haworth structure as given below.



27.

Thermoplastics		Thermosetting plastics	
(i)	These polymers are linear or slightly branched chain molecules.	(i)	These polymers are cross linked or heavily branched molecules.
(ii)	Soften on heating and harden on cooling and can be remoulded.	(ii)	On heating undergo extensive cross linking in moulds and become infusible.
(iii)	Some common examples are polyethene, PVC, polystyrene, etc.	(iii)	Some common examples are bakelite, urea-formaldehyde resins, terylene, etc.

28. (a) (i) Mole fraction of a particular component in a solution may be defined as the ratio of number of moles of that component to the sum of the moles of all the components present in the solution. For a solution of two components A and B,

Mole fraction of A,
$$x_A = \frac{n_A}{n_A - n_B}$$

Mole fraction of B, $x_B = \frac{n_B}{n_A - n_B}$

- (ii) A solution which obeys Raoult's law over the entire range of concentration and temperature, and during the formation of which no change in enthalpy and volume takes place is called an ideal solution. Thus for an ideal solution,
 - (i) Raoult's law is obeyed, i.e., $P_A = P_A x_A$ and $P_B = P_B x_B$
 - (ii) $H_{mix} = 0$
 - (iii) $V_{mix} = 0$

(b)
$$W_B = 15.0g$$
; $W_A = 450g$; $K_f = 1.86 \text{ K kg mol L}^{-1}$
 $T_f = 0^{\circ}\text{C} - (-0.34^{\circ}\text{C}) = 0.34^{\circ}\text{C or } 0.34 \text{ K}$

Substituting the values in the expression,

$$M_B = \frac{K_f - W_B - 1000}{W_A - T_f}$$

$$M_B = \frac{1.86 \text{ K kg mol}^{-1} \quad 15.0 \text{ g} \quad 1000 \text{ g kg}^{-1}}{450 \text{ g} \quad 0.34 \text{ K}} = 182.35 \text{ g mol}^{-1}$$


OR

- (a) (i) Henry's law states that the solubility of a gas in a liquid is directly proportional to the pressure of the gas.
 - (ii) The boiling point elevation constant may be defined as the elevation in boiling point when one mole of a non-volatile solute is dissolved in one thousand grams of solvent.
- (b) $W_A = 500 \text{g}$; $M_B = 92 \text{g mol}^{-1}$; $K_b = 0.512 \text{ K kg mol}^{-1}$ $T_b = 100.42 \text{°C} - 100 \text{°C} = 0.42 \text{°C} \text{ or } 0.42 \text{K}$

Substituting the values in the expression;

$$W_B = \frac{M_B}{K_b} = \frac{T_b}{1000} = \frac{W_A}{M_b}$$

$$W_B = \frac{92 \text{ g mol}^{-1} \quad 0.42 \text{ K} \quad 500 \text{ g}}{0.512 \text{ K kg mol}^{-1} \quad 1000} \quad 37.73 \text{ g}$$



XeOF.

- (b) (i) It is because S-S single bond is stronger than O-O single bond.
 - (ii) This is because the bond dissociation enthalpy of I–Cl bond is lower than that of I–I bond.
 - (iii) It is due to
 - low enthalpy of dissociation of F-F bond
 - high hydration enthalpy of F -.

OR

- (a) (i) $3\text{Cu} + 8\text{HNO}_3$ (dilute) $3\text{Cu} (\text{NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O}$
 - $(ii) \text{ XeF}_4 + \text{O}_2\text{F}_2 \qquad \text{XeF}_6 + \text{O}_2$
- (b) (i) It is because P-P single bond is stronger than N-N single bond.
 - (ii) Because of its small size, oxygen is capable of forming p -p bond and exist as diatomic O₂ molecule. The intermolecular forces in oxygen are weak van der Waals force, due to which it is a gas at room temperature. On the other hand, sulphur, due to its larger size prefers to form S-S single bond and exist as octaatomic S₈ molecule having puckered ring

- structure. Because of larger size the force of attraction holding the S_8 molecules together are much stronger. Hence sulphur is a solid at room temperature.
- (iii) All halogens are coloured. This is due to absorption of radiation in visible region which results in the excitation of outer electrons to higher energy level while the remaining light is transmitted. The colour of the halogen is the colour of transmitted light.

(b) The compound is 2-ethyl benzaldehyde and the reactions involved in the question are given below:

$$COO^{-} CH_{2}OH$$

$$CH_{2}CH_{3} + CH_{2}CH_{3}$$

$$COO^{-} CH_{2}CH_{3} + CH_{2}CH_{3}$$

$$COO^{-} CH_{2}CH_{3} + CH_{2}CH_{3}$$

$$COO^{-} CH_{2}CH_{3} + CH_{2}CH_{3}$$

$$COO^{-} CH_{2}CH_{3} + COOH$$

$$CH_{2}CH_{3} + CH_{2}CH_{3} + COOH$$

$$COOH COOH$$

$$COOH COOH$$

$$COOH CH_{2}CH_{3} + COOH$$

$$COOH COOH$$

$$COOH CH_{2}CH_{3} + COOH$$

$$COOH CH_{2}CH_{3} + COOH$$

$$COOH COOH$$

$$COOH CH_{2}CH_{3} + COOH$$

$$COOH CH_{3} + COOH$$

$$COOH CH_{4}CH_{3} + COOH$$

$$COOH CH_{4}CH_{4} + COOH$$

$$COOH CH_{4}CH_{4}$$

OR

(a) (i) Propanone on treatment with I₂/NaOH (NaOI) undergoes iodoform reaction to give yellow ppt of iodoform but propanol does not.


(ii) Benzaldehyde being an aldehyde reduces Tollens' reagent to give silver mirror test but acetophenone being a ketone does not give this test.

$$C_6H_5CHO + 2[Ag (NH_3)_2]^+ + 3\bar{O}H$$
 Heat $C_6H_5CO\bar{O} + 2H_2O + 2Ag + 4NH_3$
Benzaldehyde Tollen's reagent Silver mirror

- (b) (i) Methyl tert-butyl ketone < Acetone < Acetaldehyde
 - (ii) 4-Methoxy benzoic acid < Benzoic acid < 3,4-Dinitrobenzoic acid
 - (iii) (CH₃)₂CHCOOH < CH₃CH(Br)CH₂COOH < CH₃CH₂CH(Br)COOH

CBSE (All India) SET-II

- 1. Interstitial defect
- 2. The catalytic reaction that depends upon the pore structure of the catalyst and the size of the reactant and product molecule is known as shape-selective catalysis.
- **3.** Collectors enhance non-wettability of the mineral particles.
- **6.** 3-Phenyl prop-2-enol
- 17. The cleansing action of soap such as sodium stearate is due to the fact that soap molecules form micelle around the oil droplet in such a way that hydrophobic part of the stearate ions is in the oil droplet and hydrophilic part projects out of the grease droplet like the bristles. Since the polar groups can interact with water, the oil droplet surrounded by stearate ions is now pulled in water and removed from the dirty surface. Thus soap helps in emulsification and washing away of oils and fats.



Hard water contains calcium and magnesium salts. In hard water, soap gets precipitated as calcium and magnesium soap which being insoluble stick to the clothes as gummy mass. Therefore soaps do not work in hard water.

20. At anode: Al
$$Al^{3+} + 3e^{-}]_{x2}$$
At Cathode: $Ni^{2+} + 2e^{-}$ $Ni]_{x3}$
Cell reaction: $2Al + 3Ni^{2+}$ $2Al^{3+} + 3Ni$

$$E^{o}_{Cell} = E^{o}_{Cathode} - E^{o}_{anode} = E^{o}_{Ni^{2}/Ni} - E^{o}_{Al^{3}/Al}$$

$$= -0.25V - (-1.66 \text{ V}) = 1.41 \text{ V}$$

$$[Al^{3+}] = 1 \times 10^{-3} \text{M}; [Ni^{2+}] = 0.5 \text{ M}; n = 6$$
Substituting the values in the Nernst equation,
$$E_{cell} = E^{o}_{cell} - \frac{0.059}{n} \log \frac{[Al^{3}]^{2}}{[Ni^{2}]^{3}}$$

$$E_{cell} = 1.41 \text{ V} - \frac{0.059}{n} \log \frac{(10^{-3})^{2}}{n}$$

$$E_{\text{cell}} = E_{\text{cell}} = \frac{1.09}{n} \log \frac{100}{[\text{Ni}^2]^3}$$

$$E_{\text{cell}} = 1.41 \text{ V} - \frac{0.059}{6} \log \frac{(10^{-3})^2}{(0.5)^3}$$

$$= 1.41 \text{ V} - \frac{0.059}{6} \log 8 \times 10^{-6}$$

$$= 1.41 \text{ V} - \frac{0.059}{6} (-0.54)$$

$$E_{cell} = 1.41 \text{ V} + 0.0053 \text{ V}$$


$$= 1.4153 \text{ V}$$

- 23. (i) Interstitial compounds are well known for many of the transition elements because the transition elements are capable of entrapping small sized atoms such as H,C and N in the interstitial sites in their crystal lattices. These trapped atoms get bonded to the atoms of transition elements, for example, TiC, Fe₃H and Mn₄N, etc.
 - (ii) The decrease in metallic radius coupled with increase in atomic mass results in a general increase in density from titanium to copper in the first series of transition elements.
 - (iii) This is due to very small energy gaps between 5f, 6d and 7s subshells in actinoids.
- 27. (i) Elastomers: These are the polymers having the weakest intermolecular forces of attraction between the polymer chains. The weak forces permit the polymer to be stretched. A few 'cross links' are introduced between the chains, which help the polymer to retract to its original position after the force is released as in vulcanised rubber. Elastomers thus possess an elastic character, e.g., buna-S, buna-N, neoprene, etc.
 - (ii) Condenstaion Polymers: The condensation polymers are formed by the repeated condensation reaction between different bifunctional or trifunctional monomer units usually with elimination of small molecules such as water, alcohol hydrogen chloride, etc. Nylon 6, 6, Nylon 6, terylene are some examples.
 - (iii) Addition polymers: Addition polymers are formed by repeated addition of same or different monomer molecules. The monomers used are unsaturated compounds, e.g., alkenes, alkadienes and their derivatives. Polythene is an example of addition polymer.

$$n$$
 CH₂=CH₂ $\xrightarrow{350-570\,\mathrm{K},\,1000-2000\,\mathrm{atm}}$ (CH₂ — CH₂) \xrightarrow{n} Ethene Polyethene

Polyethene


30.



(ii) No. of electron pairs around central atom Cl = 5No. of bond pairs = 3

No of lone pairs = 2

The shape would be slightly bent T.



- (b) (i) Nitrogen is much less reactive than phosphorus because of the high bond enthalpy of N N bond.
 - (ii) This is because in H₂O hydrogen bond formed is three dimensional whereas in H-F it is linear
 - (iii) This is because S-S single bond is stronger than O-O single bond.

OR

$$(a) \quad (i) \quad \begin{picture}(60,0) \put(0.5){\line(1,0){100}} \put(0.5){\l$$

- (b) (i) Due to smaller size of O as compared to S, the bond dissociation enthalpy of O-H bond is higher than that of S-H bond. As a result, in aqueous solution, S-H bond can break more easily form H⁺ ion than O-H bond. Hence H₂S is more acidic than H₂O.
 - (ii) This is because fluorine is the most electronegative element and it does not have d orbitals.
 - (iii) This is because the valence shell orbital of He is completely filled $(1s^2)$ and it has high ionisation enthalpy and more positive electron gain enthalpy.

CBSE (All India) SET-III

- 1. Silicon or germanium doped with group 15 elements like P or As are called n type of semiconductors.
- **4.** H₃PO₂ is monobasic as it contains only one ionisable H-atom which is present as OH group.

5.
$$H_2C = C - CH_2 - Br$$

3-Bromo-2-methyl propone

7. On prolonged heating with HI, glucose gives n-hexane.

- **8.** Human beings suffer from depression when they have low levels of noradrenaline. Noradrenaline is a neurotransmiter that plays a role in mood changes. Low levels of noradrenaline lower the signal-sending activity and make human beings suffer from depression. Tranquilizers such as improniazid and phenelyzine are useful in treating this depression.
- 11. (i) In the extraction of silver, the metal is leached with dilute solution of NaCN. Dilute NaCN solution forms a soluble complex with Ag while the impurities remain unaffected which are then filtered off.

$$4Ag + 8NaCN + O_2 + 2H_2O$$
 $4Na[Ag(CN)_2] + 4NaOH$ Soluble complex

(ii) SiO_2 acts as a flux in the extraction of copper. Sulphide ore of copper contains iron as impurity which is removed as iron silicate (slag).

$$FeO + SiO_2$$
 $FeSiO_3$ $Slag$

18. Difference between antiseptics and disinfectants:


Antiseptics

- Antiseptics are chemical substances which prevent the growth of microorganisms and may even kill them but are not harmful to living tissues.
- Antiseptics are generally applied to living tissues such as wounds, cuts, ulcers and diseased skin surfaces.
- Dettol, furacine, soframicine are antiseptics.

Disinfectants


- Disinfectants are chemical substances which kill microorganisms or stop their growth but are harmful to human tissues.
- Disinfectants are applied to inanimate objects such as floor, drainage system, instrument, etc.
- Chlorine in the concentration of 0.2 to 0.4 ppm in aqueous solution and SO₂ in very low concentration are disinfectants.
- **22.** Three distinct features of chemisorptions which are not found in physisorptions:
 - (i) Enthalpy of adsorption: Enthalpy of adsorption is high (80-240 kJ mol⁻¹) as it involves chemical bond formation.
 - (ii) **High specificity:** Chemisorption is highly specific and it will only occur if there is some possibility of chemical bonding between adsorbent and adsorbate. For example, hydrogen is adsorbed by transition metals by virtue of hydride formation.
 - (iii) Irreversibility: Chemisorption is usually irreversible in nature as it involves compound formation. It is very slow at low temperatures on account of high activation energy.
- 23. (i) Cr^{2+} is reducing agent as its configuration changes from d^4 to d^3 , the later having a half-filled trg level. On the other hand, the change from Mn^{2+} to Mn^{3+} results in the extra stable half-filled, d^5 configuration. Therefore Mn^{3+} is an oxidising agent.
 - (ii) This is due to small energy gap between 5f, 6d and 7s subshells in actinoids.
 - (iii) This is because with increase in atomic number in a series, the increased nuclear charge is partly cancelled by the increased shielding effect of electrons in the *d*-orbitals of penultimate shell.

24. (i) $[Fe(CN)_6]^{4-}$ = Hexacyanoferrate(II) ion; $Fe^{2+}(3d^6)$




 d^2sp^3 -hybridisation in $[Fe(CN)_6]^{4-}$ leads to octahedral structure.

(ii) $[Cr(NH_3)_4Cl_2]^+$ = Tetraamminedichloridochromium (III) ion, $Cr^{3+}(3d^3)$



 d^2sp^3 hybridisation in $[Cr(NH_3)_4Cl_2]^+$ leads to octahedral structure.

(iii) $[Ni(CN)_4]^{2-}$ = Tetracyanonickelate(II)ion; $Ni^{2+}(II)$ (3 d^8)



dsp² hybridisation in [Ni(CN)₄]²⁻ leads to square planar structure.

- **26.** (i) A peptide linkage is an amide linkage —(CONH)— formed between —COOH group of one -amino acid and —NH₂ group of the other -amino acid by the loss of a water molecule.
 - (ii) Primary structure of proteins: The sequence in which various amino acids are arranged in a protein is called its primary structure. Any change in the sequence of amino acids creates different protein which alters biological functions.
 - (iii) Denaturation of Proteins: When a protein in its native form is subjected to a change in temperature or a change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix gets uncoiled and protein loses its biological activity. This is called denaturation of protein. During denaturation 2° and 3° structures are destroyed but 1° structures remain intact, e.g., coagulation of egg white on boiling, curdling of milk, etc.

CBSE EXAMINATION PAPERS

FOREIGN-2012

Time allowed: 3 hours [Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question nos. 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question nos. 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question nos. 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question nos. 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables if necessary, use of calculators is not allowed.

CBSE (Foreign) SET-I

- 1. What is meant by the term 'forbidden zone' in reference to band theory of solids?
- **2.** Why is the adsorption phenomenon always exothermic?
- 3. Write the reaction involved in the extraction of silver after the silver ore has been leached with NaCN.
- **4.** Although the H-bonding in hydrogen fluoride is much stronger than that in water, yet water has a much higher boiling point than hydrogen fluoride. Why?
- 5. Write the IUPAC name of the following compound:

$$\begin{array}{c} \operatorname{H_3C--CH=-CH---CH_2---CH_3} \\ \operatorname{OH} \end{array}$$

- **6.** Draw the molecular structure of the compound, 4-methylpent-3-en-2-one.
- 7. Write the full form of DNA and RNA.
- **8.** What is meant by 'narrow spectrum antibiotics'?
- **9.** (a) Define any two of the following terms:
 - (i) van't Hoff factor
 - (ii) Mole fraction
 - (iii) Ebullioscopic constant
 - (b) State Raoult's law.

OR

The density of water of a lake is 1.25 g (mL)^{-1} and one kg of this water contains 92 g of Na⁺ ions. what is the molarity of Na⁺ ions in the water of the lake? (Atomic mass of Na = 23.00 u)

- **10.** Define the following terms:
 - (i) Order of a reaction
 - (ii) Activation energy of a reaction

- Name one chief ore each of copper and aluminium. Name the method used for concentration of these two ores.
- **12.** Explain the following:
 - (i) The chemical reactivity of nitrogen is much less than that of phosphorus.
 - (ii) SF₆ is kinetically inert.
- 13. Draw the molecular structures of the following species:
 - (i) H₃PO₃
 - (ii) BrF₃
- **14.** What are ambident nucleophiles? Explain giving an example.
- 15. Explain as to why
 - (i) Alkyl halides, though polar, are immiscible with water.
 - (ii) Grignard's reagents should be prepared under anhydrous conditions.
- **16.** Describe the following giving the chemical equation in each case:
 - (i) Carbylamine reaction
 - (ii) Hofmann's bromamide reaction
- 17. Complete the following reaction equations:
 - (i) $C_6H_5N_2Cl + H_3PO_2 + H_2O$
 - (ii) C₆H₅NH₂ + Br₂(aq)
- **18.** State a reason for each of the following statements:
 - (i) Soaps do not work in hard water.
 - (ii) The use of the sweetner aspartame is limited to cold foods and drinks.
- 19. Iron has a body centred cubic (bcc) unit cell with a cell dimension of 286.65 pm. The density of iron is 7.874 g cm⁻³. Use this information to calculate Avogadro's number. (Atomic mass of Fe = 55.845 u)

OR

Silver crystallises in face centred cubic (fcc) unit cell. If the radius of silver atom is 145 pm, what is the length of each side of the unit cell?

- **20.** At 25°C the saturated vapour pressure of water is 3.165 kPa (23.75 mm Hg). Find the saturated vapour pressure of a 5% aqueous solution of urea (carbamide) at the same temperature. (Molar mass of urea = 60.05 g mol⁻¹)
- **21.** Consider the reaction:

$$2A + B$$
 $C + D$

Following results were obtained in experiments designed to study the rate of reaction:

Experiment	Initial concentr	ration (mol L ⁻¹)	Initial rate of formation
No.	[A]	[B]	
1	0.10	0.10	1.5×10^{-3}
2	0.20	0.20	3.0×10^{-3}
3	0.20	0.40	6.0×10^{-3}

- (a) Write the rate law for the reaction.
- (b) Calculate the value of rate constant for the reaction.
- (c) Which of the following possible reaction mechanism is consistent with the rate law found in (a)?

I.
$$A + B$$
 $C + E$ (slow)
$$A + E$$
 D (fast)

II. $B + C$ E (slow)
$$A + E$$
 F (fast)
$$A + F$$
 D (fast)

- **22.** Define the following terms giving one suitable example for each:
 - (i) Electrophoresis
 - (ii) Micelles
 - (iii) Peptization
- **23.** Complete the following chemical equations:
 - (i) $NH_4Cl(aq) + NaNO_2(aq)$
 - (ii) P₄ + NaOH + H₂O
 - $(iii) \begin{array}{c} {\rm Xe}(g) \\ {\rm (Xenon\ in\ excess)} \end{array} + \begin{array}{c} {\rm F_2}(g) \end{array} \quad {}^{\rm 673\ K}_{\rm 1\ Bar}$
- **24.** Explain the following:
 - (i) The -complexes are known for transition elements only.
 - (ii) Nickel(II) does not form low spin octahedral complexes.
 - (iii) [Fe(CN)₆]⁴⁻ and [Fe(H₂O)₆]²⁺ are of different colours in dilute solutions.
- 25. State the products of the following reactions:
 - (i) CH₃—CH₂—CH₂—O—CH₃ + HBr

$$(ii) \bigcirc ^{\mathrm{OC_2H_5}} + \mathrm{HBr}$$

- **26.** What is glycogen? How is it different from starch? How is starch structurally different from cellulose?
- **27.** Write the structure of the monomer of each of the following polymers:
 - (i) Nylon-6
 - (ii) Teflon
 - (iii) Neoprene
- **28.** (a) What type of a battery is the lead storage battery? Write the anode and the cathode reactions and the overall reaction occurring in a lead storage battery sending out an electric current.
 - (b) A voltaic cell is set up at 25°C with the half-cells

$$Ag^{+} \; (0.001 \; M) \mid Ag \; and \; Cu^{2+} \; (0.10M) \mid Cu$$

What should be its cell potential? ($E_{cell}^{o} = 0.46 \text{ V}$, log $10^{5} = 5$)

OR

- (a) Define the term molar conductivity and explain how molar conductivity changes with solution concentration for weak and strong electrolytes.
- (b) A strip of nickel metal is placed in a 1-molar solution of Ni(NO₃)₂ and a strip of silver metal is placed in a 1-molar solution of AgNO₃. An electrochemical cell is created when the two solutions are connected by a salt bridge and the two strips are connected by wires to a voltmeter.
 - (i) Write the balanced equation for the overall reaction occurring in the cell and calculate the cell potential
 - (ii) Calculate the cell potential, E, at 25°C for the cell if the initial concentration of $Ni(NO_3)_2$ is 0.100 molar and the initial concentration of AgNO₃ is 1.00 molar.

$$[E_{Ni^{2+}/Ni}^{o} = -0.25 \text{ V}; E_{Ag^{2+}/Ag}^{o} = 0.80 \text{ V}, \log 10^{-1} = -1]?$$

- **29.** (a) Complete the balance the following chemical equations:
 - (i) $Cr_2O_7^{2-} + I^- + H^+$
 - $(ii) \text{ MnO}_4^- + \text{SO}_3^{2-} + \text{H}^+$
 - (b) Explain the following observations:
 - (i) Transition elements and their compounds are known to act as catalysts.
 - (ii) The higher oxidation states are usually exhibited by the members in the middle of a series of transition elements.
 - (iii) The metal-metal bonding is more frequently found with the second and third series of transition elements.

OR

(a) Calculate the number of unpaired electrons in the following gaseous state ions:

Which one of these is the most stable in aqueous solutions?

(At. nos.
$$V = 23$$
, $Cr = 24$, $Mn = 25$, $Fe = 26$)

- (b) Explain the following observations:
 - (i) The transition metal ions are usually coloured in aqueous solutions.
 - (ii) Cu(I) is not stable in an aqueous solution.
 - (iii) The highest oxidation state of a transition metal is exhibited in its oxide or fluoride.
- **30.** (a) Describe the mechanism of the addition of Grignard reagent to the carbonyl group of a compound to form an adduct which on hydrolysis yields an alcohol.
 - (b) Draw the structures of the following compounds:
 - (i) 3-Methylbutanal
 - (ii) Hexane-1,6-diotic acid
 - (iii) p-Nitropropiophenone

OR

- (a) Illustrate the following reactions giving a suitable chemical equation for each:
 - (i) Cannizzaro reaction
 - (ii) Hell-Volhard-Zelinsky reaction

- (b) How would you bring about the following conversions? Write the complete equation in each case.
 - (i) Ethanol to 3-hydroxybutanal
 - (ii) Benzoic acid to m-nitrobenzyl alcohol
 - (iii) Benzaldehyde to benzophenone

CBSE (Foreign) SET-II

Questions Uncommon to Set-I

- 1. How would you distinguish between a metallic solid and an ionic solid other than by metallic lustre?
- 2. What is meant by 'shape selective catalysis'?
- **4.** Which is more acidic and why, H₂O or H₂S?
- **5.** Of the two alcohols; (a) CH₂=CH—CH₂OH and (b) CH₂=CH—CH₂—CH₂OH, which one will react more easily with conc. HCl in the presence of ZnCl₂?
- **8.** Of the two bases named below, which one is present in RNA and which one is present in DNA? Thymine, Uracil
- 13. Draw the molecular structures of the following species:
 - (i) H₂S₂O₈
 - (ii) XeF₂
- 18. What are biodegradable and non-biodegradable detergents? Give one example of each.
- **22.** Differentiate among a homogeneous solution, a suspension and a colloidal solution, giving a suitable example of each.
- **23.** Complete the following chemical equations:
 - (i) $HgCl_2 + PH_3$
 - (ii) NaOH + Cl₂

(hot & conc.)

- (*iii*) $XeF_4 + O_2F_2$ 143 K
- 24. Name the following complexes and draw the structures of one possible isomer of each:
 - (i) $[Cr(C_2O_4)_3]^{3-}$
 - (ii) $[Pt(NH_3)_2Cl_2]$
 - (iii) $[Co(en)_2Cl_2]^+$
- **26.** Explain the meaning of the following terms:
 - (i) Invert sugar
 - (ii) Polypeptides
 - (iii) Enzymes

CBSE (Foreign) SET-III

Questions Uncommon to Set-I and II

- 1. Define paramagnetism with an example.
- 3. What is the role of a depressant in Froth Floatation process for the concentration of a sulphide ore?
- **4.** Which is a stronger acid in aqueous solution, HF or HCl, and why?
- **5.** Write the IUPAC name of the following compound:

- **6.** Draw the molecular structure of 4-Hydroxy-4-methylpentan-2-one.
- **12.** State a reason for each of the following statements:
 - (i) Fluorine never exhibits any positive oxidation state.
 - (ii) Helium does not form any real chemical compound.
- **14.** Explain why
 - (i) the dipole moment of chlorobenzene is lower than that of cyclohexyl chloride.
 - (ii) haloalkanes are only slightly soluble in water but dissolve easily in organic solvents.
- **20.** Calculate the freezing point depression expected for 0.0711 m aqueous solution of sodium sulphate. If the solution actually freezes at -0.320 °C, what is the actual value of van't Hoff factor at this temperature? (K_f for water = 1.86 K kg mol⁻¹)
- **22.** Present a classification of colloids where dispersion medium is water. State the characteristics and one example of each of these classes.
- **23.** Complete the following chemical equations:

(i)
$$Sn + 2PCl_5$$
 heatin

(ii)
$$Fe^{3+} + SO_2 + H_2O$$

(iii)
$$XeF_2(s) + H_2O(l)$$

26. Write three such behaviours of glucose which cannot be explained by an open chain structure of glucose molecule. What alternative structure has been proposed for the glucose molecule?



SET-I

- 1. The energy gap between valance band and conduction is known as forbidden zone.
- **2.** As the adsorption progresses, the residual forces at the surface decreases resulting in the decrease of surface energy which appears as heat.
- **3.** $2[Ag(CN)_2](aq)$ Zn(s) $[Zn(CN)_4]^2(aq)$ 2Ag(s)
- 4. This is because hydrogen bonding is multidimensional in water whereas in HF it is linear.
- 5. Hex-4-en-3-ol

7. DNA: Deoxyribonucleic acid

RNA: Ribonucleic acid

- **8.** Antibiotics which are mainly effective against gram-positive or Gram-negative bacteria are known as narrow spectrum antibiotics, *e.g.*, ampicillin G.
- **9.** (a) Van't Hoff factor:
 - (i) May be defined as the ratio of normal molecular mass to observed molecular mass or the ratio of observed colligative property to calculated colligative property.
 - (ii) Mole fraction may be defined as the ratio of number of moles of one component to the sum of moles all the components present in a solution.
 - (iii) Ebuthoscopic constant may be defined as the elevation in boiling point when one mole of a non-volatile solute is added to 1000 grams of solvent.
 - (b) It states that for a solution of volatile liquids the partial vapour pressure of each component in the solution is directly proportional to its mole faction.

Number of moles of solute, Na⁺ ions =
$$\frac{92g}{23g \text{ mol}^{-1}}$$
 = 4 mol

Volume of solution =
$$\frac{\text{Mass of solution}}{\text{Density of solution}}$$
 $\frac{1000 \text{ g}}{1.25 \text{ g mL}^{-1}}$ 800 mL $\frac{800 \text{ mL}}{1000 \text{ mL L}^{-1}} = 0.8 \text{ L}$

Molarity =
$$\frac{\text{Number of moles Na}}{\text{Volume of solution in liter}}$$
 $\frac{4 \text{ mol}}{0.8 \text{ L}}$
= 5 mol L⁻¹ or 5 m.

10. (*i*) The sum of the powers of the concentration of the reactants in the rate law expression is called the order of the reaction. For a general reaction

Let rate =
$$K[A]^m[B]^n$$

Order of the reaction = m + n

(ii) The minimum extra energy absorbed by the reactant molecular so that their energy becomes equal threshold energy is called activation energy.

Activation energy = Threshold energy - Energy possessed by reactant molecular


11.

Metal	Chief ore	Method of concentration	
Copper	Copper pyrite (CuFeS ₂) Froth floatation		
Aluminium	Bauxite $[AlO_x(OH)_{3-2x}]$	Leaching	
	where $0 < x < 1$		

- **12.** (*i*) As N≡N triple bond (941.4 kJmol⁻¹) is much stronger than p–p single bond (213 kJmol⁻¹), therefore nitrogen is much less reactive than phosphorus.
 - (ii) This is because in SF₆, sulphur is sterically protected by six fluorine atoms.
- **13.** (*i*) H₃PO₃







14. Nucleophiles which can attack through the different nucleophilic centres present in it are called ambident nucleophiles. For example, cyanide group is resonance hybrid of the following two contributing structures:

$$\stackrel{\ominus}{:}\stackrel{\frown}{C}=\stackrel{\frown}{N}:$$
 \longleftrightarrow $:\stackrel{\frown}{C}=\stackrel{\frown}{N}:$

It can attack through carbon to form cyanide and through N to form isocyanide.

- **15.** (*i*) This is due to inability of alkyl halides to form intermolecular hydrogen bonds with water molecules.
 - (ii) This is because Grignard reagent forms alkanes by reacting with moisture.

$${\rm RMgX} \quad {\rm H_2O} \qquad \quad {\rm RH} \quad {\rm Mg(OH)X}$$

16. (*i*) Carbylamine reaction: Aliphatic and aromatic primary amines when heated with chloroform and ethanolic potassium hydroxide form carbylamines or isocyanides which are foul smelling substances. Secondary and tertiary amines do not show this reaction.

$$R - NH_2 + CHCl_3 + 3KOH (alc.) \qquad RNC + 3KCl + 3H_2O$$

$$C_6H_5NH_2 + CHCl_3 + 3KOH (alc.) \qquad C_6H_5NC + 3KCl + 3H_2O$$
Aniline
$$Phenyl$$
isocyanide

(ii) Hoffmann's bromamide reaction: When a primary acid amide is heated with bromine in an aqueous or ethanolic solution of NaOH, a primary amine is obtained. The amine so obtained contains one carbon less than that present in the amide.

$$\begin{array}{c} \text{RCONH}_2 + \text{Br}_2 + 4\text{NaOH} & \text{R-NH}_2 + \text{Na}_2 \text{ CO}_3 + 2\text{NaBr} + 2\text{H}_2\text{O} \\ \text{Acid amide} & \text{C}_6\text{H}_5\text{CONH}_2 + \text{Br}_2 + 4\text{NaOH} & \text{C}_6\text{H}_5\text{NH}_2 + \text{Na}_2\text{CO}_3 + 2\text{NaBr} + 2\text{H}_2\text{O} \\ \text{Benzamide} & \text{Aniline} & \text{Aniline} \\ \\ \textbf{17.} \quad \textit{(i)} \ \text{C}_6\text{H}_5\text{N}_2\text{Cl} + \text{H}_3\text{PO}_2 + \text{H}_2\text{O} & \text{C}_6\text{H}_6 + \text{N}_2 + \text{H}_3\text{PO}_3 + \text{HCl} \\ \text{Benzene} & \text{diazonium chloride} & \text{NH}_2 & \text{NH}_2 \\ \hline \end{array}$$

- **18.** (*i*) In hard water soaps get precipitated as calcium and magnesium soap which being insoluble in water stick to the clothes as gummy mass.
 - (ii) This is because aspartame is unstable at cooking temperature.

19. Here,
$$z = 2$$
, $M = 56 \text{ g mol}^{-1}$, $d = 7.87 \text{ g cm}^{-3}$
 $a = 286.65 \text{ pm} = 286.65 \times 10^{-10} \text{ cm} = 2.8$

Substituting the values in the expression,

$$N_A = \frac{z M}{a^3 d}$$
, we get

$$N_A = \frac{2 56 \text{ g mol}^{-1}}{(286.65 \ 10^{-10} \text{ cm})^3 \ 7.87 \text{ g cm}^{-3}}$$

$$= 6.042 \times 10^{23} \text{ mol}^{-1}$$

OR

For
$$bcc$$
 unit cell,
$$a = 2\sqrt{2}r$$

$$a = 2\sqrt{2} \times 145 \text{ pm}$$

$$= 2 \times 1.414 \times 145 \text{ pm}$$

$$a = 410.06 \text{ pm}.$$

20. W_B 5 g, W_A 95 g, M_B 60.05 g mol⁻¹, MA = 18 g mol⁻¹ P°A = 3.165 kPa

Substituting the values in the expression;

$$\frac{P_A^{\circ} P}{P_A^{\circ}} = \frac{W_B M_A}{M_B W_A}, \text{ we get}$$

$$\frac{3.165 \text{ k Pa } P}{3.165 \text{ k Pa}} = \frac{5 \text{ g}}{60.05 \text{ g mol}} = \frac{1}{95 \text{ g}} = 0.015$$

$$P = 3.165 \text{ k Pa} - 0.015 \times 3.165 \text{ k Pa}$$

$$P = 3.118 \text{ k Pa}$$

21. (a) Let Rate =
$$k[A]^m [B]^n$$

$$1.5 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1} \quad K(0.1)^m (0.1)^n \quad \dots (i)$$

3.0 10
3
 mol L^{1} min⁻¹ $K(0.2)^{m}(0.2)^{n}$... (ii)

$$6.0 \times 10^{-3} \text{ mol } L^{-1} \text{ min}^{-1} K(0.2)^m (0.4)^n \dots (iii)$$

Dividing equation (iii) by (ii), we get

$$\frac{6.0 \quad 10^{-3}}{3.0 \quad 10^{-3}} = \frac{k(0.2)^m (0.4)^n}{k(0.2)^m (0.2)^n}$$

$$2 2^n n = 1$$

Dividing equation (ii) by (i), we get

$$\frac{3.0 \quad 10^{-3}}{1.5 \quad 10^{-3}} = \frac{k(0.2)^m (0.2)^n}{k(0.1)^m (0.1)^n}$$

$$2 = 2^m 2^n \quad 2^m . 2$$

$$2^m = 1 \quad \text{or} \quad 2^m \quad 2^0 \qquad m = 0$$
Rate $K[A] \quad [B]^1 \quad \text{or Rate} = K[B]$

(b)
$$K \quad \frac{\text{Rate}}{[B]}$$

$$K = \frac{1.5 \cdot 10^{-3} \,\text{mol } L^{-1} \,\text{min}^{-1}}{0.1 \,\text{mol } L^{-1}} = 1.5 \times 10^{-2} \,\text{min}^{-1}$$

- (c) The reaction mechanism (II) is consistent with the rate law found in (a).
- **22.** (*i*) The movement of colloidal particles towards oppositely charged electrodes in an electric field is called electrophoresis.
 - (ii) There are some substances such as soap which at low concentration behave as normal electrolytes, but at higher concentration exhibit colloidal behaviour due to the formation of aggregates. The aggregated particles thus formed are known as micelles or associated colloids.
 - (iii) The process of converting a procipitate into colloidal solution by shaking it with dispersion medium in the presence of small amount of electrolyte is called peptization.

23. (i)
$$NH_4Cl(aq)$$
 $NaNO_2(aq)$ $N_2(g)$ $2H_2O(l) + NaCl(aq)$

(iii)
$$Xe(g)$$
 $F_2(g)$ $\frac{673K}{lhar}$ $XeF_2(s)$

(Xenon in excess)

- **24.** (*i*) This is because the transition metals have empty *d*-orbitals into which the electron pairs can be denoted by ligands containing and electrons, *e.g.*, C₆H₆, C₂H₄ etc.
 - (ii) As only one inner d-orbital is available in nickel for bonding in the presence of strong ligand, e.g., CO, CN.

- (iii) In both the complexes, Fe is in +2 oxide state with d^6 configuration. As the ligands CN and H₂O possess different crystal field splitting energy (O), they absorb different components of visible light for d-d transition. Hence, the transmitted colours are different in dilute solutions.
- $(i) \ \operatorname{CH_3--CH_2--CH_2--O--CH_2--CH_3--H--Br} \qquad \operatorname{CH_3--CH_2--Br---CH_3--CH_2--OH} \\ \operatorname{OC_2H_5} \qquad \qquad \operatorname{OH} \\ (ii) \qquad + \ \operatorname{HBr} \qquad \longrightarrow \qquad \operatorname{C_2H_5Br} \quad + \qquad \qquad \\$
 - (iii) (CH₃)₃C \longrightarrow OC₂H₅ HI $(CH_3)_3C-I$ C_2H_5-OH
- **26.** Glycogen is a polymer of D glucose. The carbohydrates are stored in animal body as glycogen. Starch is also a polymer of D glucose and consist of two components amylose and amylopectin. Amylose is linear chain polymer of D-glucose. Both glycogen and amylopectin are branched chain polymer of D glucose but glycogen is more highly branched than amylopectin. Strarch is the main storage polysaccharide of plants.

Starch is a polymer of D glucose whereas cellulose is a polymer -D-glucose.

27.

S.No.	Polymer	Monomer	Structure of Monomer
(i)	Nylon-6	Caprolactum	$H_{2}C$ $C=O$ $H_{2}C$ CH_{2} $H_{2}C-CH_{2}$ Caprolactum
(ii)	Teflon	Tetrafluoro ethene	CF ₂ =CF ₂
(iii)	Neoprene	Chloroprene	$CH_2 = C - CH = CH_2$ CI

28. (a) Lead storage battery is a secondary battery. The reactions occurring in lead storage battery when current is drawn from it are:

 $Pb(s) + SO_4^{2-}(aq)$ $PbSO_4(s) + 2e^{-}$ Anode: $PbO_2(s) + SO_4^{2-}(aq) + 4H^+(aq) + 2e$ Cathode:

 $PbSO_4(s) + 2H_2O(l)$

 $2\text{PbSO}_4(s) + 2\text{H}_2\text{O}(l)$ $Pb(s) + PbO_{2}(s \quad 2H_{2}SO_{4}(aq))$ Overall reaction:

Cu $Cu^{2+} + 2e^{-}$ (b) At Anode: Ag + e Ag] 2 $Cu + 2Ag^{+} Cu^{2+} + 2Ag$ At Cathode:

Cell reaction:

 $E_{\text{cell}}^{\text{o}} = 0.46 \text{ V}; [\text{Ag}^{+}] = 1 \times 10^{-3} \text{ M}, [\text{Cu}^{2+}] = 0.1 \text{M} : n = 2$

Substituting the values in the equation:

$$\begin{split} E_{\text{cell}} &= E_{\text{cell}}^{\,0} - \frac{0.059}{n} \log \frac{[\text{Cu}^{\,2}]}{[\text{Ag}]^2}, \text{ we get} \\ E_{\text{cell}} &= 0.46 \text{ V} - \frac{0.059}{2} \log \frac{(0.1)}{(10^{-3})^2} \\ &= 0.46 \text{ V} - (0.0295 \log 10^5) = 0.46 \text{V} - 0.0295 \times 5 \\ &= 0.3125 \text{ V} \end{split}$$


OR

(a) Molar conductivity of a solution at a given concentration is the conductance of the volume V of the solution containing one mole of electrolyte kept between two electrodes with area of cross section A and distance of unit length.

$$V = \frac{1000}{c}$$

where is the conductivity and V is the volume of solution containing one mole of the electrolyte and c is the molar concentration.

Molar conductivity increases with decrease in concentration or increase in dilution as the



number of ions as well as the mobility of ions increases with increase in dilution.

For strong electrolytes, the number of ions does not increase appreciably on dilution and only mobility of ions increases due to decrease in interionic attractions. Therefore, $_m$ increases a little as shown in figure by a straight line. For weak electrolytes, number of ions as well as mobility of ions increases on dilution which results in a very large increase in molar conductivity, especially at infinite dilution (*i.e.*, concentration, c 0) as shown by curve in figure.

(b) (i) At Anode: Ni Ni²⁺+ 2e⁻
At Cathode: Ag + e Ag] 2

Cell reaction: Ni + 2Ag + Ni²⁺ + 2Ag

$$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$$

$$= E_{\text{Ag}^{+}/\text{Ag}}^{\circ} \quad E_{\text{Ni}^{2+}/\text{Ni}}^{\circ} = 0.80 \text{ V} - (-0.25 \text{ V})$$

$$E_{\text{cell}}^{\circ} = 1.05 \text{ V}$$
(ii)
$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{n} \log \frac{[Ni^{2}]}{[Ag]^{2}}$$
Here, $n = 2$, $E_{\text{cell}}^{\circ} = 1.05 \text{ V}$, $[Ni^{2+}] = 0.1 \text{ M}$, $[Ag^{+}] = 1.0 \text{ M}$

$$E_{\text{cell}}^{\text{o}} = 1.05 \text{V} - \frac{0.059}{2} \log \frac{(0.01)}{(1)^2}$$

 $E_{\text{cell}} = 1.05 \text{V} - 0.0295 \log 10^{-1} = 1.05 + 0.0295 \text{ V}$
 $E_{\text{cell}} = 1.0795 \text{ V}$

(b) (i) The catalytic activity of transition metals and their compounds is attributed to the following reasons:

Due to their tendency to show variable oxidation states transition metals form instable intermediate compounds and provide a new path for the reaction with lower activation energy.

In Some cases, the transition metals provide a suitable large surface area with free valencies on which reactants are adsorbed.

- (ii) This is due to presence of maximum number of unpaired electrons in a transition metal which is present in the middle of a series.
- (iii) In the same group of *d*-block elements, the 4*d* and 5*d* transition elements are larger in size than than those of 3*d* elements. Hence, the valence electrons are less tightly held and form metal–metal bond more frequently.


OR

(*a*)

Ion	Electronic Configuration	Number of Unpaired electrons
Mn ²⁺ [Ar]	$3d^54s^0$	5
Cr ³⁺ [Ar]	$3d^34s^0$	3
V ³⁺ [Ar]	$3d^24s^0$	2
Fe ²⁺ [Ar]	$3d^64s^0$	4

Mn²⁺ ion is most stable in aqueous solution.

(b) (i) Ni(CO)₄: Tetracrabonyl nickel(o); Ni = $3d^8 4s^2$



Structure = Tetrahedral; Magnetic behaviour : Diamagnetic

(ii) Cu^{2+} (aq) is much more stable than Cu(aq). This is because high negative enthalpy of hydration of $Cu^{2+}(aq)$ easily compensates the high second ionisation enthalpy of copper. Due to his $Cu^+(aq)$ undergo disproportionation as follows:

 Cu^2 (aq) 2Cu (aa) Cu(s)

(iii) This is due to high electronegativity oxygen and fluorine.

30. (a) Mechanism:

(i) Nucleophilic addition of Grignard reagent to carbonyl group to form an adduct.

$$C \stackrel{\stackrel{-}{=} O}{=} V \stackrel{\stackrel{-}{=} V}{=} X \longrightarrow \begin{bmatrix} C \stackrel{\stackrel{-}{=} V}{=} C \stackrel{\stackrel{-}{=} V}{=} C \end{bmatrix}$$

(ii) Hydrolysis of the adduct to alcohol.

$$\begin{bmatrix} \searrow_{C} - \bar{O} - \mathring{M}g - X \end{bmatrix} \xrightarrow{H_2O} \searrow_{C} - O + Mg(OH)X$$

- (b) Name of compound
 - (i) S–Methyl butanal

(ii) Hexane-1, 6-dioic acid

$$HOOC$$
— CH_2 — CH_2 — CH_2 — CH_2 — $COOH$

(iii) P-Nitropropiophenone

$$NO_2$$

OR

(i) Cannizzaro reaction: Aldehydes which do not have an- hydrogen atom, undergo (*a*) disproportionation reaction on treatment with concentrated alkali. In this reaction, one molecule of the aldehyde is reduced to alcohol while another is oxidised to salt of carboxylic acid.

(ii) Hell-Volhard-Zelinsky reaction: Carbosylic acids having an -hydrogen are halogenated at the -position on treatment with chlorine or bromine in the presence of small amount of red phosphorus to give -halocarboxylic acids.

(b) (i)
$$CH_3CHO$$
 dil NaOH $CH_3 - CH_3 - CH - CH_2 - CHO$ Ethanol $CH_3 - CH - CH_2 - CHO$

(ii) COOH COOH CONH₂

HNO₃ (conc.)

$$H_2SO_4(conc.)$$
;


NO₂
 I_1AIH_4
 I_2AIH_4
 I_1AIH_4
 I_2AIH_4
 I_1AIH_4
 I_1AIH_4

p-Nitro benzyl alcohol

(iii) Benzaldehyde
$$K_2Cr_2O_7/H_2SO_4$$
 SOCl₂ SOCl₂ $AlCl_3$ (anhyd.) Benzophenone

SET-II

- 1. Metallic solid conducts electricity in solid state but ionic solid conducts electricity only in solution or in molten state.
- **2.** The catalytic reaction which depends upon the pore structure of the catalyst and the size of the reaction and product molecules is called shape selective catalysis.
- **4.** H₂S, due to low bond dissociation enthalpy of H–S bond.
- 5. CH₂ CH CH₂OH
- **8.** (i) Thymine is present in DNA.
 - (ii) Uracil is present in RNA.
- **13.** (i) $H_2S_2O_8$ (Peroxodisulphuric acid) (ii) XeF_2



22.

Property	Homogeneous solution	Colloidal solution	Suspension
1. Particle Size	Less than 1nm	Between 1 nm to 1000 nm	more than 1000 nm
2. Separation byordinary filtrationultra filtration	Not possible Not possible	Not possible Possible	Possible Possible
3. Settling of particles	Do not settle	Settle only on configuration	Settle under gravity
4. Appearance	Transparent	Translucent	opaque
5. Example	Glucose dissolved in water	Smoke, milk, Gold sol	Sand in water


$$23. \qquad 3 \text{HgCl}_2 \quad 2 \text{PH}_3 \qquad \text{Hg}_3 \text{P}_2 \quad 6 \text{HCl}$$

$$6 \text{NaOH} + 3 \text{Cl}_2 \qquad 5 \text{NaCl} + \text{NaClO}_3 + 3 \text{H}_2 \text{O}$$

$$(\text{Hot and conc.})$$


$$XeF_4 + O_2 F_2 \quad ^{143 \text{K}} \quad XeF_6 + O_2$$

24. (i) $[Cr(C_2O_4)_3]^{3-}$: Trioxialato chromate (III) ion




Optical isomers of [Cr(Ox) 3]3-

(ii) [Pt(NH₃)₂Cl₂]; Diamine dichlorido platinum (II)



Geometrical isomers of [Pt(NH₃)₂Cl₂]

(iii) [CO(en)₂Cl₂]⁺: Diechloridobis (ethane-1, 2-diamine) cobalt(III) ion



- **26.** (*i*) Sucrose is dextrorotatory, on hydrolysis in the presence of hydrochloric acid or enzyme invertase, it gives a mixture of D-(+)-glucose and D-(-)-fructose which is laeorotatory called invert sugar.
 - (ii) If more than find-amino acids are joined together by peptide bend (CONH) the polyamide thus formed is called polypeptide.
 - (iii) Enzymes are biocatalyst. Almost all the enzymes are globular proteins. Enzymes are specific in nature and efficiency in their action. They are required in small quantity and highly active optimum temperature range (298–310 K) and pH values 5–7.

SET-III

- 1. The substances which are weakly attracted by external magnetic field are called paramagnetic substances and the property. Thus exhibited is called paramagnetism. Paramagnetism is shown by those substances whose atoms, ions or molecules contain unpaired electron contain unpaired electrons, *e.g.*, O², Cu²⁺ Fe³⁺, Cr³⁺, etc.
- **3.** Depressant is used to prevent the formation of froth from one of the two sulphide ores.
- **4.** Since the bond dissociation enthalpy of the H–Cl bond is lower than H–F bond therefore H–Cl is stronger acid than H– F in aqueous solution.
- **5.** 1-methoxy-2-methyl propane.

- **12.** (*i*) This is because fluorine is most electronegative element and does not have d-orbitals in its valence shell.
 - (ii) Due to vary high ionisation enthalpy helium does not form any real chemical compound.
- **14.** (*i*) Dipole moment is a product of charge and distance. Since chlorobenzene has lower magnitude of negative charge on Cl atom and shorter C—Cl bond than cyclohexyl chloride due to resonance therefore chlorobenzene has lower dipole moment than cyclohexyl chloride.

(ii) This is because the new intermolecular attractions between haloalkenes and organic solvent molecules have almost the same strength ones being broken in the separate haloalkan and solvent molecules.

20.
$$T_f T_f^0 T_f = 0$$
°C $- (-0.320$ °C) $= 0.320$ °C $= 0$ or 0.320 K

$$m = 0.0711 \text{ mol kg}^{-1}, K_f = 1.86 \text{ K Kg mol}^{-1}, T_f = 0.320 K$$

Substituting these values in the expression, $T_f = iK_f m$, we get

0.320 K =
$$i$$
 1.86 K kg mol⁻¹ × 0.0711 mol kg⁻¹
 i $\frac{0.320}{0.132}$ = 2.42

- 22. When the dispersion medium is water colloids are classified as hydrophilic colloids and hydrophobic colloids. Hydrophilic colloids are solvent (water) attracting, reversible and quite stable, *e.g.*, gum, starch, gelatin etc. Hydrophobic colloids are solvent (water repelling, irreversible and less stable, *e.g.*, metal sulphides sol, metal hydroxide sol, etc.
- **23.** (i) $\operatorname{Sn} + 2\operatorname{PCl}_5$ heating $\operatorname{SnCl}_4 + 2\operatorname{PCl}_3$
 - (ii) $2Fe_3 + + SO_2 + 2H_2O$ $2Fe^{2+} + SO_4^2$ 4H
 - (iii) $2XeF_2(s) + 2H_2O(e)$ $2Xe + 4HF + O_2$
- 26. The following reactions cannot be explained by the open chain structure of glucose.
 - (i) Despite having the aldehyde group, glucose does not give 2, 4– DNP test, Schiffs test and it does not form the hydrogensulphite addition product with NaHSO₃.
 - (ii) The penta acetate of glucose does not react with hydroxyl amine indicating the absence of free —CHO group.
 - (iii) D-Glucose on treatment of methyl alcohol in the presence of dry HCl gas gives two isomeric nonomethyl derivatives known as -D-glucoside and methyl - D-glucoside. These glucosides do not reduce fehling solution and also do not react with hydrogen cyanide indicating the absence of free —CHO group.

A ring structure called pyranose structure (- or -) is proposed for the glucose molecule.

CBSE Examination Papers Delhi-2013

SET-I

Time allowed: 3 hours Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question numbers 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question numbers 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question numbers 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question numbers 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables, if necessary. Use of calculators is not allowed.
- 1. How many atoms constitute one unit cell of a face-centered cubic crystal?
- 2. Name the method used for the refining of nickel metal.
- 3. What is the covalency of nitrogen in N_2O_5 ?
- **4.** Write the IUPAC name of CH_3 —CH— CH_2 —CH= CH_2 .
- 5. What happens when CH₃—Br is treated with KCN?
- **6.** Write the structure of 3–methyl butanal.
- 7. Arrange the following in increasing order of their basic strength in aqueous solution: CH₃.NH₂, (CH₃)₃N, (CH₃)₂NH
- **8.** What are three types of RNA molecules which perform different functions?
- 9. 18 g of glucose, $C_6H_{12}O_6$ (Molar Mass = 180 g mol⁻¹) is dissolved in 1 kg of water in a sauce pan. At what temperature will this solution boil?

 $(K_b \text{ for water} = 0.52 \text{ K kg mol}^{-1}, \text{ boiling point of pure water} = 373.15 \text{ K})$

- **10.** The conductivity of 0.20 M solution of KCl at 298 K is 0.025 S cm⁻¹. Calculate its molar conductivity.
- 11. Write the dispersed phase and dispersion medium of the following colloidal systems:
 - (i) Smoke (ii) Milk

OR

What are lyophilic and lyophobic colloids? Which of these sols can be easily coagulated on the addition of small amounts of electrolytes?

- 12. Write the differences between physisorption and chemisorption with respect to the following:
 - (i) Specificity

(ii) Temperature dependence

(iii) Reversibility and

- (iv) Enthalpy change
- **13.** (a) Which solution is used for the leaching of silver metal in the presence of air in the metallurgy of silver?
 - (b) Out of C and CO, which is a better reducing agent at the lower temperature range in the blast furnace to extract iron from the oxide ore?
- 14. What happens when
 - (i) PCl₅ is heated?

(ii) H₃PO₃ is heated?

Write the reactions involved.

- **15.** (a) Which metal in the first transition series (3d series) exhibits +1 oxidation state most frequently and why?
 - (b) Which of the following cations are coloured in aqueous solutions and why?

$$Sc^{3+}$$
, V^{3+} , Ti^{4+} , Mn^{2+}

(At. nos.
$$Sc = 21$$
, $V = 23$, $Ti = 22$, $Mn = 25$)

- 16. Chlorobenzene is extremely less reactive towards a nucleophilic substitution reaction. Give two reasons for the same.
- **17.** Explain the mechanism of the following reaction:

$$^{2}\text{CH}_{3}$$
— $^{2}\text{CH}_{2}$ — ^{2}OH 3 4 13 13 14 $^{$

- **18.** How will you convert:
 - (i) Propene to Propan-2-ol?
 - (ii) Phenol to 2, 4, 6 trinitrophenol?
- **19.** (a) What type of semiconductor is obtained when silicon is doped with boron?
 - (b) What type of magnetism is shown in the following alignment of magnetic moments?



- (c) What type of point defect is produced when AgCl is doped with CdCl₂?
- 20. Determine the osmotic pressure of a solution prepared by dissolving 2.5×10^{-2} g of K_2SO_4 in 2 L of water at 25°C, assuming that it is completely dissociated.

$$(R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}, \text{ Molar mass of } \text{K}_2 \text{SO}_4 = 174 \text{ g mol}^{-1}).$$

21. Calculate the emf of the following cell at 298 K:

$$Fe\ (s)\ \big|\ Fe^{2+}\ (0.001\ M)\ \big\|\ H^{+}\ (1M)\ \big|\ H_{2}\ (g)\ (1\ bar),\ Pt\ (s)$$

(Given
$$E_{\text{cell}}^{0} = +0.44 \text{ V}$$
)

- 22. How would you account for the following?
 - (i) Transition metals exhibit variable oxidation states.
 - (ii) Zr (Z = 40) and Hf (Z = 72) have almost identical radii.
 - (iii) Transition metals and their compounds act as catalyst.

OR

Complete the following chemical equations:

(i)
$$Cr_2O_7^{2-}$$
 6Fe² 14H⁺

(ii)
$$2\text{CrO}_4^{2-}$$
 2H

(iii)
$$2MnO_4^ 5C_2O_4^{2-}$$
 16H

23. Write the IUPAC names of the following coordination compounds:

(i)
$$[Cr(NH3)3Cl3]$$

(ii)
$$K_3[Fe (CN)_6]$$

(*iii*)
$$[CoBr_2(en)_2]^+$$
, (en = ethylenediamine)

24. Give the structures of A, B and C in the following reactions:

(i)
$$C_6H_5N_2Cl^-$$
 CuCN A H_2O/H B NH_3 C

(ii)
$$C_6H_5NO_2$$
 Sn HCl A NaNO₂ HCl B H_2O/H C

25. Write the names and structures of the monomers of the following polymers:

(i) Buna-S

(ii) Neoprene

(iii) Nylon-6, 6

26. After watching a programme on TV about the adverse effects of junk food and soft drinks on the health of school children, Sonali, a student of Class XII, discussed the issue with the school principal. Principal immediately instructed the canteen contractor to replace the fast food with the fibre and vitamins rich food like sprouts, salad, fruits, etc. This decision was welcomed by the parents and the students.

After reading the above passage, answer the following questions:

- (a) What values are expressed by Sonali and the Principal of the school?
- (b) Give two examples of water-soluble vitamins.
- **27.** (*a*) Which one of the following is a food preservative? Equanil, Morphine, Sodium benzoate
 - (b) Why is bithional added to soap?
 - (c) Which class of drugs is used in sleeping pills?
- **28.** (a) A reaction is second order in A and first order in B.
 - (i) Write the differential rate equation.
 - (ii) How is the rate affected on increasing the concentration of A three times?
 - (iii) How is the rate affected when the concentrations of both A and B are doubled?
 - (b) A first order reaction takes 40 minutes for 30% decomposition. Calculate $t_{1/2}$ for this reaction. (Given log 1.428 = 0.1548)

OR

- (a) For a first order reaction, show that time required for 99% completion is twice the time required for the completion of 90% of reaction.
- (b) Rate constant 'k' of a reaction varies with temperature 'T' according to the equation:

$$\log k \quad \log A - \frac{E_a}{2.303 R} \quad \frac{1}{T}$$

Where E_a is the activation energy. When a graph is plotted for $\log k$ vs. $\frac{1}{T}$, a straight line with a slope of – 4250 K is obtained. Calculate ' E_a ' for the reaction. ($R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$)

- **29.** (*a*) Give reasons for the following:
 - (i) Bond enthalpy of F₂ is lower than that of Cl₂.
 - (ii) PH₃ has lower boiling point than NH₃.
 - (b) Draw the structures of the following molecules:
 - (i) BrF₃

(ii) (HPO₃)₃

(iii) XeF₄

OR

- (a) Account for the following:
 - (i) Helium is used in diving apparatus.
 - (ii) Fluorine does not exhibit positive oxidation state.
 - (iii) Oxygen shows catenation behaviour less than sulphur.
- (b) Draw the structures of the following molecules:

(i) XeF₂

(ii) H₂S₂O₈

- **30.** (a) Although phenoxide ion has more number of resonating structures than carboxylate ion, carboxylic acid is a stronger acid than phenol. Give two reasons.
 - (b) How will you bring about the following conversions?
 - (i) Propanone to propane
 - (ii) Benzoyl chloride to benzaldehyde
 - (iii) Ethanal to but-2-enal

OR

(a) Complete the following reactions:

- (b) Give simple chemical tests to distinguish between the following pairs of compounds:
 - (i) Ethanal and Propanal
 - (ii) Benzoic acid and Phenol

SET-II

(Questions Uncommon to Set-I)

1. What type of stoichiometric defect is shown by AgCl?

- **4.** What type of bonding helps in stabilising the -helix structure of proteins?
- **6.** What inspired N. Bartlett for carrying out reaction between Xe and PtF₆?
- 7. What happens when ethyl chloride is treated with aqueous KOH?
- **8.** Write the structure of 4–chloropentan–2–one.
- **9.** How will you convert the following?
 - (i) Propan-2-ol to propanone.
- (ii) Phenol to 2, 4, 6-tribromophenol
- 11. What is the difference between oil/water (O/W) type and water/oil (W/O) type emulsions? Give an example of each type.
- **17.** (a) Which of the following ores can be concentrated by froth floatation method and why? Fe₂O₃, ZnS, Al₂O₃.
 - (b) What is the role of silica in the metallurgy of copper?
- **18.** (a) Why does p-dichlorobenzene have a higher m.p. than its o- and m- isomers?
 - (b) Why is () Butan-2-ol is optically inactive?
- **23.** Write the names and structures of the monomers of the following polymers:
 - (i) Polystyrene

(ii) Dacron

- (ii) Teflon
- 27. Write the types of isomerism exhibited by the following complexes:
 - (*i*) [Co(NH₃)₅Cl]SO₄

(ii) $[Co(en)_3]^{3+}$

(ii) [Co(NH₃)₆] [Cr(CN)₆]

SET-III

(Questions Uncommon to Set-I and II)

- 1. What type of substances would make better permanent magnets, ferromagnetic or ferrimagnetic?
- **3.** What is the composition of 'Copper matte'?
- **5.** What is a glycosidic linkage?
- **6.** Write the IUPAC name of (CH₃)₂CH.CH (Cl)CH₃.
- 7. Which compound in the following pair undergoes faster $S_N \mathbf{1}$ reaction?

290 Xam idea Chemistry—XII

- **8.** Write the structure of p-Methylbenzaldehyde molecule.
- **9.** What is the difference between multimolecular and macromolecular colloids? Give one example of each.
- **14.** (a) Give an example of zone refining of metals.
 - (b) What is the role of cryolite in the metallurgy of aluminium?
- **17.** Account for the following:
 - (i) The C—Cl bond length in chlorobenzene is shorter than that in CH₃—Cl.
 - (ii) Chloroform is stored in closed dark brown bottles.
- **18.** How will you convert:
 - (i) Propene to Propan-1-ol?
 - (ii) Ethanal to Propan-2-o1?
- 23. Give the structures of products A, B and C in the following reactions:

- **27.** Write the names and structures of the monomers of the following polymers:
 - (i) Bakelite

(ii) Nylon-6

(iii) Polythene





SET-I

- 1. (8 corner atoms) $\frac{1}{8}$ (6 face centre atoms) $\frac{1}{2} = 1 + 3 = 4$
- 2. Mond Process.
- **3.** 4
- **4.** 4-Chloropent-1-ene.

6.
$$^{4}_{\text{CH}_{3}}$$
 $^{3}_{\text{CH}}$ $^{2}_{\text{CH}_{2}}$ $^{1}_{\text{CHO}}$ $^{1}_{\text{CHO}}$

- 7. $(CH_3)_3 N CH_3 NH_2 (CH_3)_2 NH$
- **8.** There are three types of RNAs:
 - (i) Ribosomal RNA (rRNA)
 - (ii) Messenger RNA (mRNA)
 - (iii) Transfer RNA (tRNA)
- **9.** Moles of glucose $18 \text{ g} / 180 \text{ g} \text{ mol}^{-1} = 0.1 \text{ mol}$

Mass of solvent = 1 kg

Hence, molality of glucose solution = 0.1 mol kg^{-1}

For water, change in boiling point will be

$$T_b = K_b = m$$

= 0.52 K kg mol⁻¹ × 0.1 mol kg⁻¹ = 0.052 K.

Since water boils at 373.15 K at 1.013 bar pressure, the boiling point of solution will be 373.15 + 0.052 = 373.202 K.

10.
$$_{m}$$
 $\frac{k}{c}$ $\frac{1000}{c}$ $\frac{0.025 \text{ S cm}^{-1} \quad 1000 \text{ cm}^{3} \text{ L}^{-1}}{0.20 \text{ mol L}^{-1}}$ $125 \text{ S cm}^{2} \text{ mol}^{-1}$

11.

Type of Colloid	Dispersed Phase	Dispersion Medium
(i) Smoke	Solid	Gas
(ii) Milk	Liquid	Liquid

OR

Lyophilic sols are solvents attracting sols whereas lyophobic sols are solvent repelling sols. Lyophobic sols can be easily coagulated.

12.

	Physisorption	Chemisorption
(i) Specificity	It is not specific in nature.	It is highly specific in nature.
(ii) Temperature dependence	Low temperature is favourable for adsorption. It decreases with increase of temperature.	High temperature is favourable for adsorption. It increases with the increase of temperature.
(iii) Reversibility It is reversible in nature.		It is irreversible.
(<i>iv</i>) Enthalpy change Enthalpy of adsorption is low $(20-40 \text{ kJ mol}^{-1})$ in this case.		Enthalpy of adsorption is high (80-240 kJ mol ¹) in this case.

- **13.** (a) Dilute solution of NaCN.
 - (b) CO.
- **14.** (i) When PCl₅ is heated, the less stable axial bonds break to form PCl₃.

$$PCl_5$$
 Heat $PCl_3 + Cl_2$.

(ii) Orthophosphorous acid on heating disproportionates to give orthophosphoric acid and phosphine.

- 15. (a) Cu has the electronic configuration $3d^{10} 4s^1$. It can easily lose $4s^1$ electron to give the stable $3d^{10}$ configuration. Hence, it shows +1 oxidation state.
 - (b) Mn^2 and V^3 are coloured in aqueous solutions because they have unpaired electrons.
- **16.** (*i*) Due to resonance C—Cl bond acquires a partial double bond character which is difficult to cleave.
 - (ii) Due to sp^2 hybridisation of C of C—Cl bond.
 - (iii) Due to unstable phenyl cation.
 - (iv) Due to repulsion between nucleophile and electron-rich arenes.

Н

17. (i) CH_3 CH_2 O H + H CH_3CH_2 O

$$(iii) \ \mathrm{CH_3-\!CH_2-\!C\!H_3} \quad \longrightarrow \quad \mathrm{CH_3CH_2-\!C\!H_3-\!C\!H_3-\!C\!H_3+\ H^+}$$

18. (i) Propan-2-ol can be prepared from propene by hydration as shown below:

$$\begin{tabular}{lll} ${\rm CH_3-CH}\!=\!{\rm CH_2} & ${\rm H_2O/H^+}$ & ${\rm CH_3-CH}\!-\!{\rm CH_3}$ \\ & & {\rm OH} & \\ \end{tabular}$$

- **19.** (a) p-type semiconductor.
 - (b) Ferromagnetism.
 - (c) Impurity defect/cation vacancy defect.
- **20.** K_2SO_4 dissolved = 0.025 g

Volume of solution = 2 L, T = 25°C = 298 K

Molar mass of $K_2SO_4 = 174$ g mol ¹

Since K_2SO_4 dissociates completely as K_2SO_4 $2K^+ + SO_4^2$ $i = \frac{\text{Total number of moles of particle after dissociation}}{\text{Number of moles of particles before dissociation}} \quad \frac{3}{1} = 3$

Applying van't Hoff equation,

$$= \frac{i \quad W_B \quad R \quad T}{M_B \quad V}$$

$$= \frac{3 \quad 0.025 \text{ g} \quad 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1} \quad 298 \text{ K}}{174 \text{ g mol}^{-1} \quad 2 \text{ L}} = 5.27 \times 10^{-3} \text{ atm}$$

21. Cell reaction:
$$Fe(s) + 2H$$
 (aq) $Fe^2 (aq) + H_2(g)$

 $E_{\text{cell}} \quad E_{\text{cell}}^{\text{o}} \quad \frac{0.0591}{2} \log \frac{[\text{Fe}^{2+}]}{[\text{H}^{+}]^{2}}$ Nernst equation:

$$E_{\text{cell}} = 0.44 = \frac{0.0591}{2} \log \frac{10^{-3}}{(1)^2}$$

$$0.44 \quad \frac{0.0591}{2} \quad (3) \quad 0.44 \quad 0.0886 \quad 0.52856V$$

- 22. (i) Transition elements show variable oxidation states because electrons in ns and (n are available for bond formation as they have nearly same energy.
 - (ii) This is due to filling of 4f orbitals which have poor shielding effect (lanthanoid contraction).
 - (iii) This is because of their ability to adopt multiple oxidation states and to form complexes.

(i)
$$Cr_2O_7^{2-}$$
 6Fe² 14H⁺ 2Cr³⁺ + 6Fe³⁺ + 7H₂O

(ii)
$$2Cr_2O_4^{2-}$$
 2H $Cr_2O_7^{2-}$ H₂O

(i)
$$Cr_2O_7^{2-}$$
 $6Fe^2$ $14H^+$ $2Cr^{3+} + 6Fe^{3+} + 7H_2O$
(ii) $2Cr_2O_4^{2-}$ $2H$ $Cr_2O_7^{2-}$ H_2O
(iii) $2MnO_4^{-}$ $5C_2O_4^{2-}$ $16H$ $2Mn^{2+} + 10CO_2 + 8H_2O$

294 Xam idea Chemistry—XII

23. (i) Triamminetrichloridochromium(III)

(ii) Potassium hexacyanoferrate(III)

(iii) Dibromidobis-(ethane-1,2-diammine)cobalt(III) ion

$$24$$
 (i) A = C₆H₅CN, B = C₆H₅COOH, C = C₆H₅CONH₂

(ii)
$$A = C_6H_5NH_2$$
, $B = C_6H_5N_2^+C1$, $C = C_6H_5OH$

25.

S. No.	Polymers	Monomer names	Monomer structures
(i)	Buna-S	1, 3-Butadiene,	CH ₂ =CH=CH=CH ₂
		Styrene	$C_6H_5CH=CH_2$
			Cl
(ii)	Neoprene	Chloroprene	$CH_2 = C - CH = CH_2$
(iii)	Nylon-6, 6	Hexamethylene diamine,	NH ₂ —(CH ₂) ₆ —NH ₂
		Adipic acid	COOH—(CH ₂) ₄ —COOH

- **26.** (a) Values shown by Sonali: concern for the society, socially active and helpful to others. Values shown by principal: caring, commanding and serious about the welfare of students.
 - (b) Vitamin B complex, Vitamin C.
- **27.** (a) Sodium benzoate.
 - (b) Bithional acts as an antiseptic agent and reduces the odour produced by bacterial decomposition of organic matter on the skin.
 - (c) Tranquilizers.

28. (a) (i) Rate =
$$\frac{d[R]}{dt}$$
 $k[A]^2[B]$

(ii) When concentration of A is tripled

Rate =
$$k [3A]^2[B] = 9k [A]^2[B]$$

i.e.. Rate of reaction will become 9 times.

(ii) When concentration of both A and B is doubled

Rate =
$$k [2A]^2 [2B] = 8k [A]^2 [B]$$

i.e., Rate of reaction will become 8 times.

(b) For a first order reaction

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$
when $t = 40$ minutes, $\frac{[R]_0}{[R]} = \frac{100}{100 - 30} = \frac{10}{7}$

$$k = \frac{2.303}{40} \log \frac{10}{7} = \frac{2.303}{40} \log 1.428$$

$$k = \frac{2.303}{40} = 0.1548$$

$$k = 8.91 \quad 10^{-3} \text{ min}^{-1}$$

$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{8.91 \quad 10^{-3}}$$

$$t_{1/2} = 77.78$$
 min.

OR

(a) As
$$t = \frac{2.303}{k} \log \frac{[R]_0}{[R]}$$

$$\frac{t_{99\%}}{t_{90\%}} = \frac{\frac{2.303}{k} \log \frac{100}{1}}{\frac{2.303}{k} \log \frac{100}{10}} = \frac{\log 100}{\log 10} = 2$$


Hence
$$t_{99\%}$$
 $2t_{90\%}$
 $Slope = \frac{E_a}{2.303 \ R} = 4250 \text{ K}$
 $E_a = 4250 \times 2.303 \times 8.314$
 $= 81375 \text{ J mol}^{-1}$
 $= 81.375 \text{ kJ mol}^{-1}$

- 29. (a) (i) Bond dissociation enthalpy decreases as the bond distance increases from F₂ to I₂ because of the corresponding increase in the size of the atom as we move from F to I. The F—F bond dissociation enthalpy is, however, smaller than that of Cl—Cl and even smaller than that of Br—Br. This is because F atom is very small and hence the three lone pairs of electrons on each F atom repel the bond pair holding the F-atoms in F₂ molecule. Hence, the bond dissociation enthalpy increases in the order: I₂ F₂ Br₂ Cl₂.
 - (ii) Unlike NH₃, PH₃ molecules are not associated through hydrogen bonding in liquid state. That is why the boiling point of PH₃ is lower than NH₃.




Slightly bent "T"

(ii) (HPO₃)₃




(iii) XeF₄



Square planar

- (a) (i) Helium is used as a diluent for oxygen in modern diving apparatus because of its very low solubility in blood.
 - (ii) This is because fluorine is the most electronegative element.
 - (iii) Because O—O single bond is weaker than S—S single bond.
- (b) (i) Refer to Q. 12(ii) of CBSE (All India) Paper 2013.



- **30.** (a) (i) Resonating structures of carboxylate ion are more stable than phenoxide ion.
 - (ii) Negative charge is dispersing on two electronegative oxygens in carboxylate ion whereas it is on one oxygen atom in phenoxide ion.
 - (b) (i) Propanone to Propane

$$\begin{array}{c|cccc} O & & & & & \\ CH_3-C-CH_3 & & & & & \\ Propanone & & & & & \\ & & & & & \\ Propanone & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & \\ & & & \\ &$$

(ii) Benzoyl chloride to benzaldehyde

(iii) Ethanal to but-2-enal

2CH₃CHO Dil. NaOH (Aldol condensation)
4
CH₃ 3 CHOH 2 CHO 2 CHO 3 -Hydroxybutanal 3 CH 2 CHO 3 -Hydroxybutanal 3 CH 2 CH 3 CH 3 CH 2 CH 3 CH 2 CH 3 CH 2 CH 3 CH 2 CHO But - 2 - enal 3 CH 2 CHO

(a) (i)
$$2H$$
—C—H $\xrightarrow{\text{conc. KOH}}$ H_3C —OH + H—C $\overset{O}{\circ}$ OK

(ii) CH_3COOH $\xrightarrow{Br_2/P}$ CH_2 —COOH

(iii) CHO
$$\frac{\text{HNO}_3/\text{H}_2\text{SO}_4}{273-283 \text{ K}}$$

$$\text{NO}_2$$

- (b) (i) Ethanal (CH₃CHO) and propanal (CH₃CH₂CHO): Ethanal responds to iodoform test and gives a yellow precipitate on addition of NaOH and I₂, while propanal does not.
 - (ii) **Phenol and benzoic acid:** Benzoic acid reacts with NaHCO₃ giving CO₂ gas with effervescence, whereas phenol does not Phenol decolourises Br₂ water giving white precipitate, whereas benzoic acid does not.

SET-II

(Questions Uncommon to Set-I)

- 1. Frenkel defect.
- **2.** 4-Bromo-4-methylpent-2-ene.
- **4.** The -helix structure of proteins is stabilised by intramolecular H-bonding between C=O of one amino acid residue and the N—H of the fourth amino acid residue in the chain.
- **6.** Neil Bartlett observed that PtF_6 reacts with O_2 to yield an ionic solid, O_2^+ PtF_6

$$O_2(g)$$
 $PtF_6(g)$ O_2^+ $[PtF_6]$

Here, O_2 gets oxidised to O_2^+ by PtF_6 .

Since the first ionisation enthalpy of Xe (1170 kJ mol 1) is fairly close to that of O_{2} molecule (1175 kJ mol 1), Bartlett thought that PtF_{6} should also oxidise Xe to Xe $^{+}$. This inspired Bartlett to carry out the reaction between Xe and PtF_{6} . When Xe and PtF_{6} were mixed, a rapid reaction occurred and a red solid with the formula, $Xe^{+}[PtF_{6}]^{-}$ was obtained.

OH OH OH SO₃H HNO₃ (conc.)
$$O_2N$$
 NO₂

Phenol SO₃H NO₂

Phenol-2,4 disulphonic acid (Picric acid)

- 11. (i) Oil in water (o/w) type emulsions: In such emulsions oil is the dispersed phase and water is the dispersion medium, e.g., milk, vanishing cream.
 - (ii) Water in oil (w/o) type emulsion: In such emulsions, water is the dispersed phase and oil is the dispersion medium, e.g., butter, cod liver oil.
- **17.** (a) ZnS. This is because the sulphide ore particles are preferentially wetted by oil and gangue particles are preferentially wetted by water.
 - (b) Silica acts as a flux to remove iron oxide (FeO) as slag in the metallurgy of copper.

$$\begin{array}{ccc} \text{FeO} + \text{SiO}_2 & \text{FeSiO}_3 \\ & \text{Flux} & \text{Slag} \end{array}$$

- **18.** (a) It is due to symmetry of *p*-dichlorobenzene which fits in crystal lattice better than the *o* and *m*-isomers.
 - (b) (±)-Butan-2-ol exist in two eventiomeric forms which rotate the plane of polarised light in opposite directions in equal amounts. Therefore, cancelling each other and becoming optically inactive.

23.

Polymer	Monomers and their structures	
(i) Polystyrene	CH=CH ₂	
	Styrene (Vinyl benzene)	
(ii) Dacron	OHCH ₂ —CH ₂ OH Ethyl glycol	HOOC—COOH Terephthalic acid
(ii) Teflon	$ \begin{array}{c} F \\ F \end{array} $ $ \begin{array}{c} C = C \\ F \end{array} $ Tetrafluoroethene	

- **27.** (*i*) $[Co(NH_3)_5Cl]SO_4$ Ionisation isomerism (*ii*) $[Co(en)_3]^{3+}$ Optical isomerism
 - (ii) $[Co(NH_3)_6]$ $[Cr(CN)_6]$ Coordination isomerism

SET-III

(Questions Uncommon to Set-I and II)

- 1. Ferromagnetic.
- 3. Cu₂S and FeS.
- 5. The linkage between two monosaccharide units through oxygen atom is called glycosidic linkage.
- **6.** 2-Chloro-3-methylbutane.
- 7. CI

9. Multimolecular colloids: In this type of colloids, colloidal particles are aggregates of large atoms or molecules each having size in the colloidal range, *e.g.*, sulphur sol, gold sol. Multimolecular colloids are generally lyophobic in nature.

Macromolecular colloids: These are the solution containing macromolecules in the colloidal range, *e.g.*, starch, proteins, polyethene, etc. Macromolecular colloids are generally lyophilic in nature.

- **14.** (*a*) Zone refining is used for production of semiconductors and other metals of very high purity like germanium, silicon, boron, gallium and indium.
 - (b) Cryolite lowers the melting point of the mixture./Acts as an electrolyte./Acts as solvent for alumina. (Any one)
- 17. (i) Refer to Q.23(iii) of CBSE Examination Paper (All India) 2013.
 - (ii) Because chloroform is slowly oxidised by air in the presence of light to an extremely poisonous gas phosgene.

18. (i) Propene to Propan-1-ol

(ii) Ethanal to Propan-2-o1

$$CH_3 \longrightarrow C = O \xrightarrow{CH_3MgI} CH_3 \longrightarrow CCH_3 \xrightarrow{H^+, H_2O} CH_3 \longrightarrow CCH_3 \longrightarrow CCH_3$$

$$Ethanal \qquad Ethanal \qquad E$$

- 23. (i) $A = CH_3CH_2CN$, $B = CH_3CH_2CH_2NH_2$, $C = CH_3CH_2CH_2OH$
 - (ii) A = CH₃CONH₂, B = CH₃NH₂, C = CH₃NC

27. OH
$$\bigcirc$$
 , HCHO \bigcirc , Phenol Formaldehyde

(iii) Polythene:
$$CH_2 = CH_2$$
 Ethene

CBSE Examination Papers All India-2013

SET-I

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper, Delhi-2013.

- 1. Of physisorption or chemisorption, which has a higher enthalpy of adsorption?
- 2. Name the method used for refining of copper metal.
- 3. Name two poisonous gases which can be prepared from chlorine gas.
- **4.** Write the IUPAC name of the following compound:

$$\begin{array}{c|c}
CH_3 \\
CH_3 - CH - CH - CH_3 \\
CH_3 - CI
\end{array}$$

5. Rearrange the following compounds in the increasing order of their boiling points:

$$CH_3$$
— CH_0 , CH_3 — CH_2 — OH , CH_3 — CH_2 — CH_3

- **6.** Write the structure of n-methylethanamine.
- 7. What are the products of hydrolysis of sucrose?
- 8. Is $-(CH_2_CH)$ n a homopolymer or a copolymer?
- **9.** Account for the following:
 - (i) Schottky defects lower the density of related solids.
 - (ii) Conductivity of silicon increases on doping it with phosphorus.
- **10.** Aluminium crystallises in an fcc structure. Atomic radius of the metal is 125 pm. What is the length of the side of the unit cell of the metal?
- 11. The standard electrode potential (E°) for Daniell cell is + 1.1 V. Calculate the G° for the reaction

$$Zn(s)$$
 Cu^{2} (aq) Zn^{2} (aq) $Cu(s)$ $(1 \text{ F} = 96500 \text{ C mol}^{-1}).$

12. (a) For a reaction A + B P, the rate law is given by,

$$r k[A]^{1/2}[B]^2$$
.

What is the order of this reaction?

- (b) A first order reaction is found to have a rate constant $k = 5.5 \times 10^{-14} \text{ s}^{-1}$. Find the half life of the reaction.
- **13.** (a) Name the method used for removing gangue from sulphide ores.
 - (b) How is wrought iron different from steel?
- **14.** Draw the structures of the following molecules:
 - (i) XeOF₄

- (ii) H₃PO₃
- 15. How are interhalogen compounds formed? What general compositions can be assigned to them?
- **16.** Explain the mechanism of the following reaction:

$$_{443 \text{ K}}^{\text{H}}$$
 CH_2 = $\text{CH}_2 + \text{H}_2\text{O}$.

- **17.** Write the equations involved in the following reactions:
 - (i) Reimer-Tiemann reaction
 - (ii) Williamson's ether synthesis
- **18.** Define thermoplastic and thermosetting polymers. Give one example of each.

OR

What is a biodegradable polymer? Give an example of a biodegradable aliphatic polyester.

19. The rate of a reaction becomes four times when the temperature changes from 293 K to 313 K. Calculate the energy of activation (E_a) of the reaction assuming that it does not change with temperature.

$$[R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}, \log 4 = 0.6021]$$

- **20.** What are the characteristics of the following colloids? Give one example of each:
 - (i) Multimolecular colloids
 - (ii) Lyophobic sols
 - (iii) Emulsions
- **21.** Give reasons for the following:
 - (i) Where R is an alkyl group, $R_3P = 0$ exists but $R_3N = 0$ does not.
 - (ii) PbCl₄ is more covalent than PbCl₂
 - (iii) At room temperature, N2 is much less reactive.
- 22. For the complex $[NiCl_4]^{2-}$, write
 - (i) the IUPAC name.
 - (ii) the hybridisation type.
 - (iii) the shape of the complex.

(Atomic no. of
$$Ni = 28$$
)

OR

What is meant by crystal field splitting energy? On the basis of crystal field theory, write the electronic configuration of d^4 in terms of t_{2g} and e_g in an octahedral field when

$$(i)$$
 $_0 > P$

(ii)
$$0 < P$$

302 Xam idea Chemistry—XII

- 23. Give reasons for the following:
 - (i) Ethyl iodide undergoes S_N2 reaction faster than ethyl bromide.
 - (ii) () 2-Butanol is optically inactive.
 - (iii) C-X bond length in halobenzene is smaller than C-X bond length in CH₃-X.
- **24.** Complete the following reactions:
 - (i) $CH_3CH_2NH_2 + CHCl_3 + alc. KOH$
 - $\begin{array}{ccc} (\it{ii}) \ \ C_6 H_5 N_2 \ Cl^- & & \ \ \ & \ \ & \ \ & \ \ & \ \ & \ \ \ & \ \ & \ \ & \ \ & \ \ & \ \ & \ \ & \ \ &$

$$(iii) \begin{picture}(100,0) \put(0,0){\line(1,0){100}} \put(0,0){\line(1$$

- **25.** (i) What class of drug is Ranitidine?
 - (ii) If water contains dissolved Ca²⁺ ions, out of soaps and synthetic detergents, which will you use for cleaning clothes?
 - (iii) Which of the following is an antiseptic?
 - 0.2% phenol, 1% phenol.
- **26.** Calculate the emf of the following cell at 25°C:

$$Ag(s)/Ag (10^{-3} M) | Cu^2 (10^{-1} M) Cu (s)$$

Given $E_{\text{cell}}^{\text{o}} = +0.46 \text{ V}$ and $\log 10^n = n$.

27. Shanti, a domestic helper of Mrs. Anuradha, fainted while mopping the floor. Mrs. Anuradha immediately took her to the nearby hospital where she was diagnosed to be severely 'anaemic'. The doctor prescribed an iron rich diet and multivitamins supplement to her. Mrs. Anuradha supported her financially to get the medicines. After a month, Shanti was diagnosed to be normal.

After reading the above passage, answer the following questions:

- (i) What values are displayed by Mrs. Anuradha?
- (ii) Name the vitamin whose deficiency causes 'pernicious anaemia'.
- (iii) Give an example of a water soluble vitamin.
- **28.** (a) State Raoult's law for a solution containing volatile components.

How does Raoult's law become a special case of Henry's law?

(b) 1.00 g of a non-electrolyte solute dissolved in 50 g of benzene lowered the freezing point of benzene by 0.40 K. Find the molar mass of the solute. (K_f for benzene = 5.12 kg mol⁻¹)

OR

- (a) Define the following terms:
 - (i) Ideal solution
 - (ii) Azeotrope
 - (iii) Osmotic pressure
- (b) A solution of glucose ($C_6H_{12}O_6$) in water is labelled as 10% by weight. What would be the molality of the solution?

(Molar mass of glucose = 180 g mol^{-1})

- **29.** (a) Give reasons for the following:
 - (i) Mn³⁺ is a good oxidising agent
 - (ii) $E_{\text{M}^2/\text{M}}^{\text{o}}$ values are not regular for first row transition metals (3d series).
 - (iii) Although 'F' is more electronegative than 'O', the highest Mn fluoride is MnF_4 , whereas the highest oxide is Mn_2O_7 .
 - (b) Complete the following equations:
 - (i) $2\text{CrO}_4^{2-} + 2\text{H}^+$
 - (ii) KMnO₄ heat

OR

- (a) Why do transition elements show variable oxidation states?
 - (i) Name the element showing maximum number of oxidation states among the first series of transition metals from Sc (Z = 21) to Zn (Z = 30).
 - (ii) Name the element which shows only +3 oxidation state.
- (b) What is lanthanoid contraction? Name an important alloy which contains some of the lanthanoid metals.
- **30.** (a) How will you convert the following:
 - (i) Propanone to Propan-2-ol
 - (ii) Ethanal to 2-hydroxy propanoic acid
 - (iii) Toluene to benzoic acid
 - (b) Give simple chemical test to distinguish between:
 - (i) Pentan-2-one and Pentan-3-one
 - (ii) Ethanal and Propanal

OR

(a) Write the products of the following reactions:

(i)
$$CH_3$$
— C — CH_3 CH_3 — CH_3 CH_3 — CH_3 CH_4 CH_5 CH_5 CH_5 CH_5 CH_6 CH_6 CH_6 CH_6 CH_7 CH_7 CH_7 CH_8 CH_8

- (b) Which acid of each pair shown here would you expect to be stronger?
 - (i) F— CH_2 —COOH or Cl— CH_2 —COOH

SET-II

(Questions Uncommon to Set-I)

- 1. Write the structure of 2-aminotoluene.
- 2. Which aerosol depletes ozone layer?
- 4. Ethanol is soluble in water. Why?
- 5. Write the IUPAC name of the following compound:

$$\begin{array}{c|cccc} \mathrm{CH}_3\mathrm{--CH}\mathrm{--CH}_2\mathrm{--CH}\mathrm{--CH}_3\\ & | & | \\ \mathrm{Br} & \mathrm{Cl} \end{array}$$

- 7. Write the name of linkage joining two amino acids.
- **8.** Give one example of a condensation polymer.
- **9.** (a) Why does presence of excess of lithium makes LiCl crystals pink?
 - (b) A solid with cubic crystal is made of two elements P and Q. Atoms of Q are at the corners of the cube and P at the body-centre. What is the formula of the compound?
- 14. Draw the structures of the following molecules:
 - (i) XeF₆

- (ii) H₂S₂O₇
- **18.** Outline the principles of refining of metals by the following methods:
 - (i) Zone refining

- (ii) Vapour phase refining
- **19.** Define the following terms giving an example of each:
 - (i) Associated colloids

(ii) Lyophilic sol

- (iii) Adsorption
- **22.** Write the main products of the following reactions:

(i)
$$C_6H_5N_2CI^ H_3PO_2 + H_2O$$

 NH_2

$$(ii) \qquad \xrightarrow{\operatorname{Br}_2(aq)} ?$$

- **27.** Give reasons for the following:
 - (i) Oxygen is a gas but sulphur is a solid.
 - (ii) O₃ acts as a powerful oxidising agent.
 - (iii) BiH₃ is the strongest reducing agent amongst all the hydrides of Group 15 elements.

SET-III

(Questions Uncommon to Set-I and II)

- 1. What is especially observed when a beam of light is passed through a colloidal solution?
- **2.** What is the basicity of H_3PO_3 and why?
- 3. Write the IUPAC name of the following compound:

- **5.** What are the products of hydrolysis of lactose?
- **6.** Is $(CH_2 CH)_n$ a homopolymer or a copolymer?

- **8.** Write the structure of prop-2-en-1-amine.
- **12.** Draw the structures of the following molecules:
 - $(i) N_2 O_5$

- (ii) XeF₂
- **13.** (a) What change occurs when AgCl is doped with CdCl₂?
 - (b) What type of semiconductor is produced when silicon is doped with boron?
- **18.** Name the principal ore of aluminium. Explain the significance of leaching in the extraction of aluminium.
- 19. Define the following terms with an example in each case:
 - (i) Macromolecular sol

(ii) Peptisation

- (iii) Emulsion
- **21.** Give reasons for the following:
 - (i) Though nitrogen exhibits +5 oxidation state, it does not form pentahalide.
 - (ii) Electron gain enthalpy with negative sign of fluorine is less than that of chlorine.
 - (iii) The two oxygen-oxygen bond lengths in ozone molecule are identical.
- 27. Write the main products of the following reactions:
 - (i) $CH_3CH_2NH_2$ $\frac{HNO_2}{0.C}$

(iii)
$$N-H$$
 CH_3-C-C_1 $Base$?



SET-I

- 1. Chemisorption.
- 2. Electrolytic refining.
- **3.** The poisonous gases which can be prepared from chlorine gas are phosgene gas (COCl₂)/mustard gas (CICH₂CH₂CH₂CH₂CH₂Cl)/chloropicrin or tear gas (CCl₃NO₂)/sulphur monochloride (S₂Cl₂).
- **4.** 2-Chloro-3,3-dimethylbutane or 3-Chloro-2,2-dimethylbutane.
- 5. $CH_3CH_2CH_3 < CH_3CHO < CH_3CH_2OH$.
- **6.** CH₃—NH—CH₂CH₃.
- 7. Sucrose on hydrolysis gives one molecule each of glucose and fructose.
- 8. Homopolymer.
- **9.** (*i*) In Schottky defect as the number of ions are missing from their normal lattice sites, the mass decreases whereas the volume remains the same. Due to this the density decreases.
 - (ii) This is due to availability of additional unpaired electrons on doping with phosphorous.

10. For an fcc unit cell,
$$r = \frac{a}{2\sqrt{2}}$$

 $a = 2\sqrt{2}r = 2 = 1.414 = 125 = 353.5 \text{ pm}.$
11. ${}_{r}G^{o} = nFE^{o}_{\text{(cell)}}$

In the given equation, n=2, F=96500 C mol $^{-1}$ and $E_{\text{cell}}^{\text{o}}=1.1$ V


Therefore,

212300 J mol
$$^{1} = -212.3 \text{ kJ mol}^{-1}$$

- 12. (a) Order of reaction $\frac{1}{2}$ 2 $\frac{5}{2}$
 - (b) Radioactive decay follows first order kinetics.

$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{5.5 \cdot 10^{-14}} \text{ s} = 1.26 \cdot 10^{13} \text{ s}.$$

- **13.** (a) Froth floatation.
 - (b) Wrought iron is the purest form of iron whereas steel is an alloy of iron.



- **15.** Interhalogen compounds are prepared by direct combustion or by the action of halogen on lower interhalogen compounds. They can be assigned general composition as XX, XX₃, XX₅ and XX₇ where X is halogen of larger size and X of smaller size and X is more electropositive than X.
- **16.** The mechanism of dehydration of ethanol involves the following steps:

Step 1: Formation of protonated alcohol:

Step 2: Formation of carbocation: It is the slowest step and hence, the rate determining step of the reaction.

Step 3: Formation of ethane by elimination of a proton

The acid used in step 1 is released in step 3. To drive the equilibrium to the right, ethene is removed as it is formed.

17. (i) Reimer-Tiemann reaction:

(ii) Williamson synthesis:

18. Thermoplastic polymers: These polymers are capable of repeatedly softening on heating and hardening on cooling. Example, polythene, polystyrene, etc.

Thermosetting polymers: These polymers on heating undergo extensive cross linking in moulds and again become infusible. These cannot be reused. Example, bakelite, urea-formaldehyde resins, etc.

Polymers which disintegrate by themselves over a period of time due to environmental degradation by microorganisms are called biodegradable polymers. Example, PHBV (Poly- -hydroxybutyrate-co- -hydroxyvalerate).

19. If rate at 293 K is *R*, rate at 313 K will be 4*R*.

$$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \frac{T_2 - T_1}{T_1 - T_2}$$

$$\log \frac{4R}{R} = \frac{E_a}{2.303R} \frac{313 - 293}{8.314}$$

$$\log 4 = \frac{E_a}{19.1471} \frac{20}{91709}$$

$$E_a = \frac{0.6021}{20} \frac{19.1471}{20} \frac{91709}{20}$$

 $= 52863.2177 \text{ J mol}^{-1} \text{ or } 52.863 \text{ kJ mol}^{-1}$

- **20.** (*i*) Multimolecular colloids are aggregates of atoms or molecules each having size less than 1 nm held together by van der Waals forces. Example, sulphur sol, gold sol.
 - (ii) Lyophobic sols are those sols in which the particles of the disperse phase have little affinity for the particles of the dispersion medium. Example, sols of metal and their sulphides and hydroxides, As₂S₃ sol, Fe(OH)₃ sol.
 - (iii) Emulsion is a colloidal solution in which both the dispersed phase and dispersion medium are liquids. Example, milk, cod liver oil, etc.
- 21. (i) N due to the absence of d-orbitals, cannot form p d multiple bonds. Thus, N cannot expand its covalency beyond four but in R₃N=O, N has a covalency of 5. So, the compound R₃N=O does not exist. On the other hand, P due to the presence of d-orbitals forms p d multiple bonds and hence can expand its covalency beyond 4. Therefore, P forms R₃P=O in which the covalency of P is 5.
 - (ii) Because Pb is in +4 oxidation state in PbCl₄ and has high charge/size ratio than Pb²⁺, thus polarising power of Pb⁴⁺ is greater than Pb²⁺, and hence it is more covalent.
 - (iii) N_2 is less reactive at room temperature because of strong p-p overlap resulting into the triple bond (N N), consequently high bond dissociation enthalpy.
- 22. (i) Tetrachloridonickelate(II) ion.
 - (ii) sp^3 hybridisation.
 - (iii) Tetrahedral.

OR

The difference of energy between two sets of degenerate orbitals as a result of crystal field splitting is known as crystal field splitting energy.

- (i) $t_{2g}^4 e_g^0$
- (ii) $t_{2g}^{3} e_{g}^{1}$

- 23. (i) Since I ion is a better leaving group than Br ion, hence, CH₃I reacts faster than CH₃Br in S_N2 reaction with OH⁻ ion.
 - (ii) () 2-Butanol is a racemic mixture, *i.e.*, there are two enantiomers in equal proportions. The rotation by one enantiomer will be cancelled by the rotation due to the other isomer, making the mixture optically inactive.
 - (iii) In CH₃–X the carbon atom is sp^2 hybridised while in halobenzene the carbon atom is sp^3 hybridised. The sp^2 hybridised carbon is more electronegative due to greater s-character and holds the electron pair of C–X bond more tightly than sp^3 hybridised carbon with less s-character. Thus, C–X bond length in CH₃–X is bigger than C–X in halobenzene.
- **24.** (*i*) $CH_3CH_2NH_2 + CHCl_3 + 3KOH(alc.)$ $CH_3CH_2NC + 3KCl + 3H_2O$

$$NH_2$$
 $+ HCl(aq)$
 $+ HCl(aq)$

- **25.** (i) Antacid/antihistamine.
 - (ii) Synthetic detergents
 - (*iii*) 0.2% phenol.
- **26.** The given cell notation in the question is incorrect.

The correct cell formula is given below:

$$Cu^{2+}(10^{-1} \text{ M}) | Cu(s) | | Ag^{+}(10^{-3} \text{ M}) | Ag(s)$$

According to Nernst equation

$$E_{\text{cell}} = E_{\text{cell}}^{\text{o}} - \frac{0.0591}{2} \log \frac{[\text{Cu}^2]}{[\text{Ag}]^2}$$

Given: $E_{\text{cell}}^{\text{o}} = 0.46 \text{ V}$

$$E_{\text{cell}} = 0.46 - \frac{0.0591}{2} \log \frac{[0.1]}{[10^{-3}]^2}$$

$$= 0.46 - 0.02955 \log \frac{[0.1]}{[10^{-6}]}$$

$$= 0.46 - 0.02955 \log 10^5$$

$$= 0.46 - 0.02955 \quad 5$$

$$= 0.46 - 0.148 = 0.314 V.$$
Or

$$E_{\text{cell}} = E_{\text{cell}}^{\text{o}} - \frac{0.059}{2} \log \frac{[\text{Ag}]^2}{[\text{Cu}^2]}$$
$$= 0.46 - \frac{0.0591}{2} \log \frac{[10^{-3}]^2}{[0.1]}$$

=
$$0.46 - 0.02955 \log \frac{[10^{-6}]}{[0.1]}$$

= $0.46 + 0.0295$ 5
= 0.6075 V.

- 27. (i) Helpfulness, caring, generosity, kindness attitude towards poor.
 - (ii) Vitamin B₁₂.
 - (iii) Vitamin C.
- **28.** (a) For a solution of volatile liquids, Raoult's law states that the partial vapour pressure of each component of the solution is directly proportional to its mole fraction present in solution, *i.e.*,

$$p_A \quad x_A$$
, or $p_A \quad p_A^{o} \quad x_A$

According to Henry's law, the partial pressure of a gas in vapour phase (p) is directly proportional to mole fraction (x) of the gas in the solution.

$$p = K_H x$$

On comparing it with Raoult's law it can be seen that partial pressure of the volalite component or gas is directly proportional to its mole fraction in solution.

$$p \lambda$$

only the proportionality constant K_H differs from p_A^o . Thus, it becomes a special case of Henry's law in which $K_H = p_A^o$.

(b) Substituting the values of various terms involved in equation $M_B = \frac{K_f - W_B - 1000}{T_f - W_A}$, we get

$$M_B = \frac{5.12 \, \, \mathrm{K \, kg \, mol^{-1}} \, \, \, \, 1.00 \, \, \mathrm{g} \, \, \, \, 1000 \, \, \mathrm{g \, kg^{-1}}}{0.40 \, \, \mathrm{K} \, \, \, \, \, 50 \, \, \mathrm{g}}$$
 256 g mol ⁻¹

OR

- (a) (i) Ideal solution: The solution which follows Raoult's law at all concentrations and temperatures.
 - (ii) Azeotrope: A liquid mixture which distills at constant temperature without undergoing any change in composition.
 - (iii) Osmotic pressure: The minimum excess pressure that has to be applied on the solution side when the solution and solvent are separated by a semipermeable membrane to stop osmosis.
- (b) 10% w/w solution of glucose means 10 grams of glucose is present in 100 g of solution, i.e., 90 g of water.

$$m = \frac{1000 \text{ wt\%}}{(100 - \text{wt\%})} = \frac{1000 \text{ 10}}{\text{Molecular weight of solute}} = \frac{1000 \text{ 10}}{(100 \text{ 10})} = 0.617 \text{ m}$$

- **29.** (a) (i) Mn^3 (3d⁴) is a good electron acceptor as the resulting species is more stable (3d⁵).
 - (ii) The E (M² / M) values are not regular which can be explained from the irregular variation of ionisation enthalpies ($_{i}H_{1}$ $_{i}H_{2}$) and also the sublimation enthalpies which are relatively much less for manganese and vanadium.
 - (iii) Due to multiple bond formation ability of oxygen with Mn in Mn₂O₇.

(b) (i)
$$2\text{CrO}_4^{2-} + 2\text{H}^+$$
 $\text{Cr}_2\text{O}_7^{2-} + \text{H}_2\text{O}$
(ii) 2KMnO_4 Heat $\text{K}_2\text{MnO}_4 + \text{MnO}_2 + \text{O}_2$

- (a) Transition elements show variable oxidation states because electrons in ns and (n 1) d-orbitals are available for bond formation as they have nearly same energy.
 - (i) Manganese (ii) Scandium.
- (b) The steady decrease in the atomic and ionic radii (having the same charge) with increase in atomic number as we move across the series from lanthanum to lutetium is known as lanthanoid contraction.

Misch metal.

- (i) Pentan-2-one and pentan-3-one: Pentan-2-one responds to iodoform test and gives yellow coloured precipitate with I2 and NaOH, while pentan-3-one does not.
 - (ii) Ethanal and propanal: Ethanal responds to iodoform test and gives yellow coloured precipitate with I₂ and NaOH, while propanal does not.

- (b) (i) F— CH_2 —COOH > Cl— CH_2 —COOH, because of higher -ve I effect of F.
 - OH $(ii) \ \, \text{CH}_3\text{COOH} > \ \, \ \, \ \,$, because CH_3COO^- is a more stable conjugate base.


SET-II

- 2. CCl₂F₂/Freon.
- **4.** Alcohols can form H-bonds with water and break the H-bonds already existing between water molecules. Hence, they are soluble in water.
- **5.** 2-Bromo-4-chloropentane.
- 7. Peptide linkage.
- 8. Nylon-6,6/Dacron/Glyptal.
- **9.** (a) Excess of lithium leads to metal excess defect. Lithium atoms lose electrons to form Li⁺ ions. These electrons diffuse into the crystal and form F-centres. Therefore, LiCl crystals become pink.
 - (b) Number of P atoms per unit cell = 1 (at the body centre) 1 = 1

Number of Q atoms per unit cell = 8 (at the corners) $\frac{1}{8}$

Hence, the formula of the compound PQ.

14. (*i*) XeF₆



(ii) H₂S₂O₇

- **18.** (*i*) **Zone refining:** Zone refining is based on the principle that the impurities are more soluble in the melt than in the solid state of metal.
 - (ii) Vapour phase refining: In this, metal is converted into its volatile compound and collected elsewhere. It is then decomposed to give pure metal.
- **19.** (*i*) Associated colloids: There are certain substances which at low concentrations behave as normal strong electrolytes, but at higher concentrations exhibit colloidal behaviour due to the formation of aggregates. Such colloids are known as associated colloids or micelles, *e.g.*, soaps and detergents.
 - (ii) Lyophilic sol: Lyophilic sols are those sols in which the particles of dispersed phase have great affinity for the dispersion medium, e.g., sols of gum, gelatin, starch, etc.
 - (iii) Adsorption: The accumulation of molecular species at the surface rather than in the bulk of a solid or liquid is termed adsorption, *e.g.*, water vapours on silica gel, poisonous gases on charcoal.
- **22.** (*i*) $C_6H_5N_2Cl^ H_3PO_2 + H_2O$ $C_6H_6 + N_2 + H_3PO_3 + HCl$

$$(ii) \begin{picture}(100,0) \put(0,0){\line(1,0){100}} \put(0,0){\line(1,$$

(iii)
$$CH_3$$
— C — NH_2 Br_2 $4NaOH$ CH_3 — NH_2 + Na_2CO_3 + $2NaBr$ + $2H_2O$

- 27. (i) Oxygen forms p p multiple bonds. Due to small size and high electronegativity oxygen exists as diatomic (O_2) molecules. Whereas, sulphur because of its bigger size and lower electronegativity, has a much greater tendency for catenation than oxygen and lower tendency for p p multiple bonds.
 - (ii) Due to the ease with which it liberates atoms of nascent oxygen, it acts as a powerful oxidising agent.
 - (iii) This is because Bi—H bond is the weakest among the hydrides of group 15.

SET-III

- 1. Tyndall effect/Illumination of the path of light.
- 2. As two P—OH bonds are present in the molecule, therefore, its basicity is two.
- **3.** 2, 5-Dichlorotoluene/1,4-Dichloro-2-methylbenzene.
- **5.** Glucose and galactose.
- **6.** Homopolymer.

12. (i) N_2O_5

(ii) XeF₂



Linear

- **13.** (a) A cationic vacancy is generated.
 - (b) p-type semiconductor.
- **18.** Bauxite ($Al_2O_3.2H_2O$).

Leaching is significant as it helps in removing the impurities like SiO₂, Fe₂O₃, TiO₂ etc. from the bauxite ore and thus alumina is obtained.

19. (*i*) **Macromolecular sol:** Macromolecules in suitable solvents form solutions in which the size of the macromolecules may be in colloidal range, *e.g.*, starch, cellulose, nylon, etc.

- (ii) **Peptisation:** The process of conversion of a freshly prepared precipitate into a colloidal solution by shaking it with dispersion medium in the presence of a suitable electrolyte is called peptisation. Example, freshly prepared precipitate of Fe(OH)₃ is peptised by FeCl₃.
- (iii) Emulsion: These are the colloidal systems in which both the dispersion medium and the dispersion phase are liquid, e.g., milk is an emulsion of fat in water.
- **21.** (*i*) Nitrogen does not have *d*-orbitals in its valence shell to expand its covalence beyond four. That is why it does not form pentahalide.
 - (ii) Lower value of bond dissociation enthalpy of F_2 is due to the strong repulsion between the non-bonding electrons of F atoms in the small sized F_2 molecule.
 - (iii) Because of resonance.

27. (i)
$$CH_3CH_2NH_2$$
 $HNO_2 \ OC$ CH_3CH_2OH

$$(iii) \begin{array}{c} N-H \\ \hline \\ H \end{array} \begin{array}{c} O \\ \hline \\ CH_3-C-Cl \\ \hline \\ Base \end{array} \begin{array}{c} O \\ \hline \\ N-C-CH_3 \\ \hline \\ H \end{array}$$



CBSE Examination Papers Foreign-2013

SET-I

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper, Delhi-2013.

- 1. Which group of solids is electrical conductors, as well as malleables and ductiles?
- 2. Measurement of which colligative property is preferred for determination of molar mass of biomolecules?
- 3. Name the method used for refining of germanium metal.
- **4.** Write the IUPAC name of the following compound:

- **5.** Arrange the following compounds in the increasing order of their acid strengths: 4-nitrophenol, phenol, 2,4,6-trinitrophenol
- **6.** Write the structure of 3-hydroxybutanal.
- 7. Why is methylamine more basic than aniline?
- **8.** Write the names of monomers of the polymer


$$-$$
HNH $(CH_2)_6$ NH $\stackrel{C}{\underset{O}{\text{NH}}}$ $(CH_2)_4$ $\stackrel{C}{\underset{O}{\text{Cl}}}$ $\stackrel{C}{\underset{O}{\text{NH}}}$

- 9. Henry's law constant ($_{\rm H}$) for the solution of methane in benzene at 298 K is 4.27 $\times 10^5$ mm Hg. Calculate the solubility of methane in benzene at 298 K under 760 mm Hg.
- 10. (i) Why is an increase in temperature observed on mixing chloroform and acetone?
 - (ii) Why does sodium chloride solution freeze at a lower temperature than water?
- 11. The rate constant for a first order reaction is 60 s^{-1} . How much time will it take to reduce the initial concentration of the reactant to $\frac{1}{10}$ th of its initial value?
- **12.** (a) What happens when a freshly precipitated Fe(OH)₃ is shaken with water containing a small quantity of FeCl₃?
 - (b) Why is a finely divided substance more effective as an adsorbent?
- 13. Write two differences between lyophobic and lyophilic sols. Give one example of each type of sol.
- **14.** How can you separate alumina (Al₂O₃) from silica present in bauxite ore? Write the chemical equations for the reactions involved.

OR

- (a) What are the constituents of 'copper matte'?
- (b) What is the role of depressant in froth floatation process?
- 15. Arrange the following in the order of property indicated for each set:
 - (a) HF, HCl, HBr, Hl increasing acid strength.
 - (b) NH₃, PH₃, AsH₃, SbH₃, BiH₃ increasing reducing power.
- **16.** Write the reactions involved when *D*-glucose is treated with the following reagents:
 - (a) HCN

- (b) Br₂-water
- 17. (a) What type of bonding helps in stabilising the -helix structure of proteins?
 - (b) What is the structural difference between a nucleoside and a nucleotide?
- 18. Arrange the following polymers in increasing order of their intermolecular forces:
 - (a) Nylon-6,6, Buna-S, Polythene
 - (b) PVC, Nylon-6, Neoprene
- 19. An element with molar mass 27 g mol⁻¹ forms a cubic unit cell with edge length 4.05×10^{-8} cm. If its density is 2.7 g cm⁻³, what is the nature of the cubic unit cell?
- **20.** For a chemical reaction R P, the variation in the concentration ln[R] vs. time t plot is given as



For this reaction

- (i) what is the order of the reaction?
- (ii) what is the slope of the curve?
- (iii) what is the unit of rate constant 'k'?
- 21. Draw the structures of the following molecules:
 - $(i) N_2O_5$

(ii) H₃PO₂

- (iii) XeF₆
- **22.** Account for the following:
 - (i) White phosphorus is more reactive than red phosphorus.
 - (ii) SnCl₄ is more covalent than SnCl₂.
 - (iii) O₃ is a powerful oxidising agent.

OR


Complete the following equations:

(i) PC1₃ H₂O

(ii) XeF₂ PF₅

(iii) NaN₃ heat

- **23.** (a) Write the IUPAC name of the complex $[CoBr_2(en)_2]^+$.
 - (b) What type of isomerism is shown by the complex [Co(NH₃)₅ SO₄]Br?
 - (c) Why is CO a stronger ligand than NH₃ in complexes?
- **24.** Consider the three types of replacement of group X by group Y as shown here.



This can result in giving compound (A) or (B) or both. What is the process called if

- (i) (A) is the only compound obtained?
- (ii) (B) is the only compound obtained?
- (iii) (A) and (B) are formed in equal proportions?
- **25.** (a) Explain the mechanism of the following reaction:

$$^{2}\text{CH}_{3}\text{CH}_{2}\text{OH}$$
 $^{\text{H}^{+}}$ $^{+}$ $^{-}$ $^{-}$ CH $_{3}\text{CH}_{2}$ —O—CH $_{2}\text{CH}_{3}$ + H $_{2}\text{O}$

- (b) Name the reagent used in the oxidation of ethanol to ethanoic acid.
- **26.** How will you convert the following:
 - (i) Aniline to chlorobenzene
 - (ii) Ethanoic acid to methanamine
 - (iii) Benzene diazonium chloride to phenol
- 27. Mr. Naresh works in a multi-national company. He is stressed due to his hectic schedule. Mr. Amit, his friend, comes to know that he has started taking sleeping pills without consulting the doctor. Mr. Amit requests Naresh to stop this practice and takes him to a Yoga centre. With regular Yoga sessions, Mr. Naresh is now a happy and relaxed man.

After reading the above passage, answer the following questions:

- (i) Write the values shown by Mr. Amit.
- (ii) Which class of drugs is used in sleeping pills?
- (iii) Why is it not advisable to take sleeping pills without consultation with the doctor?
- **28.** (a) State Kohlrausch's law of independent migration of ions. Mention one application of Kohlrausch's law.
 - (b) The resistance of a conductivity cell containing 10^{-3} M KCl solution at 25°C is 1500 . What is the cell constant if conductivity of 10^{-3} M KCl solution at 25°C is 1.5×10^{-4} S cm⁻¹?

OR

Calculate emf and G° for the following cell at 298 K:

$$Mg(s) \mid Mg^{2+}(10^{-3} M) \mid \mid Cu^{2+}(10^{-4} M) \mid Cu(s)$$

Given
$$E_{\text{Mg}^{2+}/\text{Mg}}^{\text{o}} = -2.36 \text{ V}$$
 and $E_{\text{Cu}^{2+}/\text{Cu}}^{\text{o}} = +0.34 \text{ V}$

$$(1 \text{ F} = 96500 \text{ C mol}^{-1})$$

318 Xam idea Chemistry—XII

- **29.** Assign reasons for the following:
 - (i) In the series Sc (Z = 21) to Zn (Z = 30), the enthalpy of atomisation of Zn is the lowest.
 - (ii) Zr and Hf have almost identical radii.
 - (iii) Transition metals show variable oxidation states.
 - (iv) The $E^{o}_{M^2/M}$ value for copper is positive (+ 0.34 V).
 - (v) Cr^{2+} is a very good reducing agent.

OR

Describe the preparation of KMnO₄ from pyrolusite ore (MnO₂). How does the acidified permanganate solution react with the following:

- (i) Fe^{2+} ions
- (ii) Oxalic acid (C₂O₄H₂)

Write the ionic equations for the reactions involved.

- **30.** (a) Give reasons for the following:
 - (i) Ethanal is more reactive than acetone towards nucleophilic addition reaction.
 - (ii) (CH₃)₃C—CHO does not undergo aldol condensation.
 - (iii) Carboxylic acids are higher boiling liquids than alcohols.
 - (b) Give a simple chemical test to distinguish between
 - (i) Acetophenone and Benzophenone
 - (ii) Benzaldehyde and Ethanal

OR

Write the structures of products of the following reactions:

$$(i) \begin{array}{c} O \\ C-CI \\ + H_2 \end{array} \xrightarrow{Pd/BaSO_4} ?$$

$$(ii) CH_3-C-H \qquad HCN \qquad ?$$

$$0 \qquad (iii) 2H-C-H \qquad Conc. KOH \qquad ?$$

$$0 \qquad (iv) \begin{array}{c} O \\ H_2NCONHNH_2 \\ \hline (v) \end{array} \begin{array}{c} O \\ \hline NaOH, CaO \end{array} ?$$

SET-II

Questions Uncommon to Set-I

- 1. Why is glass considered a supercooled liquid?
- **2.** Ethylamine is soluble in water whereas aniline is almost insoluble. Why?
- **4.** Write the structure of 4-methyl pent-3-en-2-one.

6. Write the IUPAC name of the following compound:

$$CH_2 = CH - CH_2 - CH - CH_2 - CI$$

$$CH_3$$
 CH₃
7. On what principle is chromatography based?

- **8.** Write the names of monomers of the following polymer:

$$+OCH_2-CH_2-O-C$$
 O
 O
 O

- 9. What type of deviation is shown by a mixture of ethanol and acetone? What type of azeotrope is formed by mixing ethanol and acetone?
- **14.** Write the reactions involved when *D*-glucose is treated with the following reagents:

- (ii) H₂N—OH
- **16.** What is special about the following terms:
 - (i) Kraft Temperature

- (ii) Sorption
- 22. Draw the structures of the following molecules:
 - (i) PCl₅

(ii) $H_4P_2O_7$

- (iii) ClF₃
- **25.** (a) Write the formulae for the following coordination compounds:
 - (i) Tetraammineaquachloridocobalt (III) chloride
 - (ii) Potassiumtetracyanonickelate (II)
 - (b) Write the hybridisation of the complex $[NiCl_4]^{2-}$. (Atomic number of Ni = 28)
- 27. How are the following conversions carried out:
 - (i) Aniline to fluorobenzene
 - (ii) Benzene diazonium chloride to benzene
 - (iii) Methyl chloride to ethylamine

SET-III

Questions Uncommon to Set-I and II

- 1. What is the two-dimensional coordination number of a molecule in a square close-packed layer?
- 2. Write the IUPAC name of the following compound:

3. Write the names of monomers of the following polymer:

$$\begin{array}{c|c} - C - (CH_2)_5 - N \\ \parallel & \parallel \\ O & H \end{array}$$

4. Why are diazonium salts of aromatic amines more stable than those of aliphatic amines?

320 Xam idea Chemistry—XII

- 7. Write the structural formula of 2-phenyl ethanoic acid?
- **8.** Name the method used for the refining of Titanium metal.
- **9.** Out of two 0.1 molal solutions of glucose and of potassium chloride, which one will have a higher boiling point and why?
- 14. Write the structural difference between DNA and RNA.
- **16.** Write the dispersed phase and dispersion medium of the following colloids:
 - (i) Cheese

(ii) Fog

21. (a) What type of isomerism is shown by each of the following complexes:

(*i*) [Pt(NH₃)₂ Cl₂]

(ii) [Co(NH₃)₅(NO₂)]Cl₂

- (b) Give an example of the role of coordination compounds in biological systems.
- **23.** Draw the structures of the following:
 - (i) Solid PCl₅

(ii) H₂S₂O₈

(iii) XeO3

- **26.** How are the following conversions carried out:
 - (i) Aniline to iodobenzene
 - (ii) Ethyl nitrile to ethyl amide
 - (iii) Benzene diazonium chloride to benzonitrile



2



- 1. Metallic solids.
- 2. Osmotic pressure.
- 3. Zone refining method.
- **4.** 2-Chloro-3-methylbutane.
- **5.** phenol < 4-nitrophenol < 2,4,6-trinitrophenol

OH O

- 6. H₃C CH CH₂ C H
- 7. Methylamine is more basic than aniline due to resonance in aniline and lower stability of aniline ions.
- **8.** Hexamethylenediamine $[H_2N (CH_2)_6 NH_2]$ and adipic acid $[HOOC (CH_2)_4 COOH]$.
- **9.** Here, $_{\rm H}$ 4.27 $10^5 \, \text{mm Hg}$

p 760 mm Hg

According to Henry's law, $p = \frac{H^{x} CH_{4}}{760 \text{ mm Hg}} = 1.78 - 10^{-3}$ $x_{CH_{4}} = \frac{p}{H} = \frac{760 \text{ mm Hg}}{4.27 - 10^{5} \text{ mm Hg}} = 1.78 - 10^{-3}$

Mole fraction of methane in benzene; $x_{\text{CH}_4} = 1.78 \times 10^{-3}$.

10. (*i*) The bonds between chloroform molecules and molecules of acetone are dipole—dipole interactions but on mixing, the chloroform and acetone molecules, they start forming hydrogen bonds which are stronger bonds resulting in the release of energy. This gives rise to an increase in temperature.

(ii) When a non-volatile solute is dissolved in a solvent, the vapour pressure decreases. As a result, the solvent freezes at a lower temperature.

11.
$$t = \frac{2.303}{k} \log \frac{[R]_0}{[R]} \quad \text{as} \quad [R] = \frac{[R]_0}{10}$$
$$t = \frac{2.303}{k} \log \frac{[R]_0}{\frac{[R]_0}{10}} = \frac{2.303}{60} = \log 10 = \frac{2.303}{60} = 1 \quad \textbf{3.838} \quad \textbf{10} \quad ^2\textbf{s}$$

12. (a) It is converted into colloidal state.

$$Fe(OH)_3 + FeCl_3$$
 $[Fe(OH)_3Fe]^{3+} + 3Cl^{-}$

- (b) Powdered substances have greater surface area as compared to their crystalline forms. Greater the surface area, greater is the adsorption.
- 13. (i) Lyophobic sols: Particles of dispersed phase have no affinity for dispersion medium, rather they hate dispersion medium. They are not easily prepared and need stabilizing agents for their preservation. They are irreversible. Examples: sols of gold, silver, Fe(OH)₃, As₂O₃ etc. They are also called extrinsic colloids.

- (ii) Lyophilic sols: Particles of dispersed phase have great affinity for the dispersion medium. They are self-stabilized because of strong attractive forces operating between the suspended particles and the dispersion medium. They are reversible in nature. Examples: gums, gelatin, starch, albumin, etc. They are also known as intrinsic colloids.
- **14.** Leaching of aluminium from bauxite: Finely powdered bauxite ore is digested with an aqueous solution of sodium hydroxide at 473–523 K and 35–36 bar pressure. Al₂O₃ is leached out as sodium aluminate (and SiO₂ too as sodium silicate) leaving impurities behind.

$$Al_2O_3(s)$$
 $2NaOH(aq)$ $3H_2O$ $2Na[Al(OH)_4](aq)$

The aluminate in solution is neutralised by passing CO_2 gas and hydrated Al_2O_3 is precipitated. At this stage, the solution is seeded with freshly prepared samples of hydrated Al_2O_3 which induces the precipitation.

2Na [Al(OH)₄](
$$aq$$
) CO₂(g) Al₂O₃. x H₂O 2NaHCO₃(aq)

The sodium silicate remains in the solution and hydrated alumina is filtered, dried and heated to get back pure Al_2O_3 .

$$Al_2O_3.xH_2O(s)$$
 $^{1470 \text{ K}}$ $Al_2O_3(s)$ $xH_2O(g)$ **OR**

- (a) Cu₂S and FeS.
- (b) In froth floatation process, the role of the depressant is to prevent certain type of particles from forming the froth with the air bubbles.
- **15.** (a) HF < HCl < HBr < HI.
 - (b) $NH_3 < PH_3 < AsH_3 < SbH_3 < BiH_3$.
- **16.** (a) HCN


$$\begin{array}{cccc} \text{CHO} & & \text{CH} \swarrow \text{CN} \\ | & & | & \text{OH} \\ (\text{CHOH})_4 & & \text{HCN} & (\text{CHOH})_4 \\ | & & | & | \\ \text{CH}_2\text{OH} & & \text{CH}_2\text{OH} \\ \end{array}$$

D-Glucose Glucose cyanohydrin

(b) Br₂-water

- **17.** (a) Hydrogen bonds (intermolecular) between the C=O of one amino acid residue and N—H of the fourth amino acid residue in the chain give stability to the structure.
 - (b) A nucleoside is formed of pyrimidine or purine base connected to C-1 of sugar (ribose or deoxyribose) by a -linkage.

A nucleotide contains all the three basic components of nucleic acids, i.e., a phosphoric acid group, a pentose sugar and a nitrogenous base.



- **18.** (*a*) Buna-S < Polythene < Nylon 6,6.
 - (b) Neoprene < PVC < Nylon-6.

19. Density,
$$d = \frac{Z - M}{a^3 - N_A}$$
 or $Z = \frac{d - a^3 - N_A}{M}$...(i)

Here, M 27 g mol 1

$$a = 4.05 = 10^{-8}$$
 cm

$$d = 2.7 \text{ g cm}^{-3}$$


$$N_A$$
 6.022 10^{23} mol ¹

Substituting these values in expression (i), we get
$$Z = \frac{2.7 \text{ g cm}^{-3} + (4.05 - 10^{-8} \text{ cm})^3 - 6.022 - 10^{23} \text{ mol}^{-1}}{27 \text{ g mol}^{-1}}$$

$$Z = 3.99 = 4$$

Thus, there are 4 atoms of the element present per unit cell. Hence, the cubic unit cell must be face-centred.

- **20.** For this reaction
 - (i) First order
 - (ii) k (rate constant)
 - (iii) time⁻¹ (s^{-1})



(ii)
$$H_3PO_2$$

O

P

OH

324 Xam idea Chemistry—XII

- 22. (i) This is due to polymeric structure of red phosphorus or angular strain in P₄ molecule of white phosphorus where the angle is only 60°.
 - (ii) The oxidation states of central atom Sn in SnCl₄ and SnCl₂ are +4 and +2 respectively. +4 state of Sn has higher polarising power which, inturn, increase the covalent character of bond formed between the central atom and the other atoms.
 - (iii) Due to the ease with which it liberates atoms of nascent oxygen, it acts as a powerful oxidising agent.

$$O_3$$
 O_2 + O (nascent oxygen)

OR

(i)
$$PCl_3 + 3H_2O H_3PO_3 + 3HCl$$

$$(ii)$$
 XeF₂ PF₅ [XeF] [PF₆]

(iii)
$$2NaN_3$$
 heat $2Na + 3N_2$

- 23. (a) Dibromobis (ethane-1, 2-diamine) cobalt
 - (b) Ionisation isomerism.
 - (c) Because in case of CO back bonding takes place in which the central metal uses its filled d orbital with empty anti-bonding *molecular orbital of CO.
- **24.** (*i*) Retention
 - (ii) Inversion
 - (iii) Racemisation

Н

25. (a)
$$CH_3$$
 CH_2 $OH+H$ CH_3-CH_2 OH

- (b) Tollens' reagent.
- **26.** (*i*) Aniline to chlorobenzene

(ii) Ethanoic acid to methanamine

(iii) Benzene diazonium chloride to phenol

- **27.** (*i*) Caring nature, helpful.
 - (ii) Tranquilizers.
 - (iii) Most of the drugs taken in doses higher than recommended may cause harmful effects and act as poison. Therefore, a doctor should always be consulted before taking the medicine.
- 28. (a) Kohlrausch's Law: It states that the limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the electrolyte. In general, if an electrolyte on dissociation gives cations and _ anions then its limiting molar conductivity is given by

Here, o and o are the limiting molar conductivities of cations and anions, respectively.

Applications of Kohlrausch's Law:

Calculation of molar conductivities of weak electrolyte at infinite dilution: For example, molar conductivity of acetic acid (weak acid) at infinite dilution can be obtained from the knowledge of molar conductivities at infinite dilution of strong electrolytes like HCl, CH₃COONa and NaCl as illustrated below.

$$i.e., \qquad {\overset{\text{o}}{m}}_{\text{(CH}_3\text{COOH)}} \qquad {\overset{\text{o}}{\text{CH}_3\text{COO}}} \qquad {\overset{\text{o}}{\text{H}}}^+ \qquad {\overset{\text{o}}{\text{Cl}}}^- \] \qquad {\overset{\text{o}}{\text{Na}}} \qquad {\overset{\text{o}}{\text{Cl}}}^- \]$$

(b) Conductivity () = $\frac{1}{\text{Resistance }(R)} \times \text{Cell constant}$

1.5
$$10^{-4}$$
 S cm $^{-1}$, $R = 1500$ ohm
1.5 $10^{-4} = \frac{1}{1500}$ Cell constant
Cell constant 1.5 $10^{-4} = 1500 = 0.225$ cm $^{-1}$

Cell reaction:
$$Mg + Cu^2$$
 $Mg^2 + Cu$; $n = 2$

Nernst equation: E_{cell} E_{cell}^0 $\frac{0.0591}{2} \log \frac{[Mg^{2+}]}{[Cu^{2+}]}$

$$E_{cell} = 0.34 \quad (2.36) \quad \frac{0.0591}{2} \log \frac{10^{-3}}{10^{-4}}$$

$$= 2.70 - 0.02655 = 2.67 \text{ V}$$

$$G = -nFE_{cell}^0$$

$$G = -2 \times 96500 \times 2.71$$

$$= -5.23 \times 10^5 \text{ J mol}^{-1}$$

$$= -5.23 \times 10^2 \text{ kJ mol}^{-1}$$

- **29.** (*i*) In the formation of metallic bonds, no electrons from 3*d*-orbitals are involved in case of zinc, while in all other metals of the 3*d* series, electrons from the *d*-orbitals are always involved in the formation of metallic bonds. That is why, the enthalpy of atomisation of zinc is the lowest in the series.
 - (ii) This is due to filling of 4f orbitals which have poor shielding effect (lanthanoid contraction).
 - (iii) Transition elements show variable oxidation states because electrons in ns and (n-1) d-orbitals are available for bond formation as they have nearly same energy.
 - (iv) This is because the sum of enthalpies of sublimation and ionisation is not balanced by hydration enthalpy.
 - (v) Cr^{2+} is a stronger reducing agent because after the loss of one electron Cr^{2+} becomes Cr^{3+} which has more stable t_{2g}^3 (half-filled) configuration in a medium like water.

It is prepared by fusion of pyrolusite, MnO₂, with KOH in the presence of an oxidising agent like KNO₃. This produces the dark green potassium manganate, K₂MnO₄ which disproportionates in a neutral or acidic solution to give purple permanganate.

$$2MnO_2 + 4KOH + O_2$$
 $K_2MnO_4 + 2H_2O$ $3MnO_4^{2-} + 4H^+$ $2MnO_4^- + MnO_2 + 2H_2O$

Commercially, it is prepared by alkaline oxidative fusion of MnO₂ followed by the electrolytic oxidation of manganate (VI).

$$\begin{array}{ccc} MnO_2 & Fused \ with \ KOH & MnO_4^{2-} \\ & oxidised \ with \ air \ or \ KNO_3 & Manganate \ ion \\ MnO_4^{2-} & Electrolytic \ oxidation \\ & in \ alkaline \ solution & MnO_4^{-} \\ & Permanganate \end{array}$$

In the laboratory, KMnO₄ is prepared by oxidation of manganese (II) ion salt by peroxodisulphate.

$$2Mn^{2+} + 5S_2O_8^{2-} + 8H_2O$$
 $2MnO_4^- + 10SO_4^{2-} + 16H^+$
Peroxodisulphate Permanganate

(i)
$$MnO_4 + 8H + 5Fe^2$$
 $Mn^{2+} + 4H_2O + 5Fe^{3+}$

(ii)
$$2\text{MnO}_4 + 16\text{H} + 5$$
 $2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 10\text{CO}_2$ 2COO^-

- **30.** (a) (i) This is due to steric and electronic reasons. Sterically, the presence of two relatively large substituents in ketones hinders the approach of nucleophile to carbonyl carbon than in aldehydes having only one such substituent. Electronically two alkyl groups reduce the positivity of the carbonyl carbon more effectively in ketones than in aldehydes.
 - (ii) This is because for aldol condensation to take place, at least one -hydrogen (i.e., hydrogen at carbon adjacent to carbonyl carbon) should be available, which is not present in (CH₃)₃C—CHO.
 - (iii) This is due to more extensive association of carboxylic acid molecules through intermolecular hydrogen bonding. The hydrogen bonds do not break completely even in the vapour phase.
 - (b) (i) Acetophenone and benzophenone:

Acetophenone responds to iodoform test and gives a yellow precipitate on addition of NaOH and I_2 , but benzophenone does not.

(ii) Benzaldehyde and Ethanal

Ethanal reacts with NaOI (I₂/NaOH) to form yellow precipitate of iodoform while benzaldehyde does not give this test.

$$(i) \bigcirc \begin{matrix} O \\ C - Cl \\ + H_2 \end{matrix} \xrightarrow{Pd/BaSO_4} CHO$$

(ii)
$$CH_3$$
— C — H HCN CH_3 — C — CN H

Acetaldehyde cyanohydrin

OH

SET-II

(Questions Uncommon to Set-I)

- 1. Glass is an amorphous solid and has a tendency to flow, though very slowly.
- 2. Due to hydrogen bonding ability of ethylamine.

4.
$$CH_3$$
 C $CH=CH$ CH_3 O CH_3

- **6.** 5-Chloro-4-methyl pent-1-ene.
- 7. Is is based on the principle that different components of a mixture are differently adsorbed on an adsorbent.
- 8. Ethylene glycol and terephthalic acid
- **9.** Positive deviation.


Minimum boiling azeotropes.

- **16.** (*i*) Kraft temperature is the minimum temperature above which the formation of micelles takes place.
 - (ii) Sorption is the process in which adsorption and absorption take place simultaneously, e.g., dyeing of cotton fibres by azo dyes.


22. (*i*) PCl₅



(ii) H₄P₂O₇



(iii) ClF₃



- **25.** (a) (i) [Co(NH₃)₄(H₂O)Cl]Cl₂ (ii) K₂[Ni(CN)₄]
 - (b) sp^3 hybridisation.
- 27. (i) Aniline to fluorobenzene

(ii) Benzene diazonium chloride to benzene

$$N_2$$
CI $^-$
 $+ H_3$ PO $_2 + H_2$ O $+ N_2 + H_3$ PO $_3 + H$ C
Benzene diazonium chloride

(iii) Methyl chloride to ethylamine

SET-III

- **1.** 4.
- 2. 1-chloro 2,4-dinitrobenzene.
- 3. Caprolactam.
- **4.** The diazonium salts of aromatic amines are more stable than those of aliphatic amines due to dispersal of the positive charge on the benzene ring.

- **8.** Vapour phase refining.
- 9. 0.1 M KCl solution will have higher boiling point as KCl dissociates in the solution.
- **14.** Structural difference between DNA and RNA:

DNA	RNA	
1. The sugar present in DNA is 2-deoxy <i>D</i> -(–) ribose.	1. The sugar present in RNA is <i>D</i> -(–)-ribose.	
2. DNA has double stranded -helix structure.	2. RNA has single -helix structure.	

330 Xam idea Chemistry—XII


16.


	Dispersed phase	Dispersion medium	
(i) Cheese	Liquid	Solid	
(ii) Fog	Liquid	Gas	

- **21.** (a) (i) Geometrical isomerism
 - (ii) Linkage isomerism.
 - (b) Vitamin B₁₂ (Cyanocobalamine), the antipernicious anaemia factor, is a complex of cobalt.


23. (*i*) Solid PCl₅

In solid state PCl_5 exist as $[PCl_4]^+$ $[PCl_6]^-$ in which the cation, $[PCl_4]^+$ is tetrahedral and the anion, $[PCl_6]^-$ is octahedral.






(ii) H₂S₂O₈







26. (i) Aniline to iodobenzene

$$\begin{array}{c|c} NH_2 & \stackrel{\uparrow}{V_2Cl} \\ \hline & NaNO_2/HCl \\ \hline & 273-278 \text{ K} \end{array} \qquad \begin{array}{c} \stackrel{\uparrow}{V_2Cl} \\ \hline & I \\ \hline & Iodobenzene \\ \end{array}$$

(ii) Ethyl nitrile to ethyl amide

$$\text{CH}_{3}\text{—CN} \xrightarrow{\text{H}_{2}\text{O/H}^{+}} \text{CH}_{3}\text{—COOH} \xrightarrow{\text{NH}_{3},} \text{CH}_{3}\text{CONH}_{2}$$

(iii) Benzene diazonium chloride to benzonitrile

$$\begin{array}{c|c}
\bar{N}_2\bar{C}l & CN \\
\hline
& KCN, \\
CuCN
\end{array}$$



Time allowed: 3 hours Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question numbers 1 to 8 are very short answer questions and carry 1 mark each.
- (iii) Question numbers 9 to 18 are short answer questions and carry 2 marks each.
- (iv) Question numbers 19 to 27 are also short answer questions and carry 3 marks each.
- (v) Question numbers 28 to 30 are long answer questions and carry 5 marks each.
- (vi) Use log tables, if necessary. Use of calculators is not allowed.

SET-I

- 1. Give one example each of 'oil in water' and 'water in oil' emulsion.
- 2. Which reducing agent is employed to get copper from the leached low grade copper ore?
- 3. Which of the following is more stable complex and why? $[Co(NH_3)_6]^{3+}$ and $[Co(en)_3]^{3+}$
- **4.** Write the IUPAC name of compound.

- **5.** Which of the following isomers is more volatile: o-nitrophenol or p-nitrophenol?
- **6.** What are isotonic solutions?
- **7.** Arrange the following compounds in increasing order of solubility in water:

- **8.** Which of the two components of starch is water soluble?
- 9. An element with density $11.2 \,\mathrm{g} \,\mathrm{cm}^{-3}$ forms a f.c.c. lattice with edge length of 4×10^{-8} cm. Calculate the atomic mass of the element.

(Given:
$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$
)

10. Examine the given defective crystal

A^{+}	B^-	A^{+}	B^{-}	A^{+}
B^{-}	0	B^{-}	A^{+}	B^{-}
A^{+}	B^-	A^{+}	0	A^{+}
B^{-}	A^{+}	\mathbf{B}^{-}	A^{+}	B^{-}


Answer the following questions:

- (i) What type of stoichiometric defect is shown by the crystal?
- (ii) How is the density of the crystal affected by this defect?
- (iii) What type of ionic substances show such defect?
- 11. Calculate the mass of compound (molar mass = 256 g mol⁻¹) to be dissolved in 75 g of benzene to lower its freezing point by 0.48 K ($K_f = 5.12 \text{ K kg mol}^{-1}$).
- 12. Define an ideal solution and write one of its characteristics.
- 13. Write two differences between 'order of reaction' and 'molecularity of reaction'.
- 14. Outline the principles behind the refining of metals by the following methods:
 - (i) Zone refining method
 - (ii) Chromatographic method
- 15. Complete the following chemical equations:
 - (i) Ca₃P₂ + H₂O
 - (ii) Cu + H_2SO_4 (conc.)

OR

Arrange the following in the order of property indicated against each set:

- (i) HF, HCl, HBr, HI increasing bond dissociation enthalpy
- (ii) H₂O₂H₂S₂, H₂Se₂, H₂Te increasing acidic character
- **16.** Write the IUPAC name of the complex $[Cr(NH_3)_4Cl_2]^+$. What type of isomerism does it exhibit?
- 17. (i) Which alkyl halide from the following pair is chiral and undergoes faster S_N^2 reaction?



- (ii) Out of S_N1 and S_N2, which reaction occurs with
 - (a) Inversion of configuration
- (b) Racemisation
- **18.** Draw the structure of major monohalo product in each of the following reactions:

(i)
$$\sim$$
 OH $\stackrel{SOCl_2}{\longrightarrow}$

(ii)
$$\sim$$
 CH₂ - CH = CH₂ + HBr \sim Peroxide \rightarrow

- **19.** (*a*) In reference to Freundlich adsorption isotherm write the expression for adsorption of gases on solids in the form of an equation.
 - (b) Write an important characteristic of lyophilic sols.
 - (c) Based on type of particles of dispersed phase, give one example each of associated colloid and multimolecular colloid.

- **20.** (a) Draw the structures of the following molecules:
 - (i) XeOF₄
 - (ii) H₂SO₄
 - (b) Write the structural difference between white phosphorus and red phosphorus.
- **21.** Account for the following:
 - (i) PCl₅ is more covalent than PCl₃.
 - (ii) Iron on reaction with HCl forms FeCl₂ and not FeCl₃.
 - (iii) The two O—O bond lengths in the ozone molecule are equal.
- 22. The following data were obtained during the first order thermal decomposition of SO₂Cl₂ at a constant volume:

$$SO_2Cl_2$$
 $SO_2(g) + Cl_2(g)$

Experiment	Time/second ⁻¹	Total pressure/atm
1	0	0.4
2	100	0.7

Calculate the rate constant.

(Given: $\log 4 = 0.6021$, $\log 2 = 0.3010$)

- 23. (i) Give two examples of macromolecules that are chosen as drug targets.
 - (ii) What are antiseptics? Give an example.
 - (iii) Why is use of aspartame limited to cold foods and soft drinks?
- **24.** (i) Deficiency of which vitamin causes night-blindness?
 - (ii) Name the base that is found in nucleotide of RNA only.
 - (iii) Glucose on reaction with HI gives n-hexane. What does it suggest about the structure of glucose?
- 25. After the ban on plastic bags, students of one school decided to make the people aware of the harmful effects of plastic bags on environment and Yamuna River. To make the awareness more impactful, they organised rally by joining hands with other schools and distributed paper bags to vegetable vendors, shopkeepers and departmental stores. All students pledged not to use polythene bags in future to save Yamuna River.

After reading the above passage, answer the following questions:

- (i) What values are shown by the students?
- (ii) What are biodegradable polymers? Give one example.
- (iii) Is polythene a condensation or an addition polymer?
- **26.** (a) Write the mechanism of the following reaction:

$$CH_3CH_2OH$$
 $CH_3CH_2Br + H_2O$

- (b) Write the equations involved in Reimer-Tiemann reaction.
- 27. Give the structures of A, B and C in the following reactions:

(i)
$$CH_3Br$$
 KCN A LiA/H_4 B HNO_2 C $273 K$ (ii) CH_3COOH NH $_3$ A Br_2 KOH B $CHCl_3$ NaOH C

How will you convert the following:

- (i) Nitrobenzene into aniline
- (ii) Ethanoic acid into methanamine
- (iii) Aniline into N-phenylethanamide (Write the chemical equations involved.)
- **28.** (a) Define the following terms:
 - (i) Limiting molar conductivity (ii) Fuel cell
 - (b) Resistance of a conductivity cell filled with 0.1 mol L^{-1} KCl solution is 100 . If the resistance of the same cell when filled with 0.02 mol L^{-1} KCl solution is 520 , calculate the conductivity and molar conductivity of 0.02 mol L^{-1} KCl solution. The conductivity of 0.1 mol L^{-1} KCl solution is 1.29×10^{-2} $^{-1}$ cm $^{-1}$.

OR

- (a) State Faraday's first law of electrolysis. How much charge in terms of Faraday is required for the reduction of 1 mol of Cu²⁺ to Cu?
- (b) Calculate emf of the following cell at 298 K:

$$Mg(s) \bigg| \hspace{0.1cm} Mg^{\hspace{0.1cm} 2} \hspace{0.3cm} (0.1M) \hspace{0.1cm} \bigg| \hspace{0.1cm} Cu^{\hspace{0.1cm} 2} \hspace{0.3cm} (0.01) \hspace{0.1cm} \bigg| \hspace{0.1cm} Cu(s)$$

[Given $E_{\text{cell}}^{\text{o}}$ 2.71 V, 1 F 96500 C mol⁻¹]

- **29.** (*a*) How do you prepare:
 - (i) K_2MnO_4 from MnO_2 ?
- (ii) Na $_{2}$ Cr $_{2}$ O $_{7}$ from Na $_{2}$ CrO $_{4}$?
- (b) Account for the following:
 - (i) Mn^{2+} is more stable than Fe^{2+} towards oxidation to +3 state.
 - (ii) The enthalpy of atomisation is lowest for Zn in 3d series of the transition elements.
 - (iii) Actinoid elements show wide range of oxidation states.

OR

- (i) Name the element of 3d transition series which shows maximum number of oxidation states. Why does it show so?
- (ii) Which transition metal of 3d series has positive $E^{\circ}(M^2/M)$ value and why?
- (iii) Out of Cr3+ and Mn3+, which is a stronger oxidising agent and why?
- (iv) Name a member of the lanthanoid series which is well known to exhibit +2 oxidation state.
- (v) Complete the following equation:

$$MnO_4$$
 8H 5e

30. (a) Write the products of the following reactions:

(i)
$$\bigcirc$$
 = O + H₂N — OH
(ii) 2C₆H₅CHO conc. NaOH

- (b) Give simple chemical tests to distinguish between the following pairs of compounds:
 - (i) Benzaldehyde and Benzoic acid
 - (ii) Propanal and Propanone

- (a) Account for the following:
 - (i) CH₃CHO is more reactive than CH₃COCH₃ towards reaction with HCN.
 - (ii) Carboxylic acid is a stronger acid than phenol.
- (b) Write the chemical equations to illustrate the following name reactions:
 - (i) Wolff-Kishner reduction
 - (ii) Aldol condensation
 - (iii) Cannizzaro reaction

SET-II (Questions Uncommon to Set-I)

- 1. Give one example each of sol and gel.
- **3.** Write the IUPAC name of the compound.

- 5. Some liquids on mixing form 'azeotropes'. What are 'azeotropes'?
- **6.** Arrange the following in increasing order of basic strength.

- 7. Which component of starch is a branched polymer of -glucose and insoluble in water?
- 9. State Henry's law. What is the effect of temperature on the solubility of a gas in a liquid?
- **10.** Define the following terms:
 - (i) Pseudo first order reaction
 - (ii) Half life period of reaction $(t_{1/2})$
- 11. Write the principle behind the following methods of refining:
 - (i) Hydraulic washing
 - (ii) Vapour phase refining
- **22.** (a) Draw the structures of the following:
 - (i) XeF₂
 - (ii) BrF3
- **24.** Account for the following:
 - (i) Bi(V) is a stronger oxidising agent than Sb(V).
 - (ii) N N single bond is weaker than P P single bond.
 - (iii) Noble gases have very low boiling points.
- 25. (i) Name the sweetening agent used in the preparation of sweets for a diabetic patient.
 - (ii) What are antibiotics? Give an example.
 - (iii) Give two examples of macromolecules that are chosen as drug targets.
- 27. (i) Deficiency of which vitamin causes rickets?
 - (ii) Give an example for each of fibrous protein and globular protein.
 - (iii) Write the product formed on reaction of D-glucose with Br₂ water.

SET-III (Questions Uncommon to Set-I and II)

- 1. Give one example each of lyophobic sol and lyophilic sol.
- 2. Write the IUPAC name of the compound.

- 3. What type of intermolecular attractive interaction exists in the pair of methanol and acetone?
- 5. Arrange the following in increasing order of basic strength:

$$C_6H_5NH_2$$
, $C_6H_5NHCH_3$, $C_6H_5N(CH_3)_2$

- **6.** Name the products of hydrolysis of sucrose.
- **9.** State Raoult's law for the solution containing volatile components. What is the similarity between Raoult's law and Henry's law?
- **10.** Explain the following terms:
 - (i) Rate constant (k)
 - (ii) Half life period of a reaction $(t_{1/2})$
- 11. Write the principles of the following methods:
 - (i) Froth floatation method
 - (ii) Electrolytic refining
- **20.** (a) Draw the structures of the following compounds:
 - (i) XeF₄
 - (ii) N₂O₅
- **22.** Account for the following:
 - (i) Sulphur in vapour form exhibits paramagnetic behaviour.
 - (ii) SnCl₄ is more covalent than SnCl₂.
 - (iii) H₃PO₂ is a stronger reducing agent than H₃PO₃.
- **23.** (*i*) What are disinfectants? Give an example.
 - (ii) Give two examples of macromolecules that are chosen as drug targets.
 - (iii) What are anionic detergents? Give an example.
- **24.** (i) Deficiency of which vitamin causes scurvy?
 - (ii) What type of linkage is responsible for the formation of proteins?
 - (iii) Write the product formed when glucose is treated with HI.



SET-I

- Oil in water emulsion: milk, vanishing cream.
 Water in oil emulsion: Butter, cold cream, cod liver oil.
- 2. Scrap iron, $\operatorname{Cu}^{2+}(aq)$ Fe (s) $\operatorname{Cu}(s)$ Fe 2 (aq) or H_2 gas, $\operatorname{Cu}^{2+}(aq)$ H₂(g) $\operatorname{Cu}(s)$ 2H (aq)
- **3.** In both of the given complexes, Co is in +3 oxidation state. As ethane-1, 2-diamine is bidentate chelating ligand, it forms rings and hence $[Co(en)_3]^{3+}$ is more stable complex than $[Co(NH_3)_6]^3$.
- 4. 3-hydroxybutanoic acid
- **5.** o-nitrophenol, is more volatile as intermolecular hydrogen bonding causes association molecules in p-nitrophenol.

Intramolecular hydrogen bonding

Intermolecular hydrogen bonding

- **6.** Two solutions which have the same osmotic pressure at a given temperature are called isotonic solutions.
- 7. The solubility increases in the order in which molecular mass decreases with increase in hydrogen atoms on nitrogen atom which undergo hydrogen bonding, *i.e.*,

$$C_6H_5NH_2$$
 $(C_2H_5)_2NH < C_2H_5NH_2$

- **8.** Amylose is water soluble whereas amylopectin is water insoluble component.
- **9.** For fcc lattice number of atoms per unit cell, z = 4

Here,
$$d = 11.2 \text{ g cm}^{-3}$$
, $a = 4 = 10^{-8} \text{ cm}$, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Atomic mass, $M = \frac{d - a^3 - N_A}{z}$

$$\frac{11.2 \text{ g cm}^{3} \quad (4 \quad 10^{-8} \text{ cm})^{3} \quad 6.022 \quad 10^{23} \text{ mol}^{-1}}{4} = 107.9 \text{ g mol}^{-1} \text{ or } 107.9 u$$

- **10.** (*i*) Schottky defect
 - (ii) Decreases
 - (iii) This type of defect is shown by ionic compounds which have high coordination number and small difference in size of cations and anions e.g. NaCl, KCl etc.
- **11.** Here, T_f 0.48 K, $M_B = 256 \text{ g mol}^{-1}$, $W_A = 75 \text{ g}$, $K_f = 5.12 \text{ K kg mol}^{-1}$.

Mass of solute,
$$W_B = \frac{T_f - M_B - W_A}{K_f - 1000} = \frac{0.48 \text{ K} - 256 \text{ g mol}^{-1} - 75 \text{ g}}{5.12 \text{ K kg mol}^{-1} - 1000 \text{ g kg}^{-1}} = 1.8 \text{ g}$$

- **12.** A solution which obeys Raoult's law over the entire range of concentration is called ideal solution. The important characteristics of an ideal solution are
 - (i) The enthalpy of mixing of pure components to form the solution is zero i.e., $_{\rm mix}H$ 0
 - (ii) The volume of mixing is zero i.e., $_{\rm mix}V$ 0
- 13. Differences between order and molecularity of reaction.

Order		Molecularity	
(i)	It is the sum of the powers of the concentration of the reactants in the rate law expression.	(i)	It is the number of reacting species taking part in an elementary reaction, which must collide simultaneously so as to result into a chemical reaction.
(ii)	It is determined experimentally.	(ii)	It is a theoretical concept.
(iii)	It can be zero or a fraction.	(iii)	It cannot be zero or a fraction.
(iv)	Order is applicable to elementary as well as complex reactions.	(iv)	Molecularity is applicable only for elementary reactions. For complex reactions it has no meaning.

(Any two)

- **14.** (*i*) Zone refining method is based on the principle that the impurities are more soluble in the melt than in the solid state of the metal.
 - (ii) Chromatographic method is based on the principle that different components of a mixture are adsorbed to different extent on an adsorbent. The adsorbed components are removed by using suitable solvents.
- **15.** (*i*) $\operatorname{Ca}_{3} \operatorname{P}_{2}(s) + 6\operatorname{H}_{2} \operatorname{O}(l)$

 $2PH_3 + 3Ca(OH)_2$

(ii) $Cu + 2H_2SO_4(conc.)$

 $CuSO_4 + SO_2 + 2H_2O$

OR

(i) Shorter the bond length, higher is the bond dissociation enthalpy of hydrogen halide. As the atomic size increases down the group the E–H (E = F, Cl, Br, I) bond length increases and hence the bond dissociation enthalpy increases in the reverse order i.e.,


HI HBr HCl HF

(ii) $H_2O < H_2S < H_2Se < H_2Te$


The increase in acidic character from H_2O to H_2Te is due to decrease in bond enthalpy for dissociation of H E(E O, S, Se, Te) bond down the group.

16. [Cr(NH₃)₄ Cl₂] : Tetraamminedichloridochromium (III) ion.

Geometrical isomerism



cis-Tetraamminedichlorido chromium (III) ion



trans-Tetraamminedichlorido chromium (III) ion

17. (i) (i), 2-bromobutane is a chiral molecule.

 $$B_{\Gamma}$$, primary halides undergo faster $S_{N}2$ reactions than secondary halides due to less steric hinderance.

- (ii) (a) S_N2 reaction occurs with inversion of configuration.
 - (b) S_N1 reaction occurs with racemisation.

19. (a) Freundlich adsorption isotherm equation for adsorption of gases on solids:

$$\frac{x}{m} \quad kp^{1/n}(n - 1)$$
or
$$\log \frac{x}{m} \quad \log k \quad \frac{1}{n} \log p$$


where x is the mass of the gas adsorbed on mass m of the adsorbent at pressure p, k and n are constants which depends on the nature of the gas and adsorbent at a particular temperature.

- (b) Important characteristics of lyophilic sols:
 - (i) They are reversible in nature, i.e., once the dispersed phase is separated from dispersion medium the sol can be made again by simply remixing with dispersion medium.
 - (ii) They are quite stable and are not easily coagulated.


(c)

Type of Colloid	Example
Associated colloid	Surface active agents such as soap (CMC is 10^{-4} to 10^{-3} mol L^{-1}) and synthetic detergents.
Multimolecular colloid	Sulphur sol, gold sol

20. (a) (i)




Square pyramidal




112304

Sulphuric acid

(b) White phosphorus consists of discrete tetrahedral P₄ molecule with six P—P covalent bonds.



Red phosphorus has polymeric structure in which P₄ tetrahedra are linked together through P—P covalent bond to form chain.



- 21. (i) The oxidation state of central atom phosphorus is +5 in PCl₅ whereas it is +3 in PCl₃. Higher the positive oxidation of central atom, more will be its polarising power which, in turn, increases the covalent character of bond formed between the central atom and the atoms surrounding it.
 - (ii) Iron reacts with HCl to form FeCl₂ and H₂.

H₂ thus produced prevents the oxidation of FeCl₂ to FeCl₃.

(iii) Ozone is a resonance hybrid of the following two main structures:

As a result of resonance, the two O—O bond lengths in O₃ are equal.

22.
$$k = \frac{2.303}{t} \log \frac{P_0}{2P_0 - P_t}$$

Here,
$$P_0 = 0.4$$
 atm, $t = 100$ s, $P_t = 0.7$ atm
$$k = \frac{2.303}{100} \log \frac{0.4}{2 - 0.4 - 0.7}$$
$$= \frac{2.303}{100} \log \frac{0.4}{0.1} = \frac{2.303}{100} \log 4 = \frac{2.303}{100} = 0.6021$$
$$k = 1.386 = 10^{-2} \text{ s}^{-1}$$

- **23.** (*i*) Biological macromolecules such as carbohydrates, proteins, lipids and nucleic acids are chosen as drug targets.
 - (ii) Antiseptics are the chemical substances which prevent the growth of microorganism or may even kill them but are not harmful to living human tissues, e.g., dettol, soframycin, boric acid, hydrogen peroxide etc.
 - (iii) Use of aspartame is limited to cold foods and soft drink because it is unstable at cooking temperature.
- **24.** (*i*) Vitamin A
 - (ii) Uracil
 - (iii) It suggests that all the six carbon atoms in glucose are linked in a straight chain.
- **25.** (i) Concern towards water pollution, concern for environmental protection, team work, socially aware.
 - (ii) The polymers which can be broken into small segments by enzyme catalysed reactions or to some extent by oxidation over a period of time are called biodegradable polymers. The required enzymes are produced by microorganism e.g., poly--hydroxybutyrate-co--hydroxyvalerate (PHBV), nylon-2-nylon-6 etc.
 - (iii) Polyethene is an addition polymer.

(i)
$$H-Br \rightleftharpoons H^+ + \mathbf{i}Br$$

 $CH_3-CH_2 \overset{\bullet}{\circ} H + H^+ \rightleftharpoons CH_3-CH_2 \overset{\bullet}{\circ} H_2$

Ethyl oxonium ion (Protonated 1° alcohol)

$$(ii) : \overline{Br} + \overline{CH_2} \xrightarrow{CH_3} + \overline{CH_2} \xrightarrow{CH_2} \overline{CH_3} + \overline{Br} \cdot \overline{CH_2} \cdot \overline{OH_2}$$

$$= \overline{CH_3} + \overline{Br} \cdot \overline{CH_2} - \overline{CH_3} + \overline{H_2O}$$

$$= \overline{CH_3} + \overline{CH_3} + \overline{CH_2} - \overline{CH_3} + \overline{H_2O}$$

$$= \overline{CH_3} + \overline{CH_3}$$

Intermediate

27. (i)
$$CH_3Br \xrightarrow{KCN} CH_3CN \xrightarrow{LiAlH_4} CH_3CH_2NH_2 \xrightarrow{HNO_2} CH_3CH_2OH$$

Bromomethane Ethanenitrile Ethanamine (A) (B) (C)

(ii)
$$CH_3COOH \xrightarrow{NH_3} CH_3CONH_2 \xrightarrow{Br_2+KOH} CH_3NH_2 \xrightarrow{CHCl_3+NaOH} CH_3NC$$

Ethanoic acid $CH_3COOH \xrightarrow{A} CH_3CONH_2 \xrightarrow{Br_2+KOH} CH_3NH_2 \xrightarrow{CHCl_3+NaOH} CH_3NC$
(A) $CH_3COOH \xrightarrow{A} CH_3COOH \xrightarrow{A} CH_3NH_2 \xrightarrow{CHCl_3+NaOH} CH_3NC$

OR

(i)
$$NO_2$$
 NH_2 OOD_2 OO

Nitrobenzene

Aniline

(ii)
$$CH_3COOH \xrightarrow{NH_3} CH_3CONH_2 \xrightarrow{Br_2/NaOH} CH_3NH_2$$

Ethanoic acid Ethanamide Methanamine

O Ethanoic anhydride

or

$$\begin{array}{c|cccc} & & & & & & O \\ NH_2 & O & & NH-C-CH_3 \\ & & & & & & \\ & & + CH_3-C-Cl & \xrightarrow{Pyridine} & & + HCl \\ \end{array}$$

Aniline Ethanoyl chloride

N-phenylethanamide

N-phenylethanamide

- **28.** (a) (i) The molar conductivity when concentration approaches to zero is called limiting molar conductivity.
 - (ii) A fuel cell is a device which converts energy produced during the combustion of fuels like hydrogen, methane, methyl alcohol etc. directly into electrical energy. One such successful fuel cell is hydrogen-oxygen fuel cell.
 - (b) For $0.1 \text{ mol } L^{-1} \text{ KC}l$ solution

Conductivity, =
$$1.29 \times 10^{-2}$$
 $^{-1}$ cm⁻¹, Resistance, R 100

Cell constant = Conductivity \times resistance

$$= 1.29 \times 10^{-2}$$
 $^{-1}$ cm⁻¹ × 100 $= 1.29$ cm⁻¹

For $0.02 \text{ mol } L^{-1} \text{ solution}$

Resistance = 520 , Cell constant = 1.29 cm^{-1} ,

Conductivity,
$$=\frac{\text{Cell constant}}{\text{Resistance}}$$

$$\frac{1.29 \text{ cm}^{-1}}{520} = 0.00248$$
 $^{-1} \text{ cm}^{-1}$

Molar conductivity,
$$_{m}$$

$$\frac{\text{Conductivity () } 1000 \text{ cm}^{3} \text{L}^{-1}}{\text{Molarity}}$$

$$\frac{0.00248 \quad {}^{1}\text{cm}^{-1} \quad 1000 \text{ cm}^{3} \text{ L}^{-1}}{0.02 \text{ mol L}^{-1}}$$

$$124 \quad {}^{1}\text{cm}^{2}\text{mol}^{-1}$$

OR

(a) It states that the amount of chemical reaction which occurs at any electrode during electrolysis by a current is proportional to the quantity of electricity passed through the electrolyte (solution or melt).

$$Cu^{2+} + 2e^{-}$$
 Cu

Charge required for reduction of 1 mol of $Cu^{2+} = 2F$.

 $Mq^{2+} + 2e^{-}$ (b) At anode: Mg

At cathode:
$$Cu^{2+} + 2e^{-}$$
 Cu

$$Mg + Cu^{2+}$$
 $Mg^{2+} + Cu$

At cathode:
$$Cu^{2+} + 2e^{-}$$
 Cu

$$Mg + Cu^{2+}$$
 $Mg^{2+} + Cu$

Nernst equation: $E_{cell} = E_{cell}^{o} - \frac{0.059}{n} \log \frac{[Mg^2]}{[Cu^2]}$ (at 298K)

Here,
$$E_{\text{cell}}^{o} = 2.71 \text{ V}$$
, $[Mg^{2+}] = 0.1 \text{ M}$, $[Cu^{2+}] = 0.01 \text{ M}$, $n = 2$
 $E_{\text{cell}} = 2.71 \frac{0.059}{2} \log \frac{0.1}{0.01} 2.71 0.029 1$
 $= 2.681 \text{ V}$

29. (a) (i) Pyrolusite is fused with KOH in the presence of atmospheric oxygen to give K₂MnO₄.

(ii) Na₂CrO₄ is extracted with water and acidified with H₂SO₄ to give Na₂Cr₂O₇.

- (b) (i) The electronic configuration of Mn² is [Ar] 3d⁵ which is half-filled and hence stable. So Mn² cannot lose third electron easily. On the other hand, Fe² has electronic configuration [Ar] 3d⁶. It tends to lose one electron to acquire stable [Ar] 3d⁵ electronic configuration. Hence Mn²⁺ is more stable than Fe²⁺ towards oxidation.
 - (ii) In the formation of metallic bonds, no electrons from 3*d*-orbitals are involved in case of zinc, while in all other metals of the 3*d* series, electrons from the *d*-orbitals are always involved in the formation of metallic bonds. That is why, the enthalpy of atomisation of zinc is the lowest in the series.
 - (iii) This is due to comparable energies of 5f, 6d and 7s orbitals.

- (i) Manganese $(3d^5 4s^2)$ shows maximum number of oxidation states as its atoms have 5 unpaired electrons in 3d orbitals. It shows all the oxidation states from +2 to +7.
- (ii) Copper has positive E^{o} (M²⁺/M) value, as high energy ($_{a}H$ $_{i}H$) to transform Cu(s) to Cu²⁺(aq) is not balanced by hydration enthalpy.
- (iii) Mn^{3+} is stronger oxidising agent as the charge from Mn^{3+} to Mn^{2+} results in half filled, d^5 configuration which has extra stability.
- (iv) Europium, Eu. Eu²⁺ is formed by losing the two s electrons and its (Xe) f^7 configuration accounts its formation.

30. (a) (i)
$$\longrightarrow$$
 O + H₂N-OH \longrightarrow Cyclohexanone oxime

(ii) 2C₆H₅CHO + NaOH \longrightarrow C₆H₅ - CH₂ - OH + C₆H₅COO Na (Conc.) Benzyl alcohol Sodium benzoate

(iii) CH₃COOH \longrightarrow Cl₂/P \longrightarrow Cl - CH₂ - COOH + HCl 2-chloro ethanoic acid

(b) (i) Benzoic acid being an acid reacts with NaHCO₃ solution to give brisk effervescence due to evolution of CO₂ while benzaldehyde does not response to this test.

COOH

$$+ \text{NaHCO}_3$$
 $+ \text{COONa}$
 $+ \text{CO}_2 \uparrow + \text{H}_2 O$

Sodium benzoate

 $+ \text{CHO}$
 $+ \text{NaHCO}_3$

No effervescence due to evolution of CO₂ gas

(ii) Propanal being an aldehyde reduces Tollens reagent to silver mirror but propanone being a ketone does not.

$$\label{eq:ch3} \text{CH}_3\text{CH}_2\text{CHO} + 2[\text{Ag}(\text{NH}_3)_2]^+ + 3\text{OH}^- \\ \longrightarrow \text{CH}_3\text{CH}_2\text{COO}^- + 2\text{Ag} \downarrow + 4\text{NH}_3 + 2\text{H}_2\text{O} \\ \text{Silver} \\ \text{mirror}$$

- (a) (i) The methyl group due to its +I effect reduce the magnitude of positive charge on carbonyl carbon atom. Moreover it also hinders the approach of nucleophile CN. Since in acetaldehyde there is one methyl while in acetone there are two methyl groups attached to carbonyl group therefore acetaldehyde is more reactive than acetone towards nucleophilic addition with HCN.
 - (ii) Because the release of proton from carboxylic acid is much easier than from phenol as the conjugate base carboxylate ion is much more resonance stabilised than conjugate base phenoxide ion.
- (b) (i) Wolff-Kishner reduction

$$C = O \qquad \begin{array}{c} NH_2NH_2 \\ Aldehyde \\ or ketone \end{array} \qquad C = NNH_2 \\ H_2O \qquad CH_3 \\ CH_3 \qquad C = O \qquad \begin{array}{c} CH_3 \\ H_2O \qquad CH_3 \\ CH_3 \end{array} \qquad C = NNH_2 \\ CH_3 \qquad CH_3 \qquad CH_3 \qquad CH_3 \\ CH_3 \qquad CH_3 \qquad CH_3 \qquad CH_3 \qquad CH_3 \\ Acetone \qquad CH_3 \qquad CH_2 \qquad CH_3 + N_2 \\ \end{array}$$

(ii) Aldol condensation

CH₃ C H + H CH₂ CHO
$$\stackrel{\text{dil. NaOH}}{\rightleftharpoons}$$
 CH₃ CH CH₂ CHO $\stackrel{\text{dil. NaOH}}{\rightleftharpoons}$ CH₃ CH CH₂ CHO $\stackrel{\text{dil. NaOH}}{\rightleftharpoons}$ CH₃ CH=CH CHO $\stackrel{\text{But-2-enal}}{\rightleftharpoons}$ CH₄ O OH O CH₃ CH=CH CHO $\stackrel{\text{But-2-enal}}{\rightleftharpoons}$ CH₃ C CH₂ C CH₃ $\stackrel{\text{CH}_3}{\rightleftharpoons}$ CH₃ C CH₂ C CH₃ $\stackrel{\text{CH}_3}{\rightleftharpoons}$ CH₃ CH₂ C CH₃ $\stackrel{\text{CH}_3}{\rightleftharpoons}$ CH₃ C=CH C CH₃ $\stackrel{\text{CH}_3}{\rightleftharpoons}$ CH₃ C=CH C CH₃ $\stackrel{\text{CH}_3}{\rightleftharpoons}$ CH₃ C+-methylpent-3-ene-2-one

(iii) Cannizzaro reaction

SET-II

 Sol: Paints, cell fluids Gel: Butter, cheese

3. 3-aminobutanal

5. Azeotropes are the binary mixtures of liquids having the same composition in liquid and vapour phase and boil at a constant temperature e.g. a mixture of 68% nitric acid and 32% water by mass.

6. $C_6H_5NH_2$ $C_6H_5NHCH_3$ $C_6H_5CH_2NH_2$

7. Amylopectin

9. It states that "the partial pressure of the gas in vapour phase (p) is directly proportional to the mole fraction of gas (x) in the solution".

Mathematically, $p = K_H x$

where K_H is the Henry's law constant.

The solubility of a gas in liquid decreases with rise in temperature as dissolution of a gas in a liquid is an exothermic process.

10. (*i*) A reaction which is not truely of first order but under certain conditions becomes a reaction of first order is called pseudo first order reaction *e.g.* acid hydrolysis of ethyl acetate.

$$CH_3COOC_2H_5 + H_2O$$
 H^+
 $CH_3COOH + C_2H_5OH$


Rate $[CH_3COOC_2H_5]$ as H_2O is in excess.

(ii) The half life $(t_{1/2})$ of a reaction is the time in which the concentration of reactant is reduced to one half of its initial concentration $[R]_0$.


For a first order reaction
$$t_{1/2} = \frac{0.693}{k}$$
 i.e., independent of $[R]_0$.
For a zero order reaction $t_{1/2} = \frac{[R]_0}{2k}$ *i.e.*, $t_{1/2} = [R]_0$.

- **11.** (*i*) Hydraulic washing: It is based on the differences in the gravities of the ore and the gangue particles.
 - (ii) Vapour phase refining: In this, metal is converted into its volatile compound and collected elsewhere. It is then decomposed to give pure metal. So, the two requirements are:
 - (a) The metal should form a volatile compound with an available reagent.
 - (b) The volatile compound should be easily decomposable, so that the recovery is easy.

22. (*a*) (*i*) There are two bond pairs and three lone pairs of electrons around central Xe atom in XeF₂. Therefore according to VSEPR Theory XeF₂ should be linear.



(*ii*) There are three bond pairs and two lone pairs of electrons around central Br atom in BrF₃. Therefore according to VSEPR theory BrF₃ should be slightly bent "T".



- 24. (i) Due to inert pair effect +3 oxidation state of Bi is more stable than its +5 oxidation state while +5 oxidation state of Sb is more stable than its +3 oxidation state. Therefore, Bi (V) can more easily accept a pair of electrons to form more stable Bi (III) than Sb (V) can accept a pair of electrons and hence Bi (V) is stronger oxidising agent than Sb (V).
 - (ii) N-N single bond is weaker than P-P single bond due to large interelectronic repulsion between the lone pairs of electrons present on the N atoms of N-N bond having small bond length.
 - (iii) Noble gases being monoatomic have no interatomic forces except weak London or dispersion forces and therefore, they are liquefied at very low temperatures. Hence, they have very low boiling points.
- **25.** (*i*) Saccharin, sucrolose.
 - (ii) Antibiotics are the chemical substance produced wholly or partly by chemical synthesis, which in low concentrations, either kill or inhibit the growth of microorganisms by intervening in their metabolic processes. A few examples of antibiotics are chloramphenicol, ofloxacin, penicillin.
 - (iii) Biomolecules such as carbohydrates, proteins, lipids and nucleic acids are chosen as drug targets.
- **27.** (*i*) Vitamin D
 - (ii) Fibrous protein keratin, myosin Globular protein — insulin, albumins

SET-III

- **1.** Lyophobic sol Gold sol, As₂S₃ sol Lyophilic sol — Sol of starch, sol of gum
- 2. 4-hydroxypentan-2-one
- 3. Hydrogen bonds
- 5. $C_6H_5NH_2 < C_6H_5NHCH_3 < C_6H_5N (CH_3)_2$
- **6.** $C_{12}H_{22}O_{11} + H_2O$ Sucrose $C_6H_{12}O_6 + C_6H_{12}O_6$ $D_{-(+)}$ -Glucose $D_{-(-)}$ -Fructose
- **9.** It states that for a solution of volatile liquids, the partial vapour pressure of each component in the solution is directly proportional to its mole fraction. According to Raoults law, for a volatile component, A of the solution

 $P_A = x_A$ or $P_A = P_A^{\circ} x_A$, where P_A° is the vapour pressure of pure component A.

If one of the component is so volatile that it exist as a gas then according to Henry's law

p $K_H x$, where K_H is the Henry law constant *i.e.*, the partial vapour pressure of the volatile component (gas) is directly proportional to its mole fraction in the solution.

Thus the similarity between Raoult's law and Henry's law is that both state that the partial vapour pressure of the volatile component or gas is directly proportional to its mole fraction in the solution.


- 10. (i) Rate constant, k may be defined as the rate of reaction when concentration of each reactant is taken as unity.
 - (ii) Refer to Ans. 10 (ii) Set-II CBSE Delhi 2014.
- 11. (i) This method of concentration of ore is based on the principle that the surface of sulphide ores is preferentially wetted by oils while that of gangue is preferentially wetted by water.
 - (ii) Electrolytic refining: In this method, impure metal is made to act as anode. A strip of same metal in pure form is used as cathode. They are put in a electrolytic bath containing soluble salt of same metal. On passing electric current, metal ions from the electrolyte solution are deposited at the cathode in the form of pure metal while an equivalent amount of metal dissolves from the anode and goes into the electrolyte solution as metal ions, i.e.,

At cathode: M^n (aq) ne^- M(s)

At anode: M(s) M^n (aq) ne^-

The voltage applied for the electrolysis is such that the impurities of more electropositive metals remain in the solution as ions where impurities of less basic metals settle down under the anode as anode mud. A large number of metals such as copper, gold, silver, zinc, aluminium, etc., are refined by this method.

20. (a) (i) There are four bond pairs and two lone pairs of electrons around central Xe atom in XeF₄. Therefore according to VSEPR theory XeF₄ should be square planar.



(N₂O₅)

F Xe F

(XeF₄) Square planar

- 22. (i) In vapour form sulphur partly exists as S₂ molecules which have two unpaired electrons in the antibonding * molecular orbitals like O₂ molecule and hence, exhibits paramagnetism.
 - (ii) The oxidation of central atom Sn is +4 in SnCl₄ while it is +2 in SnCl₂. Higher the positive oxidation of central atom, more will be its polarising power which, in turn increases the covalent character of bond formation between the central atom and the atoms around it.
 - (iii) Acids which contains P—H bonds have reducing character. Since H_3PO_2 contains two P—H bonds while H_3PO_3 contains only one P—H bond therefore H_3PO_2 is a stronger reducing agent than H_3PO_3 .
- **23.** (*i*) Chemical substances which kill microorganisms or stop their growth but are not safe to be applied to the living tissues are called disinfectants. For example, chlorine in concentrations of 0.2 to 0.4 ppm is used for sterilisation of water to make it fit for drinking.
 - (ii) Biomolecules such as carbohydrates, proteins, liquids and nucleic acids are chosen as drug targets.
 - (iii) Anionic detergents: These are so named because large part of their molecules are anions and it is the anionic part of the molecule which is involved in the cleansing action. These are sodium salt of sulphonated long chain alcohols or hydrocarbons. For example, sodium lauryl sulphate, sodium dodecylbenzene sulphonate, etc. Anionic detergents are used in household work and in toothpastes.
- **24.** (*i*) Vitamin C
 - (ii) Peptide linkage
 - (iii) On prolonged heating with HI, glucose gives n-hexane.



CBSE Examination Paper All India-2014

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper Delhi-2014.

SET-I

1. What is the effect of temperature on chemisorption?

- 2. What is the role of zinc metal in the extraction of silver?
- 3. What is the basicity of H_3PO_3 ?
- **4.** Identify the chiral molecule in the following pair:



Cl

5. Which of the following is a natural polymer?

Buna-S, Proteins, PVC

- **6.** The conversion of primary aromatic amines into diazonium salts is known as ______.
- 7. What are the products of hydrolysis of sucrose?
- **8.** Write the structure of p-methylbenzaldehyde.
- 9. An element with density 2.8 g cm⁻³ forms a f.c.c. unit cell with edge length 4×10^{-8} cm. Calculate the molar mass of the element.

(Given: $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$)


- 10. (i) What type of non-stoichiometric point defect is responsible for the pink colour of LiCl?
 - (ii) What type of stoichiometric defect is shown by NaCl?

OR

How will you distinguish between the following pairs of terms:

- (i) Tetrahedral and octahedral voids
- (ii) Crystal lattice and unit cell
- 11. State Kohlrausch law of independent migration of ions. Why does the conductivity of a solution decrease with dilution?

12. For a chemical reaction R P, the variation in the concentration (R) vs. time (t) plot is given as



- (i) Predict the order of the reaction.
- (ii) What is the slope of the curve?
- 13. Explain the principle of the method of electrolytic refining of metals. Give one example.
- **14.** Complete the following equations:
 - (i) $P_4 + H_2O$
 - (ii) $XeF_4 + O_2F_2$
- **15.** Draw the structures of the following:
 - (i) XeF₂

- (ii) BrF3
- **16.** Write the equations involved in the following reactions:
 - (i) Reimer-Tiemann reaction
 - (ii) Williamson synthesis
- **17.** Write the mechanism of the following reaction:

HBr

$$CH_3CH_2Br + H_2O$$

- **18.** Write the name of monomers used for getting the following polymers:
 - (i) Bakelite

- (ii) Neoprene
- **19.** (a) Calculate ${}_{r}G^{\circ}$ for the reaction

$$Mg(s) + Cu^{2+}(aq)$$

$$Mg^{2+}(aq) + Cu(s)$$

 $Mg(s) + Cu^{2+}(aq)$ $Mg^{2+}(aq) + Cu(s)$ Given: E_{cell}^{o} 2.71 V, 1 F 96500 C mol⁻¹

- (b) Name the type of cell which was used in Apollo space programme for providing electrical power.
- 20. The following data were obtained during the first order thermal decomposition of SO₂Cl₂ at a constant volume:

$$SO_2Cl_2(g)$$
 $SO_2(g) + Cl_2(g)$

Experiment	Time/s ⁻¹	Total pressure/atm
1	0	0.4
2	100	0.7

Calculate the rate constant.

(Given: $\log 4 = 0.6021$, $\log 2 = 0.3010$)

21. What are emulsions? What are their different types? Give one example of each type.

- **22.** Give reasons for the following:
 - (i) $(CH_3)_3$ P=O exists but $(CH_3)_3$ N=O does not.
 - (ii) Oxygen has less electron gain enthalpy with negative sign than sulphur.
 - (iii) H₃PO₂ is a stronger reducing agent than H₃PO₃.
- 23. (i) Write the IUPAC name of the complex $[Cr(NH_3)_4Cl_2]Cl$.
 - (ii) What type of isomerism is exhibited by the complex $[Co(en)_3]^{3+}$? (en = ethane-1,2-diamine)
 - (iii) Why is $[NiCl_4]^{2-}$ paramagnetic but $[Ni(CO)_4]$ is diamagnetic? (At nos.: Cr = 24, Co = 27, Ni = 28)
- 24. (a) Draw the structures of major monohalo products in each of the following reactions:

(i)
$$\sim$$
 CH₂OH $\xrightarrow{\text{PCl}_5}$ (ii) \sim CH₂— CH = CH₂ + HBr \longrightarrow

- (b) Which halogen compound in each of the following pairs will react faster in S_N2 reaction?
 - (i) CH₃Br or CH₃I
 - (ii) (CH₃)₃ C—C1 or CH₃—C1
- **25.** Account for the following:
 - (i) Primary amines (R-NH₂) have higher boiling point than tertiary amines (R₃N).
 - (ii) Aniline does not undergo Friedel-Crafts reaction.
 - (iii) (CH₃)₂NH is more basic than (CH₃)₃N in an aqueous solution.

Give the structures of A, B and C in the following reactions:

(i)
$$C_6H_5NO_2$$
 Sn HCl A NaNO₂ HCl B H_2O C
(ii) CH_3CN H_2O/H^+ A NH₃ B Br_2+KOH C

- **26.** Define the following terms as related to proteins:
 - (i) Peptide linkage
 - (ii) Primary structure
 - (iii) Denaturation
- 27. On the occasion of World Health Day, Dr. Satpal organised a 'health camp' for the poor farmers living in a nearby village. After check-up, he was shocked to see that most of the farmers suffered from cancer due to regular exposure to pesticides and many were diabetic. They distributed free medicines to them. Dr. Satpal immediately reported the matter to the National Human Rights Commission (NHRC). On the suggestions of NHRC, the government decided to provide medical care, financial assistance, setting up of super-speciality hospitals for treatment and prevention of the deadly disease in the affected villages all over India.
 - (i) Write the values shown by
 - (a) Dr. Satpal
- (b) NHRC
- (ii) What type of analgesics are chiefly used for the relief of pains of terminal cancer?
- (iii) Give an example of artificial sweetener that could have been recommended to diabetic patients.

- **28.** (a) Define the following terms:
 - (i) Molarity
 - (ii) Molal elevation constant (K_b)
 - (b) A solution containing 15 g urea (molar mass = 60 g mol⁻¹) per litre of solution in water has the same osmotic pressure (isotonic) as a solution of glucose (molar mass = 180 g mol⁻¹) in water. Calculate the mass of glucose present in one litre of its solution.

- (a) What type of deviation is shown by a mixture of ethanol and acetone? Give reason.
- (b) A solution of glucose (molar mass = 108 g mol^{-1}) in water is labelled as 10% (by mass). What would be the molality and molarity of the solution? (Density of solution = 1.2 g mL^{-1})
- **29.** (a) Complete the following equations:
 - (i) $Cr_2O_7^{2-} + 2OH^-$
 - (ii) $MnO_4^- + 4H^+ + 3e^-$
 - (b) Account for the following:
 - (i) Zn is not considered as a transition element.
 - (ii) Transition metals form a large number of complexes.
 - (iii) The E° value for the Mn^{3+}/Mn^{2+} couple is much more positive than that for Cr^{3+}/Cr^{2+} couple.

OR

- (i) With reference to structural variability and chemical reactivity, write the differences between lanthanoids and actinoids.
- (ii) Name a member of the lanthanoid series which is well known to exhibit +4 oxidation state.
- (iii) Complete the following equation:

$$MnO_4^- + 8H^+ + 5e^-$$

(iv) Out of Mn^{3+} and Cr^{3+} , which is more paramagnetic and why? (Atomic nos.: Mn = 25, Cr = 24)

- **30.** (a) Write the products formed when CH₃CHO reacts with the following reagents:
 - (i) HCN
 - (ii) H₂N—OH
 - (iii) CH₃CHO in the presence of dilute NaOH
 - (b) Give simple chemical tests to distinguish between the following pairs of compounds:
 - (i) Benzoic acid and Phenol
 - (ii) Propanal and Propanone

OR

- (a) Account for the following:
 - (i) Cl—CH₂COOH is a stronger acid than CH₃COOH.
 - (ii) Carboxylic acids do not give reactions of carbonyl group.

- (b) Write the chemical equations to illustrate the following name reactions:
 - (i) Rosenmund reduction
 - (ii) Cannizzaro's reaction
- (c) Out of CH₃CH₂ —CO—CH₂—CH₃ and CH₃CH₂—CH₂—CO—CH₃, which gives iodoform test?

SET-II (Questions Uncommon to SET-I)

- 1. Why is adsorption always exothermic?
- 2. Name the method that is used for refining of nickel.
- **3.** Why does NO₂ dimerise?
- **4.** Based on molecular forces what type of polymer is neoprene?
- 5. What are the products of hydrolysis of maltose?
- **6.** Write the structure of 4-chloropentan-2-one.
- **9.** Write the name of monomers used for getting the following polymers:
 - (i) Terylene
 - (ii) Nylon-6, 6
- **10.** Describe the role of the following:
 - (i) SiO₂ in the extraction of copper from copper matte
 - (ii) NaCN in froth floatation process
- 11. Complete the following equations:
 - (i) Ag + PCl₅
 - (ii) CaF₂ + H₂SO₄
- **12.** Draw the structures of the following:
 - (i) XeF₄
 - (ii) HClO₄
- **13.** (*i*) Write the type of magnetism observed when the magnetic moments are oppositively aligned and cancel out each other.
 - (ii) Which stoichiometric defect does not change the density of the crystal?
- **14.** Define the following terms:
 - (i) Fuel cell
 - (ii) Limiting molar conductivity $\binom{m}{m}$
- **19.** Define the following terms:
 - (i) Glycosidic linkage
 - (ii) Invert sugar
 - (iii) Oligosaccharides

SET-III (Questions Uncommon to Set-I and II)

- 1. What are the dispersed phase and dispersion medium in milk?
- 2. Name the method used for refining of copper metal.

354 Xam idea Chemistry—XII

- **3.** Why does NH₃ act as a Lewis base?
- **5.** Which of the following is a fibre?
 - Nylon, Neoprene, PVC
- **6.** Write the products of hydrolysis of lactose.
- **8.** Write the structure of 2-hydroxybenzoic acid.
- **9.** Complete the following equations:
 - (i) C + conc. H_2SO_4
 - (ii) $XeF_2 + H_2O$
- **10.** Draw the structures of the following:
 - (i) XeO₃
 - (ii) H₂SO₄
- 11. Write the name of monomers used for getting the following polymers:
 - (i) Teflon
 - (ii) Buna-N
- **13.** (*i*) Write the type of magnetism observed when the magnetic moments are aligned in parallel and anti-parallel directions in unequal numbers.
 - (ii) Which stoichiometric defect decreases the density of the crystal?
- **14.** Define the following terms:
 - (i) Molar conductivity $\binom{m}{m}$
 - (ii) Secondary batteries
- 17. Write the principle behind the froth floatation process. What is the role of collectors in this process?
- **23.** Define the following terms:
 - (i) Nucleotide
 - (ii) Anomers
 - (iii) Essential amino acids


SET-I

- 1. Chemisorption initially increases then decreases with rise in temperature. The initial increase is due to the fact that heat supplied acts as activation energy. The decrease afterwards is due to the exothermic nature of adsorption equilibrium.
- 2. Zinc acts as reducing agent in the extraction of silver. It reduces Ag to Ag and itself get oxidised to Zn^2 .

$$2Na[Ag(CN)_2] + Zn$$

$$Na_2[Zn(CN)_4] + 2Ag$$

3. As H₃PO₃ contains two ionizable P-OH bonds. Therefore, it is dibasic.



- 5. Proteins
- **6.** Diazotisation.
- 7. Sucrose on hydrolysis gives one molecule each of glucose and fructose.

$$C_6H_{12}O_6 + C_6H_{12}O_6$$

8.

p-methylbenzaldehyde

9. For f.c.c. unit cell number of atoms per unit cell, z = 4

$$M = \frac{d N_A a^3}{z}$$

Here,
$$d = 2.8 \text{ gcm}^{-3}$$
, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$, $a = 4 \times 10^{-8} \text{ cm}$

$$M = \frac{2.8 \text{ g cm}^{-3} + 6.022 + 10^{23} \text{ mol}^{-1} + (4 + 10^{-8} \text{ cm})^{3}}{4}$$

$$M = 26.98 \text{ g mol}^{-1}$$

- 10. (i) Metal excess defect due to anionic vacancies.
 - (ii) Schottky defect.

- (i) A void surrounded by four spheres occupying the corners of tetrahedron is called a tetrahedral void. It is much smaller than the size of spheres in the close packing. A void surrounded by six spheres along the corners of an octahedral is called octahedral void. The size of the octahedral void is smaller than that of the spheres in the close packing but larger than the octahedral void.
- (ii) The regular three dimensional arrangement of identical points in the space which represent how the constituent particles (atoms, ions, molecules) are arranged in a crystal is called a crystal lattice.

A unit cell is the smallest portion of a crystal lattice, which when repeated over and again in different directions produces the complete crystal lattice.

11. Kohlrausch law states that the limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of cation and anion of the electrolyte.

In general, if an electrolyte on dissociation gives cations and anions then its limiting molar conductivity is given by

m

and where. are the limiting molar conductivities of cations and anions respectively.

Conductivity of a solution decreases with dilution. This is due to the fact that the number of ions per unit volume that carry the current in a solution decreases with dilution.

(i) Order of the reaction is zero.

(ii) Slope =
$$\frac{d[R]}{dt}$$
 k

13. Electrolytic refining: In this method, impure metal is made to act as anode. A strip of same metal in pure form is used as cathode. They are put in an electrolytic bath containing soluble salt of same metal. On passing electric current, metal ions from the electrolyte solution are deposited at the cathode in the form of pure metal while an equivalent amount of metal dissolves from the anode and goes into the electrolyte solution as metal ions, i.e.,


 M^n (aq) ne^- At cathode:

M(s) M^n (aq) ne^- At anode:


The voltage applied for the electrolysis is such that the impurities of more electropositive metals remain in the solution as ions whereas impurities of less basic metals settle down under the anode as anode mud. A large number of metals such as copper, gold, silver, zinc, aluminium, etc., are refined by this method.

14. (*i*) P_4 + H_2O No reaction (*ii*) XeF_4 + O_2F_2 XeF_6 + O_2

15. (i) There are two bond pairs and three lone pairs electrons around central Xe atom in XeF2. Therefore according to VSEPR theory XeF2 should be linear.



(ii) There are three bond pairs and two lone pairs of electrons around central Br atom in BrF₃. Therefore according to VSEPR theory BrF₃ should be slightly bent "T".



16. (i) Reimer–Tiemann reaction:

(ii) Williamson synthesis:

- **17.** Refer to Ans 26 (*a*) Set-I, CBSE Delhi 2014.
- **18.** (*i*) Phenol and formaldehyde
 - (ii) Chloroprene

19. (a)
$$Mg(s) + Cu^{2+}(aq)$$
 $Mg^{2+}(aq) + Cu(s)$

$${}_{r}G^{\circ} = -nFE_{cell}^{\circ}$$
Here, $n = 2$, $F = 96500 \text{ C mol}^{-1}$ and $E_{cell}^{\circ} = 2.71 \text{ V}$


$${}_{r}G^{\circ} = -2 \times 96500 \text{ C mol}^{-1} \times 2.71 \text{ V}$$

$$= -523030 \text{ J mol}^{-1}$$

$$= -523.03 \text{ kJ mol}^{-1}$$

- (b) H₂—O₂ fuel cell
- 20. Refer to Ans. 22 Set-I, CBSE Delhi 2014.
- **21.** Emulsions are the colloidal solutions in which both the dispersed phase and dispersion medium are liquids. Emulsion can be classified into two types. These are:
 - (i) Oil in water (O/W) type emulsion: In this type of emulsions oil acts as disperse phase and water acts as dispersion medium e.g., milk, vanishing cream.
 - (ii) Water in oil (W/O) type emulsion: In this type of emulsions water acts as disperse phase and oil acts as dispersion medium e.g., butter, cod liver oil, cold cream.

- 22. (i) Phosphorus due to presence of d orbitals in its valence shell forms p d multiple bonds. As a result phosphorus can expand its covalency beyond four. Therefore, phosphorus form R₃P=O in which its covalency is 5. In contrast, nitrogen due to absence of d-orbitals, cannot form p d multiple bonds and hence cannot expand its covalency beyond 4. Therefore, (CH₃)₃N=O does not exist.
 - (ii) The size of O atom is smaller as compared to S. Thus when an electron is added to isolated gaseous O atom, the interelectronic repulsions encountered in the smaller 2*p*-orbitals of O are larger than those encountered in the larger 3*p*-orbitals of S.
 - As a result less energy is released in case of O than in case of S. Hence oxygen has less electron gain enthalpy with negative sign than sulphur.
 - (iii) Acids which contains P–H bonds have reducing character. Since H₃PO₂ contains two P–H bonds while H₃PO₃ contains only one P–H bond therefore H₃PO₂ is a stronger reducing agent than H₃PO₃.
- 23. (i) Tetraamminedichloridochromium (III) chloride.
 - (ii) Complex $[Co(en)_3]^{3+}$ shows optical isomerism.



(iii) In complex $[NiCl_4]^{2-}$, Ni is in +2 oxidation state with the electronic configuration $3d^8 4s^0$. As Cl^- is a weak ligand it cannot pair up the electrons in 3d orbitals therefore $[NiCl_4]^{2-}$ is paramagnetic. In $[Ni(CO)_4]$, Ni is in zero oxidation state with the electronic configuration $3d^8 4s^2$. CO is a strong ligand it causes 4s electrons to shift to 3d and pair up 3d electrons. Since there is no unpaired electron in the complex $Ni(CO)_4$, therefore, it is diamagnetic.

24. (a) (i)
$$\longrightarrow$$
 CH₂OH $\xrightarrow{\text{PCl}_5}$ \longrightarrow CH₂Cl + POCl₃ + HCl

(ii) \longrightarrow CH₂— CH = CH₂ + HBr \longrightarrow \longrightarrow CH₂— CH — CH₃

Br

- (b) (i) CH₃—I. As iodine is better leaving group due to its larger size, it will be released at a faster rate in the presence of incoming nucleophile.
 - (ii) CH₃Cl. As three bulky methyl groups hinder the approach of nucleophile in (CH₃)C—Cl.
- **25.** (*i*) In primary amines, two hydrogen atoms are present on N-atom and they undergo extensive intermolecular hydrogen bonding which results in association of molecules while in tertiary amines, no hydrogen atom is present on N-atom. Hence there is no hydrogen bonding in tertiary amines. As a result of this primary amines have higher boiling point than tertiary amines.

- (ii) Aniline being a Lewis base reacts with AlCl₃ a Lewis acid and catalyst used in Friedel crafts reaction to form salt C₆H₅NH₂AlCl₃. Due to presence of a +ve charge on N-atom in the salt, the group NH₂AlCl₃ acts as a strong deactivating group. It reduces the electron density in the benzene ring as a result of this aniline does not undergo Friedel-crafts reaction.
- (iii) The basicity of amine in aqueous solution depends upon the stability of the substituted ammonium cation. Here the combination of three factors, +ve I effect of CH₃ groups, hydrogen bonding and steric hindrance favour greater stability for ammonium cation of dimethyl amine than ammonium cation of trimethyl amine. Hence dimethylamine is stronger base than trimethyl amine.

(ii)
$$A = CH_3 - C - OH$$
, $B = CH_3 - C - NH_2$, $C = CH_3 - NH_2$
Ethanoic acid Ethanamide Methanamine O

- 26. (i) A peptide linkage is an amide (— C— NH —) linkage formed between —COOH group of one -amino acid and —NH₂ group of other -amino acid by loss of a water molecule.
 - (ii) The specific sequence in which various -amino acids present in a protein are linked to one another is called its primary structure. Any change in its primary structure creates a new protein.
 - (iii) **Denaturation of Proteins:** When a protein in its native form is subjected to a change, such as change in temperature or change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called denaturation of protein. During denaturation, 2° and 3° structures are destroyed but 1° structures remain intact, *e.g.*, coagulation of egg while on boiling, curdling of milk, etc.
- **27.** (*i*) (*a*) The values shown by Dr. Satpal are concern for the health of others, dedicated towards work, kind, compassionate.
 - (b) The values shown by NHRC are responsiveness, having understanding of public health in rural area, dutiful, caring.
 - (ii) Narcotic analgesics such as morphine, heroin, codeine are used for the relief of pains of terminal cancer.
 - (iii) Saccharin, Sucrolose.
- **28.** (a) (i) Molarity may be defined as number of moles of solute dissolved in one litre of solution.

Molarity =
$$\frac{\text{Moles of solute}}{\text{Volume of solution (in litre)}}$$

(ii) Molal elevation constant may be defined as the elevation in boiling point when one mole of solute is dissolved in 1000 grams of the solvent.

(b) Osmotic pressure,
$$=\frac{W_B R T}{M_R V}$$

Osmotic pressure of urea solution = $\frac{15 R T}{60 1}$

Osmotic pressure of glucose solution = $\frac{W_B - R - T}{180 - 1}$

As, Osmotic pressure of urea solution = Osmotic pressure of glucose solution

Therefore,
$$\frac{15 R T}{60 1} = \frac{W_B R T}{180 1}$$

Mass of glucose, $W_B = \frac{15 - 180}{60} = 45 \text{ g}$

OR

(a) A mixture of ethanol and acetone shows positive deviation from Raoults law.

In pure ethanol hydrogen bond exist between the molecules. On adding acetone to ethanol, acetone molecules get in between the molecules of ethanol thus breaking some of the hydrogen bonds and weakening molecular interactions considerably. Weakening of molecular interactions leads to increase in vapour pressure resulting in positive deviation from Raoults law

(b) Let the mass of solution = 100 g

Mass of glucose =
$$10 g$$

Number of moles of glucose =
$$\frac{\text{Mass of glucose}}{\text{Molar mass}}$$

= $\frac{10 \text{ g}}{180 \text{ g mol}^{-1}} = 0.056 \text{ mol}$

Volume of solution =
$$\frac{\text{Mass of solution}}{\text{Density of solution}}$$

= $\frac{100 \text{ g}}{1.2 \text{ g mL}^{-1}} = 83.3 \text{ mL}$
= $\frac{83.3}{1000} \text{ L} = 0.083 \text{ L}$

$$\begin{aligned} \text{Molarity} &= \frac{\text{Moles of solute}}{\text{Volume of solution in litre}} \\ &= \frac{0.056 \text{ mol}}{0.083 \text{ L}} = 0.67 \text{ mol L}^{-1} \end{aligned}$$

Mass of solvent, water =
$$100 \text{ g} - 10 \text{ g} = 90 \text{ g}$$

= $\frac{90 \text{ g}}{1000 \text{ g kg}^{-1}} = 0.09 \text{ kg}$

Molality =
$$\frac{\text{Moles of glucose}}{\text{Mass of water in kg}}$$

= $\frac{0.056 \text{ mol}}{0.09 \text{ kg}} = 0.62 \text{ mol kg}^{-1}$

29. (a) (i)
$$Cr_2O_7^{2-} + 2OH^ 2CrO_4^{2-} + H_2O$$
 (ii) $MnO_4^- + 4H^+ + 3e$ Heat $MnO_2 + 2H_2O$

- (b) (i) As zinc atom has completely filled d orbitals $(3d^{10} 4s^2)$ in its ground state as well as in its oxidised state, therefore zinc is not considered as transition element.
 - (ii) Transition metals form a large number of complexes due to following reasons:
 - Small size and high charge of the ions of transition metals.
 - Presence of vacant orbitals of appropriate energy which can accept lone pairs of electrons donated by ligand.
 - (iii) Mn^{2+} has the electronic configuration [Ar]3 d^5 which is half filled and hence very stable. Due to this the third ionisation energy of Mn is very high. Thus the much large third ionisation energy of Mn (where the required change is d^5 to d^4) is mainly responsible for this.

OR

(<i>i</i>)			
	S.No.	Lanthanoids	Actinoids
	(<i>i</i>)	Atomic or ionic radii does not show much variation.	Atomic or ionic radii show much variation.
	(ii)	Besides +3 oxidation state, they show +2 and +4 oxidation states in few cases.	Besides +3 oxidation states, they show higher oxidation states of +4, +5, +6, +7 also.

(ii) Cerium, Ce ([Xe] $4f^2 5d^0 6s^2$). Formation of Ce⁴⁺ is favoured by its noble gas configuration ([Xe] $4f^0 5d^0 6s^0$).

(iii)
$$MnO_4^- + 8H^+ + 5e$$
 $Mn^{2+} + 4H_2O$

(iv) $\operatorname{Mn}^{3+}(3d^4 4s^0)$ has 4 unpaired electrons while $\operatorname{Cr}^{3+}(3d^3 4s^0)$ has 3 unpaired electrons therefore Mn^{3+} is more paramagnetic than Cr^{3+} .

30. (a) (i)
$$\overset{\text{CH}_3}{\underset{\text{H}}{\sim}}\text{C=O}$$
 + HCN $\xrightarrow{\text{pH 9-10}}$ $\overset{\text{CH}_3}{\underset{\text{H}}{\sim}}\text{C}\overset{\text{OH}}{\underset{\text{CN}}{\sim}}$

Ethanal Hydrogen cyanide Ethanal cyanohydrin

(ii) $\overset{\text{CH}_3}{\underset{\text{H}}{\sim}}\text{C=O}$ + H₂N—OH $\xrightarrow{\text{H}^+}$ $\xrightarrow{\text{PH 3.5}}$ $\overset{\text{CH}_3}{\underset{\text{H}}{\sim}}\text{C=N-OH}$ + H₂O Ethanal oxime

$$CH_3$$
 — $CH = CH$ — C — C H But 2 enal

(b) (i) Phenol gives a violet colour with neutral FeCl₃ solution while benzoic acid gives buff coloured precipitate of ferric benzoate.


(ii) Propanal being an aldehyde reduces Tollens' reagent to produce silver mirror but propanone being a ketone does not.

$$\begin{array}{c} \text{CH}_3\text{CH}_2\text{CHO} + 2[\text{Ag}(\text{NH}_3)_2]^+ + 3\text{OH} & \text{CH}_3\text{CH}_2\text{COO} & + 2\text{Ag} & + 4\text{NH}_3 + 2\text{H}_2\text{O} \\ \text{Propanal} & \text{Silver} & \text{mirror} \end{array}$$

CH₃COCH₃ Tollens' reagent No silver mirror Propanone

OR

- (a) (i) Because of –I effect of Cl atom in ClCH₂COOH and +I effect of CH₃ group in CH₃COOH the electron density in the O—H bond in ClCH₂COOH is much lower than CH₃COOH. As a result O—H bond in ClCH₂COOH is much weaker than in CH₃COOH therefore loses a proton more easily than CH₃COOH. Hence ClCH₂COOH acid is stronger acid than CH₃COOH.
 - (ii) Carboxylic acids are resonance hybrid of the following structures:



Similarly, a carbonyl group of aldehydes and ketones may regarded as resonance hybrid of following structures.

$$>$$
C $\stackrel{\frown}{=}$ $\overset{}{\circ}$: $>$ C $\stackrel{}{=}$ $\overset{}{\circ}$: $>$ V

Because of contribution of structure (IV), the carbonyl carbon in aldehydes and ketones is electrophilic. On the other hand electrophilic character of carboxyl carbon is reduced due to contribution of structure (II). As carbonyl carbon of carboxyl group is less electropositive than carbonyl carbon in aldehydes and ketones therefore carboxylic acids do not give nucleophilic addition reactions of aldehydes and ketones.

(b) (i) Rosenmund reduction:

O

R C Cl + H₂
Acyl chloride

O

C-Cl

$$+$$
H₂
 $-$
Pd-BaSO₄
S

R C H + HCl

Aldehyde

O

C-H

Benzoyl chloride

Benzaldehyde

(ii) Cannizzaro reaction:

2HCHO conc. KOH
$$CH_3$$
 OH + HCOO K Methyl alcohol Potassium formate

CHO CH_2 —OH $COO^{-\frac{1}{2}}$

Benzaldehyde Benzyl alcohol Sodium benzoate

(c) CH_3 — CH_2 — CH_2 — $COCH_3$

SET-II


- **1.** Adsorption occurs with decrease in entropy *i.e.*, S is -ve. As G H T S and for a process to be spontaneous G must be -ve. This can be possible only when H is -ve for the process *i.e.*, adsorption is exothermic.
- **2.** Mond process (Vapour phase refining)
- 3. NO₂ is an odd electron molecule in which N has seven electrons in its valence shell and hence is less stable. To become more stable by acquiring noble gas configuration having 8 electrons in the valence shell of N, it undergoes dimerisation to form N₂O₄.
- 4. Elastomer
- 5. Maltose on hydrolysis gives two molecules of glucose.

- **9.** (i) Ethylene glycol and terephthalic acid
 - (ii) Hexamethylenediamine and adipic acid
- **10.** (*i*) Copper matte chiefly consists of Cu₂S and some FeS. Silica acts as a flux in the extraction of copper from copper matte to remove ferrous oxide as ferrous silicate slag.

(ii) NaCN is used as depressant in froth floatation process. The role of a depressant is to prevent one type of sulphide particle from forming the froth with air bubble. For example to separate lead sulphide (PbS) ore from ZnS, NaCN is used as a depressant as it forms zinc complex, Na₂[Zn(CN)₄] on the surface of ZnS thereby preventing it from forming froth. Under these conditions, only PbS forms froth and hence is separated from ZnS.


11. (i)
$$2Ag + PCl_5$$
 $2AgCl + PCl_3$ (ii) $CaF_2 + H_2SO_4$ $2HF + CaSO_4$

12. (*i*) As there are 4 bond pairs and two lone pairs of electrons around central Xe atom in XeF₄ therefore according to VSEPR theory XeF₄ should be square planar.



(XeF₄) Square planar

(ii) Perchloric acid



- 13. (i) Antiferromagnetism
 - (ii) Frenkel defect
- **14.** (*i*) A fuel cell is a device which converts the energy produced during the combustion of fuels like hydrogen, methanol, methane etc. directly into electrical energy. One of the most successful fuel cell is H₂—O₂ fuel cell.
 - (ii) When concentration approaches zero, the molar conductivity is known as limiting molar conductivity. It is represented by m.

$$_m$$
 ($_m$) when c 0

- **19.** (*i*) The linkage between two monosaccharides through oxygen atom in an oligosaccharide or a polysaccharide is known as glycosidic linkage.
 - (ii) Sucrose is dextrorotatory (+ 66.5°) but after hydrolysis it gives an equimolar mixture of D-(+)-glucose and D-(-)-fructose, which is laevorotatory. This change of specific rotation from dextrorotation to laevorotation is called inversion of sugar and the mixture obtained is called invert sugar.

(iii) Carbohydrates which on hydrolysis give two to ten molecules of monosaccharides are called oligosaccharides e.g., sucrose.

SET-III

- 1. In milk liquid fat acts as dispersed phase and water acts as dispersion medium.
- 2. Electrolytic refining
- 3. N atom in NH₃ has one lone pair of electrons which is available for donation. Therefore it acts as a Lewis base.
- 5. Nylon

6.
$$C_{12}H_{22}O_{11} + H_2O$$
 H^+ $C_6H_{12}O_6 + C_6H_{12}O_6$ Lactose Galactose

COOH

2-hydroxy-benzoic acid


9. (i) C + $2H_2SO_4(conc.)$ $CO_2 + 2SO_2 + 2H_2O$

$$CO_2 + 2SO_2 + 2H_2O_2$$


(ii) $2XeF_2(s) + 2H_2O(l)$

$$2Xe(g) + 4HF(aq) + O_2(g)$$

10. (*i*)



Pyramidal XeO₃



H₂SO₄

- **11.** (*i*) Tetrafluoroethene
 - (ii) 1, 3-butadiene and acrylonitrile
- **13.** (*i*) Ferrimagnetism
 - (ii) Schottky defect
- 14. (i) Molar conductivity, m of a solution at a dilution V is defined as the conductance of all the ions produced from one gram mole of the electrolyte dissolved in $V \text{ cm}^3$ of the solution when the electrodes are one centimetre apart and the area of the electrodes is so large that the whole of the solution is contained between them.


$$_{m}$$
 V or $_{m}$ $\frac{1000}{\text{Molarity}}$

is the conductivity and V is the volume of the solution containing 1 mole of the where electrolyte.

(ii) Secondary batteries are those batteries which can be recharged by passing electric current through them and hence can be used over again e.g. lead storage battery.

This method of concentration of ore is based upon the principle that the surface of sulphide ore is preferentially wetted by oils while that of gangue is preferentially wetted by water. Collectors enhance the non-wettability of the ore particles in the froth floatation process e.g. pine oil.

23. (*i*) A nucleotide contains all the three basic components of nucleic acid *i.e.*, a pentose sugar, a nitrogeneous base and a phosphoric acid. When nucleoside is linked to phosphoric acid at 5' position of sugar moiety, we get a nucleotide.



- (ii) A pair of stereoisomers such as -D(+) glucose and -D-(+) glucose which differ in configuration only around C_1 are called anomers.
- (iii) The amino acids which cannot be synthesised in our body and must be obtained through diet are known as essential amino acids *e.g.*, valine, lysine, histidine.



CBSE Examination Paper Foreign-2014

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper Delhi-2014.

SET-I

1. What is the function of collectors in the froth floatation process for the concentration of ores?

- 2. What type of forces are responsible for the occurrence of physisorption?
- 3. Why is the single N—N bond weaker than the single P—P bond?
- **4.** What type of isomerism is shown by the following complex: $[Co(NH_3)_6][Cr(CN)_6]$
- 5. Express the relationship between atomic radius (r) and the edge length (a) in the b.c.c. unit cell.
- **6.** Write the IUPAC name of the following compound:

7. Which of the two is more basic and why?

- **8.** Name the two components of -glucose which constitute starch.
- 9. A solution of Ni(NO₃)₂ is electrolysed between platinum electrodes using a current of 5.0 ampere for 20 minutes. What mass of nickel will be deposited at the cathode?
 (Given: At. Mass of Ni = 58.7 g mol⁻¹, 1 F = 96500 C mol⁻¹)
- 10. Define half-life of a reaction. Write the expression of half-life for
 - (i) zero order reaction and
 - (ii) first order reaction.
- 11. Write the chemical reactions involved in the extraction of silver from silver ore.
- **12.** Name the two most important allotropes of sulphur. Which one of the two is stable at room temperature? What happens when the stable form is heated above 370 K?

(i) Write the conditions to maximise the yield of H₂SO₄ by contact process.

(ii) Why is $K_{a_2} K_{a_1}$ for H_2SO_4 in water?

13. Complete the following equations:

(i)
$$2MnO_4$$
 $5S^2$ $16H$

(ii)
$$\operatorname{Cr}_2\operatorname{O}_7^2$$
 2OH

14. Write the state of hybridisation, shape and IUPAC name of the complex $[CoF_6]^3$.

(Atomic no. of Co = 27)

15. Write chemical equations when

- (i) ethyl chloride is treated with aqueous KOH.
- (ii) chlorobenzene is treated with CH₃COCl in presence of anhydrous AlCl₃.
- **16.** (a) Which alkyl halide from the following pairs would you expect to react more rapidly by S_N2 mechanism and why?

$$CH_3 - CH_2 - CH - CH_3$$
 $CH_3 - CH_2 - CH_2 - CH_2 - Br$

- (b) Racemisation occurs in S_N1 reactions. Why?
- 17. Write the mechanism of the following reaction:

- **18.** Name the reagents used in the following reactions:
 - (i) Bromination of phenol to 2, 4, 6-tribromophenol
 - (ii) Butan-2-one to Butan-2-ol
 - (iii) Friedel-Crafts alkylation of anisole
 - (iv) Oxidation of primary alcohol to carboxylic acid
- **19.** (i) What type of stoichiometric defect is shown by KCl and why?
 - (ii) What type of semiconductor is formed when silicon is doped with As?
 - (iii) Which one of the following is an example of molecular solid:

- (iv) What type of substances would make better magnets, ferromagnetic or ferrimagnetic?
- **20.** (i) Write two advantages of $H_2 O_2$ fuel cell over ordinary cell.
 - (ii) Equilibrium constant (K_c) for the given cell reaction is 10. Calculate E_{cell}.

$$\mathbf{A}(s) + \mathbf{B}^{2+}(aq) + \mathbf{B}(s)$$

21. The following data were obtained during the first order thermal decomposition of SO₂Cl₂ at a constant volume:

$$SO_2Cl_2(g)$$
 $SO_2(g) + Cl_2(g)$

Experiment	Time/s ⁻¹	Total pressure/atm
1	0	0.4
2	100	0.7

Calculate the rate constant.

(Given: $\log 4 = 0.6021$, $\log 2 = 0.3010$)

- **22.** (a) Write the expression for the Freundlich adsorption isotherm for the adsorption of gases on solids, in the form of an equation.
 - (b) What are the dispersed phase and dispersion medium of butter?
 - (c) A delta is formed at the meeting place of sea and river water. Why?
- **23.** (a) What are the different oxidation states exhibited by the lanthanoids?
 - (b) Write two characteristics of the transition elements.
 - (c) Which of the 3d-block elements may not be regarded as the transition elements and why?

OR

Assign suitable reasons for the following:

- (a) The Mn²⁺ compounds are more stable than Fe²⁺ towards oxidation to their +3 state.
- (b) In the 3d series from Sc (Z = 21) to Zn (Z = 30), the enthalpy of atomization of Zn is the lowest.
- (c) Sc³⁺ is colourless in aqueous solution whereas Ti³⁺ is coloured.
- **24.** Give the structures of A, B and C in the following reactions:
 - (i) CH_3Br KCN A LiAlH₄ B $\frac{HNO_2}{273 \text{ K}}$ C
 - (ii) CH₃COOH NH₃ A Br₂ KOH B CHCl₃ NaOH (
- **25.** Define the following terms:
 - (a) Anomers
 - (b) Denaturation of proteins
 - (c) Essential amino acids
- **26.** (*i*) Define Antihistamine with an example.
 - (ii) Which one of the following drugs is an antibiotic:

Morphine, Equanil, Chloramphenicol, Aspirin

- (iii) Why is use of aspartame limited to cold food and drink?
- 27. After the ban on plastic bags, students of one school decided to create awareness among the people about the harmful effects of plastic bags on the environment and the Yamuna river. To make it more impactful, they organised a rally by joining hands with other schools and distributed paper bags to vegetable vendors, shopkeepers and departmental stores. All students pledged not to use polythene bags in future to save the Yamuna river.

After reading the above passage, answer the following questions:

- (i) What values are shown by the students?
- (ii) What are biodegradable polymers? Give one example.
- (iii) Is polythene a homopolymer or copolymer?
- **28.** (a) State Raoult's law for a solution containing volatile components. Name the solution which follows Raoult's law at all concentrations and temperatures.
 - (b) Calculate the boiling point elevation for a solution prepared by adding 10 g of CaCl₂ to 200 g of water. (K_b for water = 0.512 K kg mol⁻¹, Molar mass of CaCl₂ = 111 g mol⁻¹)

OR

- (a) Define the following terms:
 - (i) Azeotrope
 - (ii) Osmotic pressure
 - (iii) Colligative properties
- (b) Calculate the molarity of 9.8% (w/w) solution of H_2SO_4 if the density of the solution is 1.02 g mL⁻¹. (Molar mass of $H_2SO_4 = 98 \text{ g mol}^{-1}$)
- **29.** (a) Account for the following:
 - (i) Bi is a strong oxidizing agent in the +5 state.
 - (ii) PCl₅ is known but NCl₅ is not known.
 - (iii) Iron dissolves in HCl to form FeCl₂ and not FeCl₃.
 - (b) Draw the structures of the following:
 - (i) XeOF₄
 - (ii) HClO₄

OR

- (a) Draw the structures of the following:
 - (i) H₂S₂O₈
 - (ii) Red P₄
- (b) Account for the following:
 - (i) Sulphur in vapour state exhibits paramagnetism.
 - (ii) Unlike xenon, no distinct chemical compound of helium is known.
 - (iii) H₃PO₂ is a stronger reducing agent than H₃PO₃.
- **30.** (a) Write the products formed when ethanol reacts with the following reagents:
 - (i) CH₃MgBr and then H₃O⁺
 - (ii) Zn-Hg/conc.HCl
 - (iii) C₆H₅CHO in the presence of dilute NaOH
 - (b) Give simple chemical tests to distinguish between the following pairs of compounds:
 - (i) Benzoic acid and Ethyl benzoate
 - (ii) Propanal and Butan-2-one

OR

- (a) Account for the following:
 - (i) CH₃CHO is more reactive than CH₃COCH₃ towards reaction with HCN.
 - (ii) There are two -NH₂ groups in semicarbazide (H₂NNHCONH₂). However, only one is involved in the formation of semicarbazone.
- (b) Write the chemical equation to illustrate each of the following name reactions:
 - (i) Rosenmund reduction
 - (ii) Hell-Volhard-Zelinsky reaction
 - (iii) Cannizzaro reaction

SET-II (Questions Uncommon to SET-I)

- 1. On what principle is the method of zone refining of metals based?
- 2. Name the temperature above which the formation of micelles takes place.
- 3. Arrange the following in the increasing order of their basic character:

4. What type of isomerism is exhibited by the following complex:

$$[Co(NH3)5SO4]Cl$$

- **5.** Express the relationship between atomic radius (r) and the edge length (a) of the f.c.c. unit cell.
- **6.** Which of the two is more basic and why?

- 7. Name the compound which on polymerisation gives neoprene.
- **9.** Write the role of the following:
 - (i) CO in the purification of nickel.
 - (ii) Graphite rod in the electrometallurgy of aluminium.
- 10. Complete the following equations:

$$(i) 2 \text{MnO}_4^- + 5 \text{NO}_2^- + 6 \text{H}^+$$

$$(ii) \operatorname{Cr_2O_7}^{2-} + 14 \operatorname{H}^+ + 6 e^-$$

- 11. Write the state of hybridisation, shape and IUPAC name of the complex $[Ni(CN)_4]^{2-}$. (Atomic no. of Ni = 28)
- 12. Write chemical equations when
 - (i) methyl chloride is treated with AgNO₂.
 - (ii) bromobenzene is treated with CH₃Cl in the presence of anhydrous AlCl₃.
- **13.** Name the reagents used in the following reactions:
 - (i) Nitration of phenol to 2, 4, 6-trinitrophenol
 - (ii) Butanal to Butanol
 - (iii) Friedel-Crafts acetylation of anisole
 - (iv) Oxidation of primary alcohol to aldehyde
- **20.** Define the following terms:
 - (i) Polysaccharides
 - (ii) Amino acids
 - (iii) Enzymes


SET-III (Questions Uncommon to Set-I and II)

- 1. Name the depressant which is used to separate ZnS and PbS ores in froth floatation process.
- 2. Based on the type of dispersed phase, what type of colloid is micelles?
- **4.** What type of isomerism is exhibited by the complex $[Co(NH_3)_5NO_2]^{2+}$?
- **5.** Write IUPAC name of the following compound:

$$\begin{array}{c} \mathrm{HO}-\mathrm{CH_2}-\mathrm{CH}-\mathrm{CH_2}-\mathrm{OH} \\ | \\ \mathrm{OH} \end{array}$$

372 Xam idea Chemistry—XII

6. Which of the two is more basic and why?



- 7. What are biocatalysts? Give an example.
- **9.** Write the role of the following:
 - (i) Iodine in the refining of zirconium.
 - (ii) Silica in the extraction of copper from copper matte.
- 11 Complete the following equations:
 - (i) $2MnO_4$ $5SO_3^{2-}$ 6H (ii) $2CrO_4^2$ 2H
- 12. Write the state of hybridization, shape and IUPAC name of the complex $[Co(NH_3)_6]^{3+}$. (Atomic no. of Co = 27)
- 13. Write chemical equations when
 - (i) ethyl chloride is treated with alcoholic KOH.
 - (ii) chlorobenzene is treated with CH₃Cl in the presence of anhydrous AlCl₃.
- **15.** Name the different reagents needed to perform the following reactions:
 - (i) Phenol to benzene
 - (ii) Dehydration of propan-2-ol to propene
 - (iv) Dehydrogenation of ethanol to ethanal
- **24.** Define the following terms:
 - (a) Invert sugar
 - (b) Vitamins
 - (c) Nucleoside





SET-I

- 1. Collectors (e.g., pine oil, xanthates etc.) enhance non-wettability of the ore particles.
- 2. Van der Waals forces.
- **3.** N—N bond is weaker than P—P bond because of large interelectronic repulsions between lone pairs of electrons present on the N atom of N—N bond having small bond length.
- 4. Co-ordination isomerism
- **5.** r (atomic radius) $\frac{\sqrt{3}}{4} a$ (edge length of unit cell)
- **6.** 2-hydroxybenzaldehyde
- 7. In CH₃NH₂, the +I effect of —CH₃ group makes lone pair of electrons on N-atom more available for donation. On the other hand in C₆H₅NH₂, the resonance effect causes delocalisation of lone pair of electrons over benzene ring and makes it less available for donation. Hence, CH₃NH₂ is more basic than C₆H₅NH₂.
- 8. The two component of starch are amylose and amylopectin.
- **9.** $Q I t = 5 A \times 20 \times 60 s = 6000 C$

$$Ni^2$$
 2e Ni

 2×96500 C deposit Ni = 58.7 g

6000 C will deposit Ni
$$\frac{58.7 \text{ g mol}^{-1}}{2 + 96500 \text{ C mol}^{-1}}$$
 6000 C 1.825 g

- 10. The half life $(t_{1/2})$ of a reaction is the time in which the concentration of a reactant is reduced to one half of its initial concentration.
 - (i) $t_{1/2}$ for a zero order reaction = $\frac{[R]_0}{2k}$ where $[R]_0$ = initial concentration, k = rate constant
 - (ii) $t_{1/2}$ for a first order reaction $\frac{0.693}{k}$
- 11. The chemical reaction involved in the extraction of silver from silver ore are

$$4 \mathrm{Ag}(s) \quad 8 \mathrm{NaCN}(\mathit{aq}) \ 2 \mathrm{H}_2 \mathrm{O}(\mathit{l}) \quad \mathrm{O}_2(\mathit{g})$$

$$4\text{Na} \left[\text{Ag(CN)}_{2} \right] (aq) + 4\text{NaOH}(aq)$$

Sodium dicyanoargentate (I) (Soluble complex)

$$2\text{Na}\left[\text{Ag(CN)}_{2}\right](aq) + \text{Zn}(s) \qquad \qquad \text{Na}_{2}\left[\text{Zn(CN)}_{4}\right](aq) + 2\text{Ag}(s)$$

12. Two most important allotropes of sulphur are rhombic sulphur and monoclinic sulphur. The stable form at room temperature is rhombic sulphur. When rhombic sulphur is heated above 370 K it gets converted into monoclinic sulphur.

(i) The key step in the manufacture of H₂SO₄ by contact process is

$$2SO_2(g)$$
 $O_2(g) \rightleftharpoons 2SO_3(g)$ $_rH$ $196.6 \text{ kJ mol}^{-1}$

The forward reaction is exothermic and proceed with decrease in number of moles. Therefore, low temperature (optimum temperature 720 K), high pressure (in practice 2 bar) and use of catalyst V_2O_5 to increase the rate of reaction at low temperature are the favourable conditions for maximum yield of H_2SO_4 .

(ii) H₂SO₄ is a dibasic acid, it ionizes in two steps and has two dissociation constants.

$$\begin{split} & \text{H}_2\text{SO}_4(aq) + \text{H}_2\text{O}(l) \iff \text{H}_3\text{O}^+(aq) + \text{HSO}_4(aq); \text{K}_a \quad 10 \\ & \text{HSO}_4(aq) \quad \text{H}_2\text{O}(l) \iff \text{H}_3\text{O}^+(aq) + \text{SO}_4^2 \quad (aq); \text{K}_{a_2} \quad 1.2 \times 10^{-2} \end{split}$$

 K_{a_2} K_a , because the negatively charged HSO_4 ion has much less tendency to donate a proton to H_2O as compared to neutral H_2SO_4 .

13. (i)
$$MnO_4$$
 8H 5e Mn^2 $4H_2O] \times 2$

$$S^2$$
 S $2e$] 5
$$2MnO_4$$
 $5S^2$ $+ 16H$ $2Mn^2$ $5S$ $8H_2O$
(ii) $Cr_2O_7^2$ $2OH$ $2CrO_4^2$ H_2O

14. $[CoF_6]^3$: Hexafluoridocobaltate (III) ion; Co^3 [Ar] $3d^6$

<i>3d</i>	48	4 <i>p</i>	4 <i>d</i>
$\uparrow\downarrow$ \uparrow \uparrow \uparrow		XX XX XX	XX XX

XX = Electron pair from ligand F

Hybridisation =
$$sp^3d^2$$
, Shape = Octahedral

15. (i)
$$CH_3$$
— CH_2 — $Cl + KOH(aq)$ \longrightarrow CH_3 — CH_2 — $OH + KCl$ Ethyl chloride

(ii) Cl O Anhyd.AlCl₃ Cl O Cl Chlorobenzene Acetyl Chloride Acetyl Chloride Acetyl Ninor)
$$O$$
 Cl O Cl

- **16.** (a) CH₃—CH₂—CH₂—CH₂—Br. Being primary alkyl halide, there would be less steric hinderance.
 - (b) This is due to the fact that carbocations are the intermediates in $S_N 1$ reactions. Carbocations being sp^2 hybridised are planar species, therefore, the attack of the nucleophile on it occurs from both the faces (front and rear) with almost equal ease resulting in the formation of 50:50 mixture of two enantiomers *i.e.*, racemic mixture.
- 17. Refer to Ans. 26 (a) Set-I CBSE Delhi 2014.
- **18.** (*i*) Aqueous bromine (Br_2/H_2O)
 - (ii) Lithium aluminium hydride (LiAlH₄) or sodium borohydride (NaBH₄).
 - (iii) Anhydrous aluminium chloride (anhyd. AlCl₃)

- (*iv*) Acidified potassium dichromate solution (K₂Cr₂O₇/H₂SO₄) or alkaline potassium permanganate solution (KMnO₄/OH⁻) followed by hydrolysis with dil. H₂SO₄.
- **19.** (i) KCl shows schottky defect as the cation, K⁺ and anion, Cl⁻ are of almost similar sizes.
 - (ii) n-type semiconductor.
 - (iii) CO₂
 - (iv) Ferromagnetic
- **20.** (i) The two main advantages of H_2 – O_2 fuel cell over ordinary cell are as follows:
 - It has high efficiency of 60%–70%.
 - It does not cause any pollution.

(ii) A(s) B² (aq)
$$\rightleftharpoons$$
 A² (aq) B(s)
Here, n 2, K = 10
$$E_{\text{cell}}^{0} = \frac{0.059}{n} \log K_{c}$$

$$E_{\text{cell}}^{0} = \frac{0.059}{2} \times \log 10 = \frac{0.059}{2} = 1 = 0.0295 \text{ V}$$

- 21. Refer to Ans. 22, Set-I CBSE Delhi 2014.
- 22. (a) Freundlich adsorption isotherm equation for adsorption of gases on solids:

$$\frac{x}{m} kp^{1/n}(n - 1).$$
or $\log \frac{x}{m} \log k - \frac{1}{n} \log p$

where x is the mass of the gas adsorbed on mass m of the adsorbent at pressure p, k and n are constants which depends on the nature of the gas and adsorbent at a particular temperature.

- (b) In butter, water acts as disperse phase and oil acts as dispersion medium.
- (c) River water is a colloidal solution of clay. Sea water contains a variety of electrolytes. When river water meets the sea water, the electrolytes present in sea water coagulate the colloidal solution of clay resulting in its deposition with the formation of delta.
- **23.** (a) The common oxidation state of lanthanoid is +3. Besides this some of the lanthanoid also exhibit +2 and +4 oxidation states.
 - (b) The important characteristics of the transition elements are:
 - (i) All transition elements are metallic in nature.
 - (ii) Transition elements exhibit variable oxidation states.
 - (iii) Most of the transition elements form coloured compounds.
 - (iv) A number of transition elements and their compounds show catalytic properties.
 - (c) In 3d transition series zinc is not regarded as transition element as Zn atom has completely filled d orbitals $(3d^{10})$ in its ground state as well as in its oxidised state.

OR

(a) The electronic configuration of Mn² is [Ar] $3d^5$ which is half filled and hence stable. So Mn² cannot lose third electron easily. On the other hand, Fe² has electronic configuration [Ar] $3d^6$. It tends to lose one electron to acquire stable [Ar] $3d^5$ electronic configuration. Hence Mn²⁺ is more stable than Fe²⁺ towards oxidation.

- (b) In the formation of metallic bonds, no electrons from 3d-orbitals are involved in case of zinc, while in all other metals of the 3d series, electrons from the d-orbitals are always involved in the formation of metallic bonds. That is why, the enthalpy of atomisation of zinc is the lowest in the series.
- (c) Ti^{3+} has one electron in *d*-orbitals $(3d^1)$ which can absorb light in visible region for d-d transition. Hence, it is coloured in aqueous solution. Sc^{3+} has no d electron $(3d^0)$, therefore, no light is absorbed for d-d transition. Hence, it is colourless in aqueous solution.
- 24. Refer to Ans 27 Set-I CBSE Delhi 2014.
- **25.** (a) The carbohydrates which differ in configuration at the glycosidic carbon (i.e., C_1 in aldoses and C_2 in ketoses) are called anomers e.g., -D-(+)-glucose and -D-(+)-glucose.
 - (b) **Denaturation of Proteins:** When a protein in its native form is subjected to a change, such as change in temperature or change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called denaturation of protein. During denaturation, 2° and 3° structures are destroyed but 1° structures remain intact, e.g., coagulation of egg while on boiling, curdling of milk, etc.
 - (c) The amino acids which cannot be synthesised in our body and must be obtained through diet are known as essential amino acids e.g. valine, lysine, histidine.
- **26.** (*i*) The drugs which interfere with the natural action of histamine by competing with histamine for binding of receptor where histamine exerts its effect are called anti histamine *e.g.*, brompheniramine, terfenadine, cetrizine etc.
 - (ii) Chloramphenicol
 - (iii) Use of aspartame is limited to cold foods because it is unstable at cooking temperature.
- 27. (i) Concern towards water pollution, concern for environmental protection, team work, socially aware
 - (ii) The polymers which can be broken into small segments by enzyme catalysed reactions or to some extent by oxidation over a period of time are called biodegradable polymers. The required enzymes are produced by microorganism e.g., poly--hydroxybutyrate-co--hydroxyvalerate (PHBV), nylon-2-nylon-6 etc.
 - (iii) Polyethene is an addition polymer.
- **28.** (a) It states that for a solution of volatile liquids, the partial vapour pressure of each component in the solution is directly proportional to its mole fraction. Thus, for a solution of volatile liquids A and B, P_A x_A and P_B x_B or $P_A = P_A^{\circ} x_A$ and $P_B = P_B^{\circ} x_B$ where P_A and P_B are partial vapour pressures, x_A and x_B are mole fractions, P_A° and P_B° are vapour pressure of pure components A and B respectively.

Ideal solution.

(b) The given quantities are:

$$W_B = 10 \text{ g, } W_A = 200 \text{ g, } K_b = 0.512 \text{ K Kg mol}^{-1},$$
 $M_B = 111 \text{ g mol}^{-1}$
 $CaCl_2 = Ca^2 = 2Cl$

On assuming complete dissociating of $CaCl_2$, $i = \frac{3}{1} - 3$

Substituting these values in the formula, $T_b = \frac{i - K_b - W_B - 1000}{M_B - W_A}$, we get


OR

- (a) (i) The binary mixtures of liquids having same composition in liquid and vapour phase and boil at a constant temperature are called azeotropes.
 - (ii) The excess of pressure which must be applied to the solution side to prevent the passage of solvent into it through a semipermeable membrane is called osmotic pressure.
 - (iii) The properties of solutions which depend only on the number of solute particles in the solution but independent of their nature are called colligative properties.
- (b) Let the mass of solution = 100 g


$$\begin{aligned} \text{Mass of H_2SO}_4 &= 9.8 \text{ g} \\ \text{Number of mole of H_2SO}_4 &= \frac{\text{Mass of H_2SO}_4}{\text{Molar mass}} \\ &= \frac{9.8 \text{ g}}{98 \text{ g mol}^{-1}} = 0.1 \text{ mol} \\ &\frac{\text{Mass of solution}}{\text{Density of solution}} \\ &\frac{100 \text{ g}}{1.02 \text{ g mL}^{-1}} \cdot \frac{100}{1.02} \text{ mL} \\ &\frac{100}{1.02} \text{ mL} \cdot \frac{1}{1000 \text{ mL L}^{-1}} \\ &\frac{1}{10.2} \text{L} \\ &\text{Molarity} &= \frac{\text{Moles of solute, H_2SO}_4}{\text{Volume of solution (in L)}} \\ &\frac{0.1 \text{ mol}}{1.02} \\ &\frac{1 \text{ L}}{10.2} \\ &\frac{1}{10.2} \text{ mol} \\ &\frac{1}{10.2} \text{ m$$

- **29.** (a) (i) Due to inert pair effect Bi in +3 state is much more stable than in +5 state. Therefore, Bi in +5 state accepts two electrons and get reduced to more stable +3 state. Hence Bi is strong oxidising in +5 state.
 - (ii) Nitrogen with n = 2, has s and p orbitals only. It does not have d orbitals to expand its covalency beyond four. Hence, NCl₅ is not known. PCl₅ is known as P has vacant 3d orbital to which 3s electrons can be excited to make available five half filled orbitals needed for the formation of five P—Cl bonds.
 - (iii) Iron reacts with HCl to form FeCl2 and H2.

H₂ thus produced prevents the oxidation of FeCl₂ to FeCl₃.



Perchloric acid




- (b) (i) In vapour form sulphur partly exists as S_2 molecules which have two unpaired electrons in the antibonding * molecular orbitals like O_2 molecule and hence, exhibits paramagnetism.
 - (ii) This is due to the following reasons:

He does not have d-orbitals in the valence shell and hence electron cannot be excited to higher energy levels like in Xe to form bonds.

Ionisation enthalpy of He is sufficiently higher than those of oxygen and fluorine.

(iii) Acids which contains P—H bonds have reducing character. Since H₃PO₂ contains two P—H bonds while H₃PO₃ contains only one P–H bond therefore H₃PO₂ is a stronger reducing agent than H₃PO₃.



(b) (i) Benzoic acid being an acid reacts with NaHCO₃ solution to produce brisk effervescence due to evolution of CO₂ while ethyl benzoate does not give this test.

(ii) Propanal being an aldehyde reduces Tollens' reagent to silver mirror but butan-2-one being a ketone does not.

OR


(a) (i) The methyl group due to its +I effect reduce the magnitude of positive charge on carbonyl carbon atom. Moreover it also hinder the approach of nucleophile CN. Since in acetaldehyde there is one methyl while in acetone there are two methyl groups attached to carbonyl group therefore acetaldehyde is more reactive than acetone towards nucleophilic addition with HCN.

Semicarbazide has two $-\mathrm{NH}_2$ groups but one of them (*i.e.*, directly attached to C = O) is involved in resonance as shown above. Thus, electron density on this NH $_2$ group decreases hence it does not act as a nucleophile. In contrast, the lone pair of electrons on the other NH $_2$ group (i.e., attached to —NH) is not involved in resonance and hence is available for nucleophilic attack on the C O group of aldehydes and ketones.

0

(b) (i) Rosenmund reduction:

Butan-2-one



(iii) Cannizzaro reaction:

SET-II

- 1. Zone refining is based on the principle that the impurities are more soluble in melt than in the solid state of metal.
- 2. Kraft temperature
- 3. Order of basic character BiH₃ < SbH₃ < AsH₃ < PH₃ < NH₃
- 4. Ionisation isomerism
- **5.** Atomic radius (r) Edge length (a) $\frac{2\sqrt{2}}{2\sqrt{2}}$
- **6.** CH₃NH₂ is more basic than NH₃. CH₃ group due to its +ve I effect pushes electron towards nitrogen in CH₃ NH₂ and this makes the unshared electron pair more available for sharing with the proton of the acid.
- 7. Chloroprene (CH₂ C CH CH₂
- **9.** (*i*) Impure nickel when heated in a current of CO forms volatile complex nickel tetracarbonyl, Ni(CO)₄ leaving behind impurities.

The nickel tetracarbonyl complex thus obtained is then heated to a higher temperature so that it is decomposed to give pure metal.

$$Ni(CO)_4$$
 450 – 470 K Ni + 4CO

(ii) Graphite rod acts as anode and graphite lined iron acts as cathode in the electrometallurgy of aluminium.

Carbon reacts with oxygen liberated at anode producing CO and CO_2 otherwise oxygen liberated at the anode may oxidise some of the liberated aluminium back to Al_2O_3 .

At Anode:
$$C(s) + O^2$$
 (melt) $CO(g) + 2e$

$$C(s) + 2O^2$$
 (melt) $CO_2(g) + 4e$
At Cathode: Al^{3+} (melt) $+ 3e$ $Al(l)$

10. (i)
$$2MnO_4^- + 5NO_2^- + 6H^+$$
 $2Mn^{2+} + 5NO_3^- + 3H_2O$ (ii) $Cr_2O_7^{2-} + 14H^+ + 6e^ 2Cr^{3+} + 7H_2O$

11. $[Ni(CN)_4]^{2-}$: Tetracyanonickelate (II) ion, Ni^{2+} (3d⁸)

$$3d \qquad 4s \qquad 4p$$

$$\uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \boxed{XX} \qquad \boxed{XX} \qquad \boxed{XX} \boxed{XX} \boxed{XX} \boxed{XX}$$

$$XX = Electron pair from CN^- ion$$

 dsp^2 hybridisation in [Ni (CN)₄]²⁻ leads to square planar shape.

- 13. (i) Conc. HNO₃/conc. H₂SO₄
 - (ii) Lithium aluminium hydride (LiAlH₄) or sodium borohydride (NaBH₄)
 - (iii) Acetyl chloride (CH₃COCl) in the presence of catalyst anhydrous AlCl₃.
 - (iv) Pyridinium chlorochromate, PCC (C₅H₅ N HCrO₃Cl in CH₂Cl₂).
- **20.** (*i*) Carbohydrates that yield a large number of monosaccharide units on hydrolysis are called polysaccharides *e.g.*, starch, cellulose, gums etc.
 - (ii) Amino acids are the compounds, whose molecule contains both the carboxylic acid (-COOH) group and the amino (-NH₂) group of the various amino acids, the -amino acids are most important because they are the building blocks of proteins.
 - (iii) **Enzyme:** Enzymes are complex nitrogenous organic compounds produced in living cells of plants and animals. Chemically, enzymes are globular proteins with high molar mass ranging from 15,000 to 1,000,000 g mol⁻¹.

SET-III

- 1. Sodium cyanide (NaCN)
- 2. Associated colloids
- 4. Linkage isomerism
- **5.** Propane–1, 2, 3-triol

$$NH_2$$
 is stronger base than as CH_3 group is electron releasing by +I effect and hyperconjugation effect.

- 7. Enzymes are termed as biocatalysts as they catalyse numerous reactions that occur in the bodies of animals and plants to maintain life process e.g., invertase, pepsin, urease.
- **9.** (*i*) When crude zirconium is heated in an evacuated vessel with iodine, volatile ZrI₄ is formed which vapourises leaving behind impurities. It is then decomposed by heating over a tungsten filament at 2075K to give pure zirconium.


(ii) Copper matte chiefly consist of Cu₂S and FeS. In the converter FeS gets converted into FeO. Silica (SiO₂) acts as a flux to remove FeO impurity as FeSiO₃ slag.

$$2FeS + 3O_2 2FeO + 2SO_2$$

11 (i)
$$2MnO_4$$
 $5SO_3^{2-}$ 6H $2Mn^{2+} + 5SO_4^{2-} + 3H_2O$

(ii)
$$2CrO_4^2$$
 2H $Cr_2O_7^{2-} + H_2O$

12. $[Co(NH_3)_6]^{3+}$: Hexaamminecobalt (III), $Co^{3+}(3d^6)$



 d^2sp^3 hybridisation in $[Co(NH_3)_6]^{3+}$ leads to octahedral shape.


13. (i)
$$CH_3CH_2Cl$$
 + alc. KOH $CH_2 = CH_2 + KCl + H_2O$
Ethyl chloride Ethene

(ii) + CH_3Cl Anhyd. $AlCl_3$ + CH_3Cl Chlorobenzene

1-chloro-2-methylbenzene
(Minor)

1-chloro-4-methylbenzene

- **15.** (*i*) Zinc dust
 - (ii) 85% H₂SO₄ at 440 K
 - (iv) Heat copper at 573 K (Cu/573 K)
- **24.** (a) Sucrose is dextrorotatory (+66.5°) but after hydrolysis it gives an equimolar mixture of D-(+)-glucose and D-(-)-fructose, which is laevorotatory. This change of specific rotation from dextrorotation to laevorotation is called inversion of sugar and the mixture obtained is called invert sugar.
 - (b) Vitamins are generally regarded as organic compounds required in the diet in small amounts to perform specific biological functions for normal maintenance of optimum growth and health of the organism.
 - (c) Nucleoside is a unit formed by the combination of nitrogen containing heterocyclic base and pentose sugar.



(Major)

CBSE Examination Paper Delhi-2015 (Code No. 56/1/1/D)

Time allowed: 3 hours Maximum marks: 70

General Instructions:

- (i) All questions are compulsory.
- (ii) Question numbers 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Question numbers 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Question numbers 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Question number 23 is a value-based question and carries 4 marks.
- (vi) Question numbers 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables, if necessary. Use of calculators is not allowed.
 - 1. What is the basicity of H_3PO_4 ?
 - **2.** Write the IUPAC name of the given compound:

3. Which would undergo S_N2 reaction faster in the following pair and why?

$$CH_3$$
 $CH_3 - CH_2 - Br$ and $CH_3 - C - CH_3$
 Br

- **4.** Out of BaCl₂ and KCl, which one is more effective in causing coagulation of a negatively charged colloidal sol? Give reason.
- **5.** What is the formula of a compound in which the element Y forms ccp lattice and atoms of X occupy 1/3rd of tetrahedral voids?
- **6.** What are the transition elements? Write two characteristics of the transition elements.
- 7. (i) Write down the IUPAC name of the following complex:

 $[Cr(NH_3)_2Cl_2(en)]Cl$ (en = ethylenediamine)

(ii) Write the formula for the following complex: Pentaamminenitrito-o-Cobalt (III)

8. Name the reagents used in the following reactions:

(ii)
$$C_6H_5 - CH_2 - CH_3$$
 ? $C_6H_5 - COO^- K^+$

9. What is meant by positive deviations from Raoult's law? Give an example. What is the sign of $\min H$ for positive deviation?

OR

Define azeotropes. What type of azeotrope is formed by positive deviation from Raoult's law? Give an example.

$$Ag^{+}(aq) + e^{-}$$
 $Ag(s)$ $E^{\circ} = +0.80 \text{ V}$
 $H^{+}(aq) + e^{-}$ $\frac{1}{2}H_{2}(g)$ $E^{\circ} = 0.00 \text{ V}$

On the basis of their standard reduction electrode potential (E°) values, which reaction is feasible at the cathode and why?

- (b) Define limiting molar conductivity. Why conductivity of an electrolyte solution decreases with the decrease in concentration?
- 11. 3.9 g of benzoic acid dissolved in 49 g of benzene shows a depression in freezing point of 1.62 K. Calculate the van't Hoff factor and predict the nature of solute (associated or dissociated).

(Given: Molar mass of benzoic acid = 122 g mol⁻¹, K_f for benzene = 4.9 K kg mol⁻¹)

- **12.** (*i*) Indicate the principle behind the method used for the refining of zinc.
 - (ii) What is the role of silica in the extraction of copper?
 - (iii) Which form of the iron is the purest form of commercial iron?
- 13. An element with molar mass 27 g mol⁻¹ forms a cubic unit cell with edge length 4.05×10^{-8} cm. If its density is 2.7 g cm⁻³, what is the nature of the cubic unit cell?
- **14.** (a) How would you account for the following:
 - (i) Actinoid contraction is greater than lanthanoid contraction.
 - (ii) Transition metals form coloured compounds.
 - (b) Complete the following equation:

$$2MnO_{4}^{-} + 6H^{+} + 5NO_{2}^{-}$$

- 15. (i) Draw the geometrical isomers of complex $[Pt(NH_3)_2Cl_2]$.
 - (ii) On the basis of crystal field theory, write the electronic configuration for d^4 ion if 0 < P.
 - (iii) Write the hybridisation and magnetic behaviour of the complex $[Ni(CO)_4]$.

(At. no. of
$$Ni = 28$$
)

16. Calculate emf of the following cell at 25°C:

Fe | Fe²⁺(0.001 M) || H⁺(0.01 M) | H₂(g) (1 bar) | Pt(s)
E°(Fe²⁺ | Fe) =
$$-0.44$$
 V, E°(H⁺ | H₂) = 0.00 V

- **17.** Give reasons for the following observations:
 - (i) Leather gets hardened after tanning.
 - (ii) Lyophilic sol is more stable than lyophobic sol.
 - (iii) It is necessary to remove CO when ammonia is prepared by Haber's process.
- **18.** Write the names and structures of the monomers of the following polymers:
 - (i) Nylon-6, 6
- (ii) PHBV

- (iii) Neoprene
- **19.** Predict the products of the following reactions:

$$\begin{array}{ccc} \textit{(i)} \ \, \text{CH}_{3} & \xrightarrow{\text{C} = \text{O}} & \xrightarrow{\textit{(i)} \ \text{H}_{2}\text{N} - \text{NH}_{2}} \\ \text{CH}_{3} & \xrightarrow{\textit{(ii)} \ \text{KOH/Glycol,}} \, \, \, ? \end{array}$$

(ii)
$$C_6H_5$$
— CO — CH_3 $NaOH/I_2$? +?

- **20.** How do you convert the following:
 - (i) Phenol to anisole
 - (ii) Propan-2-ol to 2-methylpropan-2-ol
 - (iii) Aniline to phenol



OR

(a) Write the mechanism of the following reaction:

$$2CH_3CH_2OH$$
 H^+ $CH_3CH_2 - O - CH_2CH_3$

- (b) Write the equation involved in the acetylation of Salicylic acid.
- **21.** (i) Which one of the following is a disaccharide: Starch, Maltose, Fructose, Glucose?
 - (ii) What is the difference between fibrous protein and globular protein?
 - (iii) Write the name of vitamin whose deficiency causes bone deformities in children.
- **22.** Give reasons:
 - (a) n-Butyl bromide has higher boiling point than t-butyl bromide.
 - (b) Racemic mixture is optically inactive.
 - (c) The presence of nitro group (-NO₂) at o/p positions increases the reactivity of haloarenes towards nucleophilic substitution reactions.
- 23. Mr. Roy, the principal of one reputed school organized a seminar in which he invited parents and principals to discuss the serious issue of diabetes and depression in students. They all resolved this issue by strictly banning the junk food in schools and to introduce healthy snacks and drinks like soup, lassi, milk etc., in school canteens. They also decided to make compulsory half an hour physical activities for the students in the morning assembly daily. After six months, Mr. Roy conducted the health survey in most of the schools and discovered a tremendous improvement in the health of students.


After reading the above passage, answer the following:

- (i) What are the values (at least two) displayed by Mr. Roy?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) What are tranquilizers? Give an example.
- (iv) Why is use of aspartame limited to cold foods and drinks?
- **24.** (a) Account for the following:
 - (i) Acidic character increases from HF to HI.
 - (ii) There is large difference between the melting and boiling points of oxygen and sulphur.
 - (iii) Nitrogen does not form pentahalide.
 - (b) Draw the structures of the following:

$$(i)$$
 CIF₃ (ii) X

OR

- (i) Which allotrope of phosphorus is more reactive and why?
- (ii) How the supersonic jet aeroplanes are responsible for the depletion of ozone layers?
- (iii) F₂ has lower bond dissociation enthalpy than Cl₂. Why?
- (iv) Which noble gas is used in filling balloons for meteorological observations?
- (v) Complete the equation: $XeF_2 + PF_5$
- 25. An aromatic compound 'A' of molecular formula C₇H₇ON undergoes a series of reactions as shown below. Write the structures of A, B, C, D and E in the following reactions:





OR

- (a) Write the structures of main products when aniline reacts with the following reagents:
 - (i) Br₂ water

- (ii) HCl
- (iii) (CH₃CO)₂O / pyridine
- (b) Arrange the following in the increasing order of their boiling point:

C₂H₅NH₂, C₂H₅OH, (CH₃)₃N

- (c) Give a simple chemical test to distinguish between the following pair of compounds: (CH₃)₂NH and (CH₃)₃N
- **26.** For the hydrolysis of methyl acetate in aqueous solution, the following results were obtained:

t/s	0	30	60
[CH ₃ COOCH ₃]/mol L ⁻¹	0.60	0.30	0.15

- (i) Show that it follows pseudo first order reaction, as the concentration of water remains constant.
- (ii) Calculate the average rate of reaction between the time interval 30 to 60 seconds.

(Given $\log 2 = 0.3010$, $\log 4 = 0.6021$)

OR

(a) For a reaction A + B

P, the rate is given by

Rate = $k[A][B]^2$

- (i) How is the rate of reaction affected if the concentration of B is doubled?
- (ii) What is the overall order of reaction if A is present in large excess?
- (b) A first order reaction takes 30 minutes for 50% completion. Calculate the time required for 90% completion of this reaction.


 $(\log 2 = 0.3010)$





Solutions •

1. Tribasic, as the structure of H₃PO₄ has three P—OH bonds.



- **2.** 2, 5-Dinitrophenol.
- Br. Being a primary halide, there will be less crowding around -carbon. CH_2
- **4.** BaCl₂, Ba²⁺ ion has greater coagulating power than K ion as Ba²⁺ ion has higher charge.
- Let the number of atoms of Y in ccp arrangement = N

Number of tetrahedral voids = 2N

Number of
$$X$$
 atoms $\frac{1}{3}$ 2N
 $X:Y=\frac{2}{3}$ N: N
 $X:Y=2:3$

So, the formula of the compound is X_2Y_3 .

Transition elements are those elements which have partly filled d-orbitals in their ground states or any of their common oxidation states.

Two characteristics of transition elements are:

- (i) Transition elements show variable oxidation states.
- (ii) Most of the transition elements form coloured compounds.
- 7. (i) Diamminedichloridoethane-1, 2-diaminechromium (III) chloride.
 - (ii) $[Co(NH_3)_5(ONO)]^{2+}$
- (i) Sodium borohydride (NaBH₄) or Lithium aluminium hydride (LiAlH₄).
 - (ii) Alkaline potassium permanganate (KMnO₄ KOH)
- The deviation is said to be positive, if the vapour pressure of the solution is greater than the pressure expected on the basis of Raoult's law for the same composition. Mixtures of ethanol and acetone behave in this manner. For such a solution $_{mix}H$ 0, i. e., positive.

OR

Azeotropes are binary liquid mixtures having the same composition in liquid and vapour phase and boil at a constant temperature. The solution which shows positive deviation from Raoult's law forms minimum boiling azeotrope at a specific composition. For example, a mixture of ethanol and water containing 95% ethanol by volume.

- **10.** (a) The reaction, $Ag^+(aq) + e$ Ag (s) is feasible at cathode as cathodic reaction is one which has higher standard reduction electrode potential (E_{red}).
 - (b) The value of molar conductivity when the concentration approaches zero is known as limiting molar conductivity. Conductivity of an electrolytic solution decreases with decrease in concentration as the number of ions per unit volume that carry the current in the solution decreases with decrease in concentration.



11. The given quantities are

$$W_B$$
 3.9 g, W_A 49 g, T_f 1.62 K, M_B 122 g mol 1 and K_f 4.9 K kg mol 1

Substituting these values in the equation $M_B = \frac{K_f - W_B - 1000}{T_f - W_A}$, we get

$$M_B = \frac{4.9 \text{ K kg mol}^{-1} \quad 3.9 \text{ g}^{-1000 \text{ g kg}^{-1}}}{1.62 \text{ K}^{-49 \text{ g}}}$$

$$M_B = 240.74 \text{ g mol}^{-1}$$

Thus, observed molecular mass of benzoic acid in benzene = 240.74 g mol 1

Normal molecular mass of benzoic acid 122 g mol 1

$$i = \frac{\text{Normal molecular mass}}{\text{Observed molecular mass}}$$

$$i = \frac{122 \text{ g mol}^{-1}}{240.74 \text{ g mol}^{-1}} = 0.506$$

As i < 1, therefore, benzoic acid will undergo association in benzene.

- **12.** (*i*) Distillation, as zinc is a low boiling metal.
 - (ii) Silica (SiO₂) acts as a flux to remove iron oxide (FeO) impurity as slag FeSiO₃ in the metallurgy of copper. FeS present in the copper matte is oxidised to FeO which combine with silica (SiO2) to form FeSiO₃.

$$\begin{array}{ccc} \text{2FeS} + 3\text{O}_2 & \text{2FeO} + 2\text{SO}_2 \\ \text{FeO} & + \text{SiO}_2 & \text{FeSiO}_3 \\ \text{Impurity} & \text{Flux} & \text{Slag} \end{array}$$

(iii) Wrought iron is the purest form of commercial iron.

13. Number of atoms per unit cell,
$$z = \frac{d - a^3 - N_A}{M}$$
 ...(i)

Here, $d=2.7 \text{ g cm}^{-3}$, $M=27 \text{ g mol}^{-1}$, $N_A=6.022 \cdot 10^{23} \cdot \text{mol}^{-1}$ and $a=4.05 \cdot 10^{-8} \cdot \text{cm}$

Substituting these values in the expression (i), we get


$$z = \frac{2.7 \text{ g cm}^{-3} + (4.05 + 10^{-8} \text{ cm})^3 + 6.022 + 10^{23} \text{ mol}^{-1}}{27 \text{ g mol}^{-1}}$$


As there are 4 atoms of the elements present per unit cell therefore the cubic unit cell must be face-centered.

- 14. (a) (i) This is because 5f electrons provide more poor shielding effect than 4f electrons as 5f orbitals are more diffused than 4f orbitals.
 - (ii) This is due to d-d transition of electrons occurring in transition metal ions by absorption of visible light.

(b)
$$\frac{\text{MnO}_4 + 8\text{H}^+ + 5e}{\text{NO}_2 + \text{H}_2\text{O}} \frac{\text{Mn}^{2+} + 4\text{H}_2\text{O}}{\text{NO}_3 + 2\text{H}^+ + 2e} \frac{1}{2} \frac{5}{2\text{MnO}_4 + 5\text{NO}_2 + 6\text{H}^+} \frac{2\text{Mn}^{2+} + 5\text{NO}_3 + 3\text{H}_2\text{O}}{2\text{Mn}^{2+} + 5\text{NO}_3 + 3\text{H}_2\text{O}}$$








Geometrical isomers of [Pt(NH₃)₂Cl₂]

- P, then fourth electron enters one of the e_g orbitals giving the configuration $t_{2g}^3 e_g^1$.
- (iii) The complex [Ni(CO)₄] involves sp³ hybridisation.



The complex is diamagnetic as evident from the absence of unpaired electrons. Fe^{2+} 2e

At cathode: $\frac{\text{Fe} + 2\text{H}^+ \qquad \text{Fe}^{2+} + \text{H}_2}{\text{Fe}^{2+} + \text{H}_2}$

$$E_{cell}$$
 E_{cell} $\frac{0.059}{n} \log \frac{[\text{Fe}^2]}{[\text{H}]^2}$...(i)

 E_{cell} $E_{cathode}$ E_{anode} $E_{
m H~/H_2}$ $E_{
m Fe^{2^+/Fe}}$ Here

$$E_{cell}$$
 0.00 V - (- 0.44 V) = 0.44 V

Substituting [Fe²] 1 10 ³ M, [H⁺] = 1 10 ² M, n 2 and E_{cell} 0.44 V in expression (i), we get

$$E_{cell}$$
 0.44 $\frac{0.059}{2} \log \frac{10^{-3}}{(10^{-2})^2}$
 E_{cell} 0.44 0.0295 $\log 10$ 0.44 - 0.0295
 E_{cell} 0.4105 V = 0.41 V

- (i) Animal hide is colloidal in nature and contains positively charged particles. When a hide is soaked in a **17.** tannin which contain negatively charged colloidal particles, mutual coagulation takes place and leather becomes hard.
 - (ii) As the stability of a lyophobic sol is only due to the presence of charge on the colloidal particles whereas the stability of lyophilic sol is due to the two factors, the charge and the solvation of colloidal particles.
 - (iii) As CO acts as a catalytic poison.




18.

	Polymers	Monomer Structures	Monomer Names
(i)	Nylon-6, 6	$H_2N(CH_2)_6NH_2$	Hexamethylene diamine
		HOOC—(CH ₂) ₄ COOH	Adipic acid
(ii)	PHBV	CH ₃ CH(OH)CH ₂ COOH	3-hydroxybutanoic acid
		CH ₃ CH ₂ CH(OH)CH ₂ COOH	3-hydroxypentanoic acid
		Cl	
(iii)	Neoprene	H ₂ C C CH CH ₂	Chloroprene

- CH₃ (Propane) **19.** (*i*) CH₃ CH₂
 - (ii) C₆H₅COO Na + (Sodium benzoate) and CHI₃ (Iodoform)
 - (iii) CH₄ (Methane)

20. (i)
$$\stackrel{-+}{\longrightarrow}$$
 $\stackrel{-+}{\longrightarrow}$ $\stackrel{--}{\longrightarrow}$ \stackrel



- (a) Mechanism:
 - (i) Protonation of ethanol

(ii) Nucleophilic (S_N 2) attack of ethanol molecule on oxonium ion



(iii) Deprotonation

21. (*i*) Maltose is a disaccharide.

(ii)

S.No.	Fibrous Proteins	Globular Proteins
(i)		This structure results when the chains of polypeptides coil around to give spherical shape, <i>e.g.</i> , insulin, albumin, etc.
(ii)	These are insoluble in water.	These are usually soluble in water.

- (iii) Vitamin D
- 22. (i) n-Butyl bromide being a straight chain alkyl halide has larger surface area than tert.butyl bromide. Larger the surface area larger the magnitude of the van der Waal's forces and hence higher is the boiling point.
 - (ii) A racemic mixture contains the two enantiomers d and l in equal proportions. As the rotation due to one enantiomer is cancelled by equal and opposite rotation of another enantiomer therefore, it is optically
 - (iii) The presence of NO₂ group at o/p position in haloarenes helps in the stabilisation of resulting carbanion by -R and -I effects and hence increases the reactivity of haloarenes towards nucleophilic substitution reactions.
- **23.** (*i*) Values displayed by Mr. Roy:
 - Caring in nature
 - Concern for student's health
 - Use of knowledge of chemistry in daily life.
 - (ii) Awareness for health can be spread by:
 - performing play in school assembly
 - performing *nukkad natak* in community
 - displaying posters/slogans
 - conducting seminars
 - (iii) Tranquilizers are a class of chemical compounds used for the treatment of stress and mild or even severe mental diseases. These relieve stress, anxiety, irritability or excitement by inducing a sense of well-being, e.g., meprobamate, equanil, luminal, etc.
 - (iv) Because aspartame is unstable at cooking temperature.
- 24. (a) (i) As the size of halogen atom increases from F to I, the bond dissociation enthalpy of H—X bond decreases from H—F to H—I. Due to this acidic character increases from HF to HI.
 - (ii) Because of small size and high electronegativity oxygen forms p -p multiple bonds and exists as a diatomic, O2 molecule. These molecules are held together by weak vander Waal forces. Sulphur on




the other hand due to its higher tendency for catenation and lower tendency for p -p multiple bond formation, forms octa-atomic, S₈ molecule. Because of bigger size of S₈ molecule than O₂ molecule the force of attraction holding the S₈ molecules together are much stronger than O₂ molecules. Hence, there is large difference between the melting and boiing points of oxygen and sulphur.

(iii)Nitrogen with n 2, has s and p-orbitals only. It does not have d-orbitals to expand its covalence beyond four. Due to this it does form pentahalide.





CIF₃: Slightly bent "T".



XeF₄: Square planar.

OR

- (i) White or yellow phosphorus is more reactive than the other allotropes because it is less stable due to angular strain in the P₄ molecule of white phosphorus where the angles are only 60°.
- (ii) Nitrogen oxides (particularly nitric oxide) emitted from the exhaust system of supersonic jet aeroplanes are responsible for depletion of ozone layer.

$$NO(g) + O_3(g)$$
 $NO_2(g) + O_2(g)$

- (iii) Bond dissociation enthalpy of F2 is lower than Cl2 due to small size of fluorine and relatively larger electron-electron repulsion among the lone pairs in F2 molecule where they are much closer to each other than in case of Cl₂.
- (iv) Helium, as it is non-inflammable and light gas.

$$(v) \text{ XeF}_2 + \text{PF}_5$$

$$[XeF] [PF_6]$$

25.

$$A = \bigcup_{\substack{C - NH_2 \\ Benzamide}} ; B = \bigcup_{\substack{+ \\ N \\ Benzene \\ diazonium \\ chloride}} ; C = \bigcup_{\substack{+ \\ N \\ Benzene}} ; D = \bigcup_{\substack{+ \\ N \\ Benzene \\ isocyanide}} ; E = \bigcup_{\substack{+ \\ Iodobenzene \\ isocyanide}}$$

(a) (i)
$$+ 3Br_2(water) \longrightarrow$$

Aniline

OR
$$\begin{array}{c} NH_2 \\ Br \\ Br \\ \end{array} + 3HBr$$

2,4,6-Tribromo aniline

$$(ii)$$
 NH_2 NH_3 $C\bar{l}$

Aniline NH_3 NH_3



(iii)
$$\begin{array}{c} NH_2 \\ CH_3 - C \\ CH_3 - C \\ O \end{array}$$
 $\begin{array}{c} O \\ Pyridine \\ O \end{array}$ $\begin{array}{c} O \\ NH - C - CH_3 \\ O \\ Acetanilide \\ O \end{array}$ $\begin{array}{c} O \\ CH_3 - C - OH \\ O \end{array}$

Acetic anhydride

- (b) $(CH_3)_3 N < C_2 H_5 NH_2 < C_2 H_5 OH$
- (c) Dimethyl amine and trimethyl amine can be distinguished by using Hinsberg's reagent, i.e., C₆H₅SO₂Cl. When treated with Hinsberg's reagent dimethylamine gives N, N-dimethyl benzene sulphonamide which is insoluble in aqueous KOH solution while trimethyl amine does not react with Hinsberg's reagent.

26. (i)
$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$
 ...(i)

Substituting $[R]_0$ 0.60 mol L¹, [R] 0.30 mol L¹ and t = 30 s in equation (i), we get

$$k = \frac{2303}{30} \log \frac{0.60}{030}$$

$$k = \frac{2.303}{30} \log 2 = \frac{2303}{30} = 03010$$

$$k = 0.0231 \text{ s}^{-1}$$

0.60 mol L¹, [R] 0.15 mol L¹ and t 60 s in equation (i), we get Again substituting, $[R]_0$

$$k = \frac{2303}{60} \log \frac{0.60}{0.15}$$

$$k = \frac{2303}{60} \log 4 = \frac{2303}{60} = 0.6021$$

$$k = 0.0231 \text{ s}^{-1}$$

As the value of k is same in both the cases, therefore, hydrolysis of methylacetate in aqueous solution follows pseudo first order reaction.

(ii) Average rate
$$\frac{ [CH_3COOCH_3]}{t}$$

$$\frac{ [0.15 \quad 0.30]}{60 \quad 30} \frac{0.15}{30}$$

Average rate 0.005 mol L ¹ s ¹



(a) (i) Rate,
$$r = k [A] [B]^2$$

If concentration of B is doubled, then

$$r = k [A] [2B]^2$$

r 4k [A] [B]², i.e., the rate will increase by 4 times.

(ii) As A is present in large excess therefore, order of the reaction w.r.t. A will be zero.

Rate
$$k [A]^0 [B]^2 k [B]^2$$

Order of reaction 2

(b) $t_{1/2}$ 30 min.

[R] [R]₀ - 90% of [R]₀ [R]₀
$$\frac{90 [R]_0}{100}$$

[R] $\frac{[R]_0}{10}$
 $k = \frac{0.693}{t_{1/2}} = \frac{0.693}{30} = 0.0231 \text{ min}^{-1}$
 $t = \frac{2303}{k} \log \frac{[R]_0}{[R]}$
 $t = \frac{2303}{0.0231} \log \frac{[R]_0}{[R]_0} = \frac{2303}{0.0231} \log 10$

99.7 min

CBSE Examination Paper Ajmer Region-2015 (Code No. 56/1/A)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination paper, Delhi-2015.

1. Out of AlCl₃ and NaCl, which is more effective in causing coagulation of a negative sol and why?

- 2. Write the formula of a compound in which the element Y forms *ccp* lattice and atoms of X occupy 1/3rd of tetrahedral voids.
- **3.** Write the formulae of any two oxoacids of phosphorus.
- **4.** Write the IUPAC name of the given compound:

$$CH_3$$
 $CH_3 - C - CH_2 - OH$
 CH_3

5. Which would undergo S_N 2 reaction faster in the following pair:

$$C_6H_5 - CH_2 - CH_2 - Br$$
 and $C_6H_5 - CH - CH_3$

Br

- **6.** (i) Why are aquatic species more comfortable in cold water than in warm water?
 - (ii) What happens when we place the blood cell in saline water solution (hypertonic solution)? Give reason.
- 7. Calculate the time to deposit 1.5 g of silver at cathode when a current of 1.5 A was passed through the solution of AgNO₃. (Molar mass of Ag = 108 g mol^{-1} , $1 \text{ F} = 96500 \text{ C mol}^{-1}$)
- **8.** Why do transition elements show variable oxidation states? How is the variability in oxidation states of *d*-block different from that of the *p*-block elements?
- 9. (i) Write down the IUPAC name of the following complex:

$$[Pt(NH_3)(H_2O)Cl_2]$$

- (ii) Write the formula for the following complex: tris(ethane-1,2-diamine)chromium(III) chloride
- 10. Write the reagents used in the following reactions:

Arrange the following compounds in increasing order of their property as indicated:

- (i) CH₃CHO, C₆H₅CHO, HCHO (reactivity towards nucleophilic addition reaction)
- (ii) 2, 4-dinitrobenzoic acid, 4-methoxybenzoic acid, 4-nitrobenzoic acid (acidic character)
- 11. Vapour pressure of water at 20°C is 17.5 mm Hg. Calculate the vapour pressure of water at 20°C when 15 g of glucose (Molar mass = 180 g mol^{-1}) is dissolved in 150 g of water.
- 12. Examine the given defective crystal:

$$X^{+}$$
 Y^{-} X^{+} Y^{-} Y^{-}

- (i) Is the above defect stoichiometric or non-stoichiometric?
- (ii) Write the term used for the electron occupied site.
- (iii) Give an example of the compound which shows this type of defect.
- **13.** Calculate E_{cell}^0 and ${}_{\text{r}}G^0$ for the following reaction at 25°C:

$$A^2$$
 B A^3

$$A^3$$

Given:
$$K_c = 10^{10}$$
, 1F = 96500 C mol⁻¹

- 14. Define adsorption with an example. Why is adsorption exothermic in nature? Write the types of adsorption based on the nature of forces between adsorbate and adsorbent.
- **15.** (*i*) Name the method used for the refining of titanium.
 - (ii) What is the role of Zn in the extraction of silver?
 - (iii) Reduction of metal oxide to metal becomes easier if the metal obtained is in liquid state. Why?
- **16.** (i) E^0 value for the Mn^{3+}/Mn^{2+} couple is positive (+ 1.5 V) whereas that of Cr^{3+}/Cr^{2+} is negative (-0.4 V). Why?
 - (ii) Transition metals form coloured compounds. Why?
 - (iii) Complete the following equation:

$$2MnO_4^-$$
 16H $5C_2O_4^2$

- 17. (i) What type of isomerism is shown by [Co(NH₃)₅ONO]Cl₂?
 - (ii) On the basis of crystal field theory, write the electronic configuration for d^4 ion, if P.
 - (iii) Write the hybridization and shape of $[Fe(CN)_6]^{3-}$. (Atomic number of Fe = 26)
- **18.** How do you convert the following:
 - (i) Prop-l-ene to Propan-2-ol
 - (ii) Bromobenzene to 2-bromoacetophenone
 - (iii) 2-bromobutane to But-2-ene

What happens when

- (i) ethyl chloride is treated with NaI is the presence of acetone,
- (ii) chlorobenzene is treated with Na metal in the presence of dry ether,
- (iii) methyl chloride is treated with KNO₂?

Write chemical equations in support of your answer.

- **19.** Give reasons for the following:
 - (i) p-nitrophenol is more acidic than p-methylphenol.
 - (ii) Bond length of C–O bond in phenol is shorter than that in methanol.
 - (iii) (CH₃)₃C—Br on reaction with sodium methoxide (Na⁺-OCH₃) gives alkene as the main product and not an ether.
- **20.** Predict the products of the following reactions:

(ii)
$$C_6H_5 - CH_2 - CH_3 \frac{(a)KMnO_4 / KOH}{(b) H}$$
?

$$(iii) \begin{picture}(100) \put(0.05){\line(1,0){100}} \put(0.05){\line($$

- 21. Write the names and structures of the monomers of the following polymers:
 - (i) Nylon-6,6

- (ii) Bakelite
- (iii) Polystyrene



- **22.** (i) Which one of the following is a disaccharide:
 - starch, maltose, fructose, glucose
 - (ii) What is the difference between acidic amino acid and basic amino acid?
 - (iii) Write the name of the linkage joining two nucleotides.
- 23. Seeing the growing cases of diabetes and depression among young children, Mr. Lugani, the principal of one reputed school organised a seminar in which he invited parents and principals. They all resolved this issue by strictly banning junk food in schools and introducing healthy snacks and drinks like soup, *lassi*, milk, etc. in school canteens. They also decided to make compulsory half an hour of daily physical activities for the students in the morning assembly. After six months, Mr. Lugani conducted the health survey in most of the schools and discovered a tremendous improvement in the health of the students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Lugani?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) What are antidepressant drugs? Give an example.
- (iv) Name the sweetening agent used in the preparation of sweets for a diabetic patient.
- **24.** An aromatic compound 'A' of molecular formula C_7H_7ON undergoes a series of reactions as shown below. Write the structures of A, B, C, D and E in the following reactions:

OR

- (a) Write the structures of the main products when aniline reacts with the following reagents:
 - (i) Br₂ water

(ii) HCl

- (iii) (CH₃ CO)₂O/pyridine
- (b) Arrange the following in the increasing order of their boiling point:

(c) Give a simple chemical test to distinguish between the following pair of compounds:

$$(CH_3)_2$$
—NH and $(CH_3)_3$ N

25. For the hydrolysis of methyl acetate in aqueous solution, the following results were obtained:

t/s	0	30	60
[CH ₃ COOCH ₃]/mol L ⁻¹	0.60	0.30	0.15

- (i) Show that it follows pseudo first order reaction, as the concentration of water remains constant.
- (ii) Calculate the average rate of reaction between the time interval 30 to 60 seconds.

OR

(a) For a reaction A + B P, the rate is given by

Rate =
$$k [A]^2 [B]$$

- (i) How is the rate of reaction affected if the concentration of A is doubled?
- (ii) What is the overall order of reaction if B is present in large excess?
- (b) A first order reaction takes 23.1 minutes for 50% completion. Calculate the time required for 75% completion of this reaction.

(Given:
$$\log 2 = 0.301$$
, $\log 3 = 0.4771$, $\log 4 = 0.6021$)



- **26.** (a) Account for the following:
 - (i) Bond angle in NH₄ is greater than that in NH₃.
 - (ii) Reducing character decreases from SO₂ to TeO₂.
 - (iii) HClO₄ is a stronger acid than HClO.
 - (b) Draw the structures of the following:
 - (i) H₂S₂O₈
 - (ii) XeOF₄

OR

- (a) Which poisonous gas is evolved when white phosphorus is heated with conc. NaOH solution? Write the chemical equation.
- (b) Write the formula of first noble gas compound prepared by N. Bartlett. What inspired N. Bartlett to prepare this compound?
- (c) Fluorine is a stronger oxidising agent than chlorine. Why?
- (d) Write one use of chlorine gas.
- (e) Complete the following equation:

$$CaF_2 + H_2SO_4$$





- 1. AlCl₃, as charge on Al³ ion is greater than Na ion.
- **2.** Let the number of atom of element Y in ccp lattice = N

Number of tetrahedral voids 2 N

Number of atoms of element $X = \frac{1}{3}$ 2N $\frac{2}{3}$ N

$$X: Y = \frac{2}{3} N: N = 2:3$$

Formula of the compound X_2Y_3

- **3.** H₃PO₂, H₃PO₃, H₃PO₄, H₄P₂O₇
- **4.** 2, 2-Dimethylpropan-1-ol.
- **5.** C_6H_5 CH_2 CH_2 Br, due to less crowding at -carbon atom.
- **6.** (i) As solubility of a gas in water decreases with increase in temperature therefore less oxygen is available in warm water than cold water.
 - (ii) Blood cells will shrink. Due to osmosis water will flow out of the blood cell and they would shrink.
- 7. Substituting W 1.5 g, M 108 g mol 1 , n 1, F 96500 C mol 1 , I = 1.5 amp in the expression $W = \frac{M}{n} \frac{I}{F}$, we get

$$1.5 = \frac{108 \quad 1.5 \quad t}{1 \quad 96500}, t \quad \frac{1.5 \quad 96500}{108 \quad 1.5} \quad 893.52 \text{ s}$$
$$t \quad \frac{893.52}{60} \quad 14.89 \text{ min}$$

or

8. In transition elements, the energies of (*n* 1)*d* orbitals and *ns* orbitals are nearly same. Therefore, electrons from both can participate in bond formation and hence show variable oxidation states.

In transition elements, the oxidation states differ from each other by unity, e.g., Fe² and Fe³, Cu and Cu², etc. while in p-block elements the oxidation state differ by units of two, e.g., Sn² and Sn⁴, Pb² and Pb⁴, etc. In transition elements the higher oxidation states are more stable for heavier elements in a group e.g., Mo(VI) and W(VI) are more stable than Cr(VI) in group 6 whereas in p-block elements the lower oxidation states are more stable for heavier elements due to the inert pair effect, e.g., Pb(II) is more stable than Pb (IV) in group 16.

- **9.** (*i*) Ammineaquadichloridoplatinum (II)
 - (ii) [Cr(en)₃]Cl₃
- 10. (i)Zn—Hg, conc. HCl or H₂NNH₂ and KOH/ethylene glycol, Heat
 - (ii) PCl₅ or SOCl₂

OR

- (i) C₆H₅CHO CH₃CHO HCHO
- (ii) 4-methoxybenzoic acid < 4-nitrobenzoic acid < 2,4-dinitrobenzoic acid.
- **11.** Substituting p_A 17.5 mm Hg, W_B 15 g, M_B 180 g mol 1 , M_A 18 g mol 1 , W_A 150 g in the

expression

$$\frac{p_A}{p_A} = \frac{W_B}{M_B} = \frac{M_A}{W_A}$$
, we get $\frac{17.5 - P}{17.5} = \frac{15}{180} = 0.01$

$$17.5 - P = 0.175$$
 or $P = 17.5 - 0.175 = 17.325$ mm Hg



- 12. (i) Non-stoichiometric defect
 - (ii) F-centre
 - (iii) When crystals of NaCl are heated in an atmosphere of Na vapours or when crystals of KCl are heated in an atmosphere of K vapours.
- 13. A^2 A^3 B; n = 1

Substituting $K_c = 10^{10}$, 1 F = 96500 C mol⁻¹, T = (25 - 273)K = 298 K, R = 8.314 JK ⁻¹ mol⁻¹ in $G = -2.303 \text{ RT log K}_c$, we get $G = -2.303 \times 8.314 \text{ J K}^{-1} \text{ mol}^{-1} 298 \text{ K} \times \log 10^{10}$

 $G = -57058.48 \text{ J mol}^{-1} = -57.058 \text{ kJ mol}^{-1}$

57058.48 J mol 1 , F = 96500 C mol 1 , n 1, in the expression Substituting *G*

$$G = -nFE_{\text{cell}}^{\text{o}}$$
, we get
 $-57058.48 = -1 \times 96500 E_{\text{cell}}^{\text{o}}$
 $E_{\text{cell}}^{\text{o}} = \frac{57058.48}{96500} = 0.591 \text{ V}$

14. The accumulation of the molecular species at the surface rather than in the bulk of a solid or liquid is known as adsorption. For example, water vapour are adsorbed by silica gel.

When a gas is adsorbed on the surface of a solid its entropy decreases i.e., S becomes -ve. Since adsorption is a spontaneous process therefore G(HTS) must be negative. As TS is +ve, G can be negative only if H has sufficiently high –ve value. Hence, adsorption is exothermic in nature.

There are two types of adsorption based on the nature of forces between adsorbate and adsorbent.

- (i) Physical adsorption, when accumulation of gas on the surface of a solid occurs due to weak van der Waal forces.
- (ii) Chemical adsorption, when the gas molecules or atoms are held to the surface of solid by chemical bonds.
- **15.** (*i*) van Arkel method (vapour phase refining)
 - (ii) Zn reduces [Ag(CN)₂] formed into Ag. $2[Ag(CN)_2]$ (aq) Zn(s) 2Ag(s) $[Zn(CN)_4]^2$ (aq)
 - (iii) Refer to Ans. 15 (iii) Panchkula Region 2015.
- 16. (i) The positive E value for the couple Mn^3 / Mn^2 is due to much higher third ionisation enthalpy of Mn, where the change is d^5 (extra stable) to d^4 . As Cr^3 (t_{2g}^3) is more stable than Cr^{2+} , therefore, E^0 value for the couple Cr^{3+}/Cr^{2+} is negative.
 - (ii) Refer to Ans. 26 (OR) (a) (i) Chennai Region 2015.
 - $Mn^2 4H_2O] 2$ 8H (iii) [MnO₄
- 17. (i) Linkage isomerism and the linkage isomer is $[Co(NH_3)_5 NO_2]Cl_2$.
 - (ii) If $_0$ P, the fourth electron enters one of two e_g orbitals giving the configuration $t_{2g}^3 e_g^1$.
 - (*iii*) $Fe^3 : 3d^5 4s^0$

3d(Inner orbital complex)

Hybridisation = d^2sp^3 , shape = octahedral



(i)
$$CH_3$$
 CH_2 Cl NaI CH_3 CH_2 I $NaCl$ $Ethyl chloride$ CH_3 CH_2 I CH_3 CH_2 CH_3 CH_4 CH_5 CH

- (iii) CH₃ Cl KNO₂ CH_3O N=O**KCl** Methyl chloride Methyl nitrite
- 19. (i) Due to -I and -R effect of NO₂ group electron density in the O—H bond decreases in p-nitrophenol whereas due to hyperconjugation effect and +I effect of CH₃ group, the electron density in the O—H bond increases in p-methylphenol. Consequently O—H bond in p-nitrophenol becomes much weaker than O—H bond in p-methyl phenol and hence p-nitrophenol is more acidic than p-methylphenol.
 - (ii) This is due to:
 - (a) partial double bond character as a result of the conjugation of unshared electron pair of oxygen with the aromatic ring and
 - (b) sp^2 hybridised state of carbon to which oxygen is attached.
 - (iii) Refer to Ans. 11 (c) Chennai Region 2015.
- **20.** (*i*) CH₃ C N NHCONH₂

 CH_3 Propanone semicarbazone

(ii) C₆H₅COOH Benzoic acid

(iii)
$$NO_2$$
 m-nitro benzoic acid



21.

	Name of polymer	Name of monomer	Structure of monomer
(<i>i</i>)	Nylon-6, 6	Hexamethylenediamine	$H_2N (CH_2)_6 NH_2$
		Adipic acid	HOOC (CH ₂) ₄ COOH
(ii)	Bakelite	Phenol	ОН
			О
		Formaldehyde	н с н
(iii)	Polystyrene	Styrene	_СН=СН2

- 22. (i) Maltose
 - (ii) Refer to Ans. 13 (ii) Foreign 2015.
 - (iii) Phosphodiester linkage.
- **23.** Refer to Ans. 23 Patna Region 2015.

24.
$$A = \bigcup_{\text{Benzamide}}^{+} ; B = \bigcup_{\text{Benzonitrile}}^{+} ; C = \bigcup_{\text{Benzonitrile}}^{+} ; D = \bigcup_{\text{Phenyl}}^{+} ; E = \bigcup_{\text{phenol}}^{+} ; E = \bigcup_{\text{$$

OR

Refer to Ans. 25 (OR) Delhi 2015.

25. Refer to Ans. 26 Delhi 2015.

OR

- (a) Refer to Ans. 26 (OR) (a) Delhi 2015.
- (b) Refer to Ans. 25 (OR) (b) Panchkula Region 2015.

Alternatively,


 $t_{50\%} = 23.1 \text{ min}$

For a first order reaction, $t_{75\%} = 2 \times t_{50\%}$ min

 $t_{75\%} = 2 \times 23.1 \text{ min} = 46.2 \text{ min}$

- (i) Both are sp³ hybridised. In NH₄ all the four orbitals are bonded whereas in NH₃ there is a lone pair of electrons on N, which is responsible for lone pair -bond pair repulsion in NH₃ reducing the bond angle to less tetrahedral angle 109°28'.
 - (ii) Due to inert pair effect the stability of +6 oxidation state decreases down the group from S to T_E. Therefore, reducing character decreases from SO_2 to TeO_2 .
 - (iii) Greater the number of oxygen atoms in the conjugate base greater will be the dispersal of negative charge through p -d bonding and hence greater will be the stability of the conjugate base. As the conjugate base CIO_4 is more stable than $\text{CIO}\,$, therefore HCIO_4 is a stronger acid than HCIO.





(a) Phosphine, PH₃

(b) Xenon hexafluoroplatinate, $Xe [PtF_6]^-(s)$.

As PtF_6 oxidises O_2 to O_2 , Bartlett thought that PtF_6 should also oxidise Xe to Xe^+ because the first ionisation enthalpy of O_2 (1175 kJ mol $^{-1}$) was almost identical with Xe (1170 kJ mol $^{-1}$).

- (c) It is due to
 - (i) low enthalpy of dissociation of F—F bond
 - (ii) high hydration enthalpy of F.
- (d) Uses of chlorine gas
 - (i) In sterilising drinking water
 - (ii) For bleaching wood pulp, textiles and cotton
 - (iii) In the preparation of poisonous gases such as phosgene, tear gas, mustard gas.
- (e) CaF₂ H_2SO_4 CaSO₄ 2HF





CBSE Examination Paper Allahabad Region-2015 (Code No. 56/1/RU)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination paper, Delhi-2015.

1. Write the formulae of any two oxoacids of sulphur.

2. Write the IUPAC name of the given compound:

$$CH_3 - CH - CH_2 - O - CH_2 - CH_3$$
 CH_3

- 3. A delta is formed at the meeting point of sea water and river water. Why?
- **4.** Which would undergo S_N1 reaction faster in the following pair:

$$\mathrm{CH_3} - \mathrm{CH_2} - \mathrm{CH_2} - \mathrm{Br}$$
 and $\mathrm{CH_3} - \mathrm{CH} - \mathrm{CH_3}$

- **5.** What is the formula of a compound in which the element Y forms *ccp* lattice and atoms of X occupy 2/3rd of tetrahedral voids?
- 6. Write one similarity and one difference between the chemistry of lanthanoids and that of actinoids.
- 7. (i) Write down the IUPAC name of the following complex:

$$\left[\text{Co(NH}_3)_5\text{Cl}\right]^{2+}$$

(ii) Write the formula for the following complex:

Potassium tetrachloridonickelate (II)

8. Write the reagents required in the following reactions:

(i)
$$CH_2 = CH - CH_2OH$$
 ? $CH_2 = CH - CHO$

(ii)
$$CH_3 - COOH$$
 ? $CH_3 - CONH_2$

OR

Arrange the following compounds in increasing order of their property as indicated:

- (i) CH₃COCH₃, C₆H₅COCH₃, CH₃CHO (reactivity towards nucleophilic addition reaction)
- (ii) Cl CH₂–COOH, F–CH₂ –COOH, CH₃–COOH (acidic character)
- **9.** (*i*) On mixing liquid X and liquid Y, volume of the resulting solution decreases. What type of deviation from Raoult's law is shown by the resulting solution? What change in temperature would you observe after mixing liquids X and Y?
 - (ii) What happens when we place the blood cell in water (hypotonic solution)? Give reason.
- **10.** Calculate the time to deposit 1.27 g of copper at cathode when a current of 2 A was passed through the solution of CuSO₄.

(Molar mass of $Cu = 63.5 \text{ g mol}^{-1}$, $1 \text{ F} = 96500 \text{ C mol}^{-1}$)

11. A solution is prepared by dissolving 10 g of non-volatile solute in 200 g of water. It has a vapour pressure of 31.84 mm Hg at 308 K. Calculate the molar mass of the solute.

(Vapour pressure of pure water at 308 K = 32 mm Hg)

- **12.** (*i*) Name the method of refining to obtain silicon of high purity.
 - (ii) What is the role of SiO_2 in the extraction of copper?
 - (iii) What is the role of depressants in froth floatation process?
- **13.** (*i*) Which one of the following is a polysaccharide: starch, maltose, fructose, glucose
 - (ii) Write one difference between -helix and -pleated sheet structures of protein.
 - (iii) Write the name of the disease caused by the deficiency of vitamin B_{12} .
- 14. (i) What type of isomerism is shown by the complex $[Cr(H_2O)_6]Cl_3$?
 - (ii) On the basis of crystal field theory, write the electronic configuration for d^4 ion if $_0 > P$.
 - (iii) Write the hybridisation and shape of $[CoF_6]^{3-}$. (Atomic number of Co = 27)
- **15.** How can the following conversions be carried out:
 - (i) Aniline to bromobenzene
 - (ii) Chlorobenzene to 2-chloroacetophenone
 - (iii) Chloroethane to butane

OR

What happens when

- (i) chlorobenzene is treated with Cl₂/FeCl₃,
- (ii) ethyl chloride is treated with AgNO₂,
- (iii) 2-bromopentane is treated with alcoholic KOH?

Write the chemical equations in support of your answer.

16. Examine the given defective crystal:

X^{+}	Y^{-}	X^{+}	Y ⁻	X^{+}
Y^{-}	O	Y ⁻	X^{+}	Y^{-}
X^{+}	Y^{-}	X^{+}	O	X^{+}
Y ⁻	X^{+}	\mathbf{Y}^{-}	X ⁺	Y ⁻

Answer the following questions:

- (i) Is the above defect stoichiometric or non-stoichiometric?
- (ii) Write the term used for this type of defect. Give an example of the compound which shows this type of
- (iii) How does this defect affect the density of the crystal?
- 17. Conductivity of 2.5×10^{-4} M methanoic acid is 5.25×10^{-5} S cm⁻¹. Calculate its molar conductivity and degree of dissociation.

Given: ${}^{0}(H^{+}) = 349.5 \text{ S cm}^{2} \text{ mol}^{-1} \text{ and } {}^{0}(HCOO^{-}) = 50.5 \text{ S cm}^{2} \text{ mol}^{-1}.$

- **18.** Write any three differences between physisorption and chemisorption.
- **19.** Give reasons for the following:
 - (i) Phenol is more acidic than methanol.
 - (ii) The C-O-H bond angle in alcohols is slightly less than the tetrahedral angle (109°28').
 - (iii) (CH₃)₃C-O-CH₃ on reaction with HI gives (CH₃)₃C-I and CH₃-OH as the main products and not (CH₃)₃C-OH and CH₃-I.
- **20.** Predict the products of the following reactions:

(i)
$$CH_3 - C$$
 O $H_2N - NH_2$? CH_3



- **21.** (a) Account for the following:
 - (i) Cu⁺ is unstable in an aqueous solution.
 - (ii) Transition metals form complex compounds.
 - (b) Complete the following equation:

$$Cr_2O_7^2$$
 8H⁺ 3NO₂

- 22. Write the names and structures of the monomers of the following polymers:
 - (i) Terylene

(ii) Buna-S

- (iii) Neoprene
- 23. Seeing the growing cases of diabetes and depression among young children, Mr. Chopra, the principal of one reputed school organised a seminar in which he invited parents and principals. They all resolved this issue by strictly banning junk food in schools and introducing healthy snacks and drinks like soup, *lassi*, milk, etc. in school canteens. They also decided to make compulsory half an hour of daily physical activities for the students in the morning assembly. After six months, Mr. Chopra conducted the health survey in most of the schools and discovered a tremendous improvement in the health of the students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Chopra?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) Why should antidepressant drugs not be taken without consulting a doctor?
- (iv) Give two examples of artificial sweeteners.
- **24.** (a) Account for the following:
 - (i) Acidic character increases from HF to HI.
 - (ii) There is a large difference between the melting and boiling points of oxygen and sulphur.
 - (iii) Nitrogen does not form pentahalide.
 - (b) Draw the structures of the following:
 - (i) ClF₃


(ii) XeF₄

OR

- (i) Which allotrope of phosphorus is reactive and why?
- (ii) How are the supersonic jet aeroplanes responsible for the depletion of ozone layer?
- (iii) F₂ has lower bond dissociation enthalpy than Cl₂. Why?
- (iv) Which noble gas is used in filling balloons for meteorological observations?
- (*v*) Complete the following equation:

$$XeF_2 + PF_5$$

25. An aromatic compound 'A' of molecular formula $C_7H_6O_2$ undergoes a series of reactions as shown below. Write the structures of A, B, C, D and E in the following reactions:





OR

- (a) Write the structures of main products when benzene diazonium chloride reacts with the following reagents:
 - (i) $H_3PO_2 + H_2O$
 - (ii) CuCN/KCN
 - (iii) H₂O
- (b) Arrange the following in the increasing order of their basic character in an aqueous solution:

(c) Give a simple chemical test to distinguish between the following pair of compounds:

26. For the hydrolysis of methyl acetate in aqueous solution, the following results were obtained:

t/s	0	10	20
[CH ₃ COOCH ₃]/mol L ⁻¹	0.10	0.05	0.025

- (a) Show that it follows pseudo first order reaction, as the concentration of water remains constant.
- (b) Calculate the average rate of reaction between the time interval 10 to 20 seconds.

(Given: $\log 2 = 0.3010$, $\log 4 = 0.6021$)

OR

- P, the rate is given by Rate = $k [A][B]^2$ (a) For a reaction A + B
 - (i) How is the rate of reaction affected if the concentration of B is doubled?
 - (ii) What is the overall order of reaction if A is present in large excess?
- (b) A first order reaction takes 30 minutes for 50% completion. Calculate the time required for 90% completion of this reaction.





Solutions

- 1. H₂SO₄ (sulphuric acid), H₂S₂O₈ (peroxodisulphuric acid)
- 2. 1-ethoxy-2-methyl propane
- 3. River water is a colloidal solution of clay and sea water contains a number of electrolytes. When river water meets the sea water, the electrolytes present in the sea water coagulate the colloidal solution of clay resulting in its deposition with the formation of delta.
- 4. CH₃ CH CH₃, because the secondary carbocation formed in the slowest step is more stable than the

Br

primary carbocation.

5. Let the number of particles of element Y in ccp lattice = N

Number of tetrahedral voids = 2N

Number of particles of element $X = \frac{2}{3}$ 2N $\frac{4}{3}$ N

$$X:Y = \frac{4}{3} N:N$$

$$X:Y$$
 4:3

Formula of the compound X_4Y_3

6. Similarities between lanthanoids and actinoids

- (i) Both lanthanoids and actinoids mainly show an oxidation state of +3.
- (ii) Actinoids show actinoid contraction like lanthanoid contraction is exhibited by lanthanoids.
- (iii) Both lanthanoids and actinoids are electropositive.

(any one)

Differences between lanthanoids and actinoids

- (i) The members of lanthanoid exhibit less number of oxidation states than the corresponding members of actinoid series.
- (ii) Lanthanoid contraction is smaller than the actinoid contraction.
- (iii) Lanthanoids except promethium are non-radioactive metals while actinoids are radioactive metals.

(any one)

- 7. (i) Pentaamminechloridocobalt (III) ion
 - (ii) $K_2[NiCl_4]$
- **8.** (*i*) Pyridinium chlorochromate (C₅H₅NHCrO₃Cl) or Cu at 573 K.
 - (ii) NH3, (Heat)

OR

- (i) C₆H₅COCH₃ < CH₃COCH₃ < CH₃CHO
- (ii) $CH_3COOH < Cl CH_2COOH < F CH_2COOH$
- 9. (i) The solution will show negative deviation from Raoult's law.
 - Temperature will rise.
 - (ii) Due to osmosis water enters into the cell and blood cell will swell.
- **10.** $Cu^{2+} + 2e$ Cu
 - 63.5 g of copper is deposited by 2 96500 C.

1.27 g of copper will be deposited by $\frac{2 - 96500 - 1.27}{635}$ C 3860 C



$$I 2 A, Q 3860 C$$

 $t \frac{Q}{I} \frac{3860}{2} = 1930 s$


11. W_B 10 g; W_A 200 g; P 31.84 mm Hg; P_A 32 mm Hg; M_A 18 g mol ¹ Substituting these values in the expression,

$$M_B = \frac{P_A}{P_A - P} = \frac{W_B - M_A}{W_A}$$
, we get

 $M_B = \frac{32 \text{ mm Hg}}{32 \text{ mm Hg} - 31.84 \text{ mm Hg}} = \frac{10 \text{ g} - 18 \text{ g mol}^{-1}}{200 \text{ g}}$
 $M_B = \frac{32}{016} = \frac{10 - 18}{200} \text{ g mol}^{-1} = 180 \text{ g mol}^{-1}$

- **12.** (i) Zone refining
 - (ii) SiO₂ acts as a flux to remove iron oxide impurity as slag, FeSiO₃ in the metallurgy of copper.

- (iii) In froth floatation process, the role of the depressant is to prevent one type of sulphide ore particles from forming the froth with air bubbles.
- (i) Starch is a polysaccharide.
 - (ii) In -helix structure of proteins the polypeptide chains are stabilized by intramolecular hydrogen bonding whereas -pleated sheet structure of proteins is stabilized by intermolecular hydrogen bonding.
 - (iii) Pernicious anaemia is caused due to deficiency of vitamin B₁₂.
- 14. (i) Hydration isomerism is shown by complex $[Cr(H_2O)_6]Cl_3$.
 - (ii) If $_0$ P, electronic configuration becomes $t_{2g}^4 e_g^0$.
 - (iii) $Co^3 : 3d^6 4s^0$



Hybridisation sp^3d^2 ; Shape Octahedral

15.

$$(i) \qquad \begin{array}{c} NH_2 \\ NaNO_2/HCl \\ \hline 273-278 \text{ K} \\ \end{array} \qquad \begin{array}{c} Cu_2Br_2/HBr \\ \hline \\ CI \\ \hline \\ (ii) \\ \end{array} \qquad \begin{array}{c} CI \\ CH_3 - C - Cl \\ Acetyl \ chloride \\ \end{array} \qquad \begin{array}{c} CI \\ Anhyd \ AlCl_3 \\ \hline \\ CHoroethane \\ \end{array} \qquad \begin{array}{c} CI \\ CH_3 - C - Cl \\ Acetyl \ chloride \\ \end{array} \qquad \begin{array}{c} CI \\ CH_3 - C - Cl \\ Acetyl \ chloride \\ \end{array} \qquad \begin{array}{c} CI \\ CH_3 - C - CH_3 \\ \hline \\ CH_3 - C - Cl_3 \\ \end{array} \qquad \begin{array}{c} CH_2 - CH_2 - CH_3 + 2NaCl_3 \\ \hline \\ CH_3 - CH_2 - CH_3 - CH$$



Cl
$$Cl_{2}$$
 Anhyd FeCl $_{3}$ + Cl_{2} Anhyd FeCl $_{3}$ + Cl_{2} Cl_{2} + Cl_{2} Chlorobenzene Cl_{3} + Cl_{2} Cl_{2} + Cl_{2} Cl_{3} + Cl_{2} + Cl_{2} Cl_{3} + Cl_{2} Cl_{3} + Cl_{2} Cl_{3} + Cl_{2} Cl_{3} +

Pent -1 - ene (minor product)

(ii) • Schottky defect

16. (i) Stoichiometric defect

• NaCl, CsCl

(iii) It decreases the density of the crystal.

18. Differences between physisorption and chemisorption

S. No.	Physisorption	Chemisorption	
(<i>i</i>)	It arises due to van der Waal's forces.	It arises because of chemical bond formation.	
(ii)	Enthalpy of adsorption is low (20-40 kJ mol $^{-1}$).	Enthalpy of adsorption is high (80-240 kJ mol ¹).	
(iii)	It is not specific in nature.	It is highly specific in nature.	
(iv)	It forms multimolecular layers.	It forms unimolecular layer.	
(v)	It is reversible in nature.	It is irreversible in nature.	
(vi)	No appreciable activation energy is needed.	High activation energy is sometimes needed.	



- 19. (i) In phenol, oxygen acquires positive charge as a result of resonance and releases H ion easily whereas there is no resonance in methanol. Moreover, due to I effect of CH₃ group the electron density in O—H bond increases which makes release of H difficult.
 - (ii) It is due to the repulsion between the lone pair of electrons on oxygen atoms,

$$\geq_{C}$$

- (iii) The reaction between $(CH_3)_3 COCH_3$ and HI follows $S_N 1$ mechanism. For a $S_N 1$ reaction, the formation of product is controlled by stability of the carbocation formed in the slowest step. Since tert.butyl carbonium ion ((CH₃)₃ C) formed after the cleavage of C—O bond in the slowest step is more stable than methyl carbonium ion (CH₃) therefore (CH₃)₃ C I and CH₃ OH are the main products.
- CH_3 C $NNH_2 + H_2O$ **20.** (*i*) CH₃ Propanone hydrazone Propanone (a) $KMnO_4 / KOH$ (ii) $C_6H_5CH_3$ C_6H_5COOH (b) H+ Benzoic acid .COOH COOH (iii) + HBr Benzoic acid

3-Bromo benzoic acid

21. (a) (i) In aqueous solution Cu undergoes disproportionation to form more stable Cu²⁺ ion.

$$2Cu^+$$
 (aq) Cu^{2+} $(aq) + Cu$ (s)

The higher stability of Cu^{2+} ion in aqueous solution is due to its greater negative hvd than Cu. It compensates the second ionisation enthalpy of Cu involved in the formation of Cu²⁺ ions.

(ii) Transition metals form complex compounds due to presence of vacant d-orbitals of suitable energy for bond formation, comparatively small size of metal ions, high ionic charge.

22.

	Polymer	Name of the monomers	Structure of the monomers
(<i>i</i>)	Terylene	Ethylene glycol	HOCH ₂ CH ₂ OH
		Terephthalic acid	ноос-Соон
(ii)	Buna-S	1, 3-Butadiene	CH ₂ CH CH CH ₂
		Styrene	CH=CH ₂
(iii)	Neoprene	Chloroprene	CH ₂ C CH CH ₂
			Cl



23. (*i*) The values displayed by Mr. Chopra are :

- Caring in nature, empathy
- Social awareness
- Concern for student's health
- (ii) Awareness for health can be spread by
 - performing *nukkad natak* in community.
 - performing play in the school.
 - displaying posters/cartoon/slogans.
 - conducting seminars.
- (iii) Because wrong choice and overdose may be harmful.
- (iv) Saccharin, sucrolose, aspartame are the examples of artificial sweeteners.

24. Refer to Ans. 24 OR Delhi 2015.

25.

$$A = \bigcup_{\substack{C \\ C \\ Benzoic \\ acid}} ; B = \bigcup_{\substack{NH_2 \\ Aniline \\ ethanamide}} ; C = \bigcup_{\substack{NH_2 \\ N-Phenyl \\ ethanamide}} ; C = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ Br \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ Br \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 2,4,6-Tribromo \\ aniline}} ; E = \bigcup_{\substack{Renzylamine \\ SP \\ 3,4,6-Tribromo \\ 3,4,6-Tri$$

OR

(a) (i)

Benzene
diazonium chloride

H₃PO₂ + H₂O

Benzene

+ H₃PO₃ + N₂ + HCl

Benzene

CN

CuCN/KCN

CuCN/KCN

Benzonitrile

OH

H₂O

$$A$$

Phenol

- (b) $C_2H_5NH_2 < (C_2H_5)_3N < (C_2H_5)_2NH$
- (c) Aniline being a primary amine gives carbylamine test, *i.e.*, when warmed with an alcoholic solution of KOH and CHCl₃ it gives foul smell of phenyl carbylamine. In contrast, N-methyl aniline (C₆H₅NHCH₃) being a secondary amine does not give this test.



$$\begin{array}{c} C_6H_5NH_2 + CHCl_3 + 3KOH \, (alc.) \\ \text{Aniline} \\ \end{array} \begin{array}{c} C_6H_5NC \\ \text{Phenyl} \\ \text{isocyanide} \\ \text{(Phenyl carbylamine)} \\ \end{array} \\$$

0.1 mol L⁻¹, [R] 0.05 mol L⁻¹, t 10 s in the expression **26.** (a) Substituting $[R]_0$ $k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$, we get

$$k = \frac{2.303}{10} \log \frac{0.1}{0.05} = \frac{2.303}{10} \log 2 = \frac{2.303}{10} = 0.3010$$

Again substituting $[R]_0$ 0.10 mol L ¹, [R] 0.025 mol L ¹, t 20 s in the expression $k = \frac{2303}{t} \log \frac{[R]_0}{[R]} \text{, we get}$

$$k = \frac{2.303}{20} \log \frac{0.1}{0.025} = \frac{2.303}{20} \log 4 = \frac{2.303}{20} = 0.6021$$

As the value of rate constant is same in both the readings therefore hydrolysis of methyl acetate follows pseudo first order reaction.

(b) Average rate $\frac{-\;[0.025\;\text{mol}\;L^{-1}-0.05\;\text{mol}\;L^{-1}]}{20\;\text{s}-10\;\text{s}} \quad \frac{0.025\;\text{mol}\;L^{-1}}{10\;\text{s}}$

 $0.0025 \text{ mol L}^{-1} \text{ s}^{-1}$ Average rate

OR

Refer to Ans. 26 (OR) Delhi 2015.



CBSE Examination Paper

Bhubaneshwar Region -2015 (Code No. 56/1/B)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper, Delhi-2015.

What is the type of charge on AgI colloidal sol formed when AgNO₃ solution is added to KI solution?

- What is the formula of a compound in which the element Y forms hcp lattice and atoms of X occupy 1/3rd of octahedral voids?
- Write the formulae of any two oxoacids of chlorine.
- Write the IUPAC name of the given compound:

Which would undergo S_N2 reaction faster in the following pair:

6. Name the reagents used in the following reactions:

(i)
$$C_6H_5$$
 — COCI $^?$ $^?$ $^?$ $^?$ C $_6H_5$ — CHO (ii) $^?$ CH $_4$

OR

Arrange the following compounds in increasing order of their property as indicated:

- (i) CH₃COCH₃, C₆H₅—CO—C₆H₅,CH₃CHO (reactivity towards nucleophilic addition reaction)
- (ii) Cl CH COOH, $Cl CH_2 COOH$, $CCl_3 COOH$ (acidic character)
- (i) Why do actinoids show wide range of oxidation states?
 - (ii) Why is actinoid contraction greater than lanthanoid contraction?
- Calculate the time of deposit 1.17 g of Ni at cathode when a current of 5A was passed through the solution of $Ni(NO_3)_2$.

(Molar mass of Ni = 58.5 g mol^{-1} , $1\text{F} = 96500 \text{ C mol}^{-1}$)

- (i) Write down the IUPAC name of the following complex: $K_3[Fe(CN)_6]$
 - (ii) Write the formula for the following complex:

Pentaamminenitrito-N-cobalt(III)

- 10. (i) On mixing liquid X and liquid Y, the volume of the resulting solution increases. What type of deviation from Raoult's law is shown by the resulting solution? What change in temperature would you observe after mixing liquids X and Y?
 - (ii) How can the direction of osmosis be reversed? Write one use of reverse osmosis.
- **11.** (*i*) Which one of the following is an oligosaccharide:

starch, maltose, fructose, glucose

- (iii) Write the name of the disease caused by the deficiency of Vitamin B₁.
- **12.** Calculate E_{cell} for the following reaction at 25°C:

A + B²⁺ (0.001 M)
$$A^{2+}$$
 (0.0001 M) + B
Given $E_{cell} = 2.6805$, 1 F = 96500 C mol⁻¹

13. Differentiate between the following:

- (i) Solution and Colloid
 - (ii) Homogeneous catalysis and Heterogeneous catalysis
 - (iii) O/W emulsion and W/O emulsion
- 14. A solution is prepared by dissolving 5 g of non-volatile solute in 95 g of water. It has a vapour pressure of 23.375 mm Hg at 25°C. Calculate the molar mass of the solute. (Vapour pressure of pure water at 25°C is 23.75 mm Hg)
- **15.** (i) Name the method of refining to obtain low boiling metal like mercury.
 - (ii) What is the role of pine oil in froth floatation process?
 - (iii) Reduction of metal oxide to metal becomes easier if the metal obtained is in liquid state. Why?
- **16.** Examine the given defective crystal:

$$X^{+}$$
 Y^{-} X^{+} Y^{-}

Answer the following questions:

- (i) Is the above defect stoichiometric or non-stoichiometric?
- (ii) Write the term used for this type of defect.
- (iii) Why do silver halides show this type of defect?
- 17. Predict the products of the following reactions:

(i)
$$CH_3 - C$$
 O $\stackrel{HCN}{\downarrow}$

(i)
$$CH_3 - C$$
 O HCN ?

H

(ii) $C_6H_5 - MgBr$ (a) CO_2 ?

(b) H_3O

(iii)
$$CH_3 - CN$$
 LiAl H_4 ?

- **18.** Write the names and structures of the monomers of the following polymers:
 - (i) Glyptal

- (ii) Teflon
- (iii) Nylon-6
- **19.** (*i*) MnO is basic whereas Mn_2O_7 is acidic in nature. Why?
 - (ii) Transition metals form alloys. Why?
 - (iii) Complete the following equation:

$$2MnO_2$$
 4KOH O_2

- **20.** (i) What type of isomerism is shown by the complex $[Ag(NH_3)_2][Ag(CN)_2]$?
 - (ii) On the basis of crystal field theory, write the electronic configuration for d^4 ion if d^4
 - (iii) Write the hybridisation and shape of $[Ni(CN)_4]^{2-}$. (Atomic number of Ni = 28)
- **21.** Give reasons for the following:
 - (i) Benzyl chloride is highly reactive towards the S_N1 reaction.
 - (ii) 2-bromobutane is optically active but 1-bromobutane is optically inactive.

- (iii) Electrophilic reactions in haloarenes occur slowly.
- 22. How do you convert the following:
 - (i) Aniline to phenol
 - (ii) Prop-1-ene to Propan-1-ol
 - (iii) Anisole to 2-methoxytoluene

OR

What happens when

- (i) ethanol is treated with Cu at 573 K,
- (ii) phenol is treated with CH₃ COCl/anhydrous AlCl₃,
- (iii) ethyl chloride is treated with NaOCH₃?

Write chemical equations in support of your answer.

23. Mr. Roy, the principal of one reputed school organized a seminar in which he invited parents and principals to discuss the serious issue of diabetes and depression in students. They all resolved this issue by strictly banning junk food in schools and introducing healthy snacks and drinks like soup, lassi, milk, etc. in school canteens. They also decided to make compulsory half an hour of daily physical activities for the students in the morning assembly. After six months, Mr. Roy conducted the health survey in most of the schools and discovered a tremendous improvement in the health of the students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Roy?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) What are tranquilizers? Give an example.
- (iv) Why is use of aspartame limited to cold foods and drinks?
- **24.** For the hydrolysis of methyl acetate in aqueous solution, the following results were obtained:

t/s	0	20	40
[CH ₃ COOCH ₃]/mol L ⁻¹	0.40	0.20	0.10

- (a) Show that it follows pseudo first order reaction, as the concentration of water remains constant.
- (b) Calculate the average rate of reaction between the time interval 20 to 40 seconds.

OR

- (a) Define the following terms:
 - (i) Collision frequency
 - (ii) Rate constant (k)
- (b) The rate constant of a first order reaction increases from 4×10^{-2} to 24×10^{-2} when the temperature changes from 300 K to 350 K. Calculate the energy of activation (E_a) .

$$(\log 2 = 0.301, \log 3 = 0.4771, \log 4 = 0.6021, \log 6 = 0.7782)$$

- **25.** (a) Account for the following:
 - (i) Bi(V) is a stronger oxidizing agent that Sb(V).
 - (ii) H—O—I is a weaker acid than H—O—Cl.
 - (iii) Bond angle decreases from H₂O to H₂S.
 - (b) Draw the structures of the following:
 - (*i*) SF₄
 - (ii) XeF₂


OR

- (i) Why does PCl₅ fume in moisture?
- (ii) Write the name of the allotrope of sulphur which is stable at room temperature.
- (iii) Chlorine water on standing loses its yellow colour. Why?
- (iv) Write the disproportionation reaction of H₃PO₃.

(v) Complete the following equation:

$$F_2$$
 H_2O

26. Write the structures of A, B, C, D and E in the following reactions :



OR

- (i) Write the structures of the main products when benzene diazonium chloride reacts with the following reagents:
 - (a) KI
 - (b) CH₃CH₂OH
 - (c) Cu / HCI
- (ii) Arrange the following in the increasing order of their basic character in an aqueous solutions:

$$CH_3NH_2$$
, $(CH_3)_2NH$, $(CH_3)_3N$

(iii) Give a simple chemical test to distinguish between the following pair compounds:



Solutions -

- 1. Negatively charged sol, AgI/I is formed when AgNO₃ solution is added to KI solution.
- **2.** Let the number of atoms of element Y in hcp lattice N

Number of octahedral voids 1 N N

Number of atoms of element $X = \frac{1}{3}$ N

$$X: Y = \frac{1}{3} N: N = 1:3$$

Formula of the compound XY_3

- 3. HOCl, HOClO, HOClO₂, HOClO₃.
- 4. 1-phenylpropan-2-ol
- 5. CH₃ CH CH₂ CH₂ Br, the presence of methyl group away from the halide group will decrease

the steric hinderance and hence the rate.

- **6.** (i) H_2/Pd -BaSO₄
 - (ii) NaOH/CaO, Heat

OR

- $(i) C_6H_5COC_6H_5 < CH_3COCH_3 < CH_3CHO$
- (ii) Cl CH₂ COOH < Cl CH COOH < CCl₃ COOH

C

- 7. (i) Because 5f, 6d and 7s orbitals are of comparable energies.
 - (ii) As 5f electrons provide poorer shielding effect than 4f electrons, therefore, actinoid contraction is greater than lanthanoid contraction.
- **8.** At cathode: $Ni^{2+} + 2e$

58.5 g of Ni deposited by 2F 2 96500 C

1.17 g of Ni will be deposited by
$$\frac{2 - 96500 - 1.17 \text{ C}}{58.5}$$
 3860 C

Substituting Q 3860 C, I 5 A in the expression, t $\frac{Q}{I}$, we get

$$t = \frac{3860 \text{ C}}{5 \text{ A}}$$
 772 s or $t = \frac{772}{60}$ 12.87 min

- **9.** (*i*) Potassium hexacyanidoferrate (III)
 - (ii) $[Co(NH_3)_5 NO_2]^{2+}$
- **10.** (*i*) The resulting solution will show positive deviation from Raoult's law and it would have lower temperature.
 - (ii) The direction of osmosis can be reversed by applying an external pressure greater than the osmotic pressure on the solution side. Reverse osmosis is used in desalination of sea water.
- **11.** (*i*) Maltose
 - (ii) DNA contains cytosine and thymine as pyrimidine bases whereas RNA contains cytosine and uracil as pyrimidine bases.

DNA controls the transmission of hereditary effects whereas RNA controls the synthesis of protein.

(iii) Beri-Beri

12. At anode:
$$A A^2$$

Substituting E_{cell} 2.6805 V, n 2, $[A^2]$ 1 10 4 M, $[B^2]$ 1 10 3 M in the Nernst

equation for the above cell reaction,
$$E_{\text{cell}} = E_{\text{cell}}^{\text{o}} = \frac{0.059}{n} \log \frac{[A^2]}{[B^2]}$$
, we get

2.6805
$$E_{\text{cell}}^{\text{o}} = \frac{0.059}{2} \log \frac{10^{-4}}{10^{-3}}$$

$$E_{\text{cell}}^{\text{o}}$$
 2.6805 0.0295 log 10 ¹ = 2.6805 - 0.0295

$$E_{\text{cell}}^{\text{o}}$$
 2.651 V

13. (i) Differences between Solution and Colloid

	Property	Solution	Colloid
•	Nature	Homogeneous	Heterogeneous
•	Particle size	Less than 1 nm	Between 1 nm to 1000 nm
•	Appearance	Transparent	Translucent
•	Filterability	Pass through ordinary filter paper as well as animal membrane.	Pass through ordinary filter paper but not through animal membrane.

(ii) Differences between Homogeneous catalysis and Heterogeneous catalysis

	Homogeneous catalysis	Heterogeneous catalysis
•	In this process reactants and catalyst are in the same phase.	In this process reactants and catalyst are in different phases.
	Example: $2SO_2(g) + O_2(g)$ $SO_3(g)$ $2SO_3(g)$	Example: $N_2(g) + 3H_2(g)$ Fe(s) $N_3(g)$ $NH_3(g)$

(iii) Differences between O/W Emulsion and W/O Emulsion

	O/W Emulsion	W/O Emulsion
•	Oil is the dispersed phase and water is the dispersion medium.	Water is the dispersed phase and oil is the dispersion medium.
•	Emulsifying agents used are gums, proteins, natural and synthetic detergents, etc.	Emulsifying agents used are lamp black, heavy metal salts of fatty acids, long chain alcohols, etc.
•	Examples: milk, vanishing cream, etc.	Examples: butter, cold cream, etc.

14. Substituting P_A 23.75 mm Hg, P 23.375 mm Hg, W_B 5 g, W_A 95 g, W_A 18 g mol 1 in the

expression
$$M_B = \frac{P_A}{P_A - P} = \frac{W_B - M_A}{W_A}$$
 , we get

$$M_B = \frac{23.75 \text{ mm Hg}}{23.75 \text{ mm Hg}} = \frac{5 \text{ g}}{23.75 \text{ mm Hg}} = \frac{5 \text{ g}}{95 \text{ g}} = \frac{18 \text{ g mol}}{95 \text{ g}}$$
 $M_B = \frac{23.75}{0.375} = \frac{5}{95} = \frac{18}{95} =$

- **15.** (*i*) Distillation
 - (ii) Pine oil (collector) enhance the non-wettability of the mineral particles.
 - (iii) Refer to Ans. 15 (iii) Panchkula Region-2015

- **16.** (*i*) Stoichiometric defect.
 - (ii) Frenkel defect.
 - (iii) Because Ag⁺ ion due to its small size fits into the interstitial site.

17. (*i*) CH₃ CH_3 OH Η Cyanohydrin $(ii) \ \mathrm{C_6H_5MgBr} \atop (b) \ \mathrm{H_3O^+} \ \mathrm{C_6H_5} \quad \begin{array}{c} \mathrm{COOH} \\ \mathrm{Benzoic\ acid} \end{array}$ CH_2

18.

	Name of Polymer	Name of Monomers	Structure of Monomers
(<i>i</i>)	Glyptal	Ethylene glycol	HO CH ₂ CH ₂ OH
		Phthalic acid	O O HO—C C—OH
(ii)	Teflon	Tetrafluoroethene	F C C F
(iii)	Nylon-6	Caprolactum	$H_{2}C$ N C $H_{2}C$ CH_{2} $H_{2}C$ CH_{2}

- (i) The oxide in the lower oxidation state of a metal is basic and in the higher oxidation state of the metal it is acidic. That is, why MnO is basic whereas $\mathrm{Mn}_2\mathrm{O}_7$ is acidic.
 - (ii) The transition metals have similar atomic radii and other characteristics due to this the atoms of one metal can easily take up the positions in the crystal lattice of the other and hence they form a large number of alloys, e.g., stainless steel, brass, etc.

(iii) 2MnO₂ + 4KOH + O₂ 2K₂MnO₄ + 2H₂O Potassium manganate

- **20.** (ii) If $_0$ P, the fourth electron enters one of the e_g orbitals giving the configuration $t_{2g}^3 e_g^1$.
 - (*iii*) $Ni^{2+}: 3d^8 4s^0$

Hybridisation dsp^2 , Shape Square planar.

(i) Benzyl chloride is highly reactive towards the S_N 1 reaction because the intermediate benzyl carbocation 21. formed in slowest step is stabilized through resonance.

$$\stackrel{\stackrel{+}{\text{CH}_2}}{\longleftrightarrow} \stackrel{\text{CH}_2}{\longleftrightarrow} \stackrel{\text{CH}_2}{\longleftrightarrow} \stackrel{\text{CH}_2}{\longleftrightarrow}$$

Benzyl cation stabilised by resonance

(ii) 2-bromobutane is a chiral molecule as it contains an asymmetric carbon atom therefore, it is optically active whereas 1-bromobutane is an achiral molecule as it does not contain asymmetric carbon atom therefore it is optically inactive.

$$\begin{array}{cccc} & & Br \\ & CH_3 & CH_2 & C & CH_3 \\ & & H \\ & carbocation of 2-Bromobutane \end{array}$$

(iii) Halogen in haloarenes withdraws electrons through I effect and release electrons through R effect. The inductive effect is stronger than resonance effect and causes net electron withdrawal. As a result, the electrophilic substitution reactions in haloarenes occur slowly.

22. (i) NaNO₂/HCl
$$\frac{1}{273-278 \text{ K}}$$
 NaNO₂/HCl $\frac{1}{273-278 \text{ K}}$ NaNO₂/HCl $\frac{1}{273-278 \text{ K}}$ NaNO₂/HCl $\frac{1}{273-278 \text{ K}}$ NaNO₂/HCl $\frac{1}{273-278 \text{ K}}$ NaNO₂/HCl $\frac{1}{283 \text{ K}}$ Phenol $\frac{1}{283 \text{ K}}$ Phenol $\frac{1}{283 \text{ K}}$ Phenol $\frac{1}{283 \text{ K}}$ CH₂ CH₂ CH₂ OH $\frac{1}{283 \text{ K}}$ CH₃ CH₂ CH₂ OH $\frac{1}{283 \text{ K}}$ CH₃ CH₂ CH₃ CH₃ CH₃ CH₃ CH₄ CH₃ CH₄ CH₃ CH₄ CH₅ CH₅

CH₃CH₂

O

Ethyl methyl ether

(iii) CH₃

 CH_2

Ethyl chloride

Cl + NaOCH₃

Sodium

methoxide

4-Hydroxy acetophenone

CH₃ + NaCl

- 23. Refer to Ans. 23 Delhi Region 2015
- **24.** (a) For a first-order reaction

$$k = \frac{2303}{t} \log \frac{[R]_0}{[R]} = \dots(i)$$

Substituting
$$[R]_0$$
 0.40 mol L⁻¹, $[R]$ 0.20 mol L⁻¹, t 20 s in equation (i), we get

$$k = \frac{2303}{20} \log \frac{0.40}{0.20} = \frac{2303}{20} \log 2 = \frac{2303}{20} = 0.3010$$

Again substituting
$$[R]_0$$
 0.40 mol L¹, $[R]$ 0.10 mol L¹, t 40 s in equation (i), we get

$$k = \frac{2303}{40} \log \frac{0.40}{0.10} = \frac{2.303}{40} \log 4 = \frac{2303}{20} = 0.602$$

As constant values of k are obtained by applying first order integrated rate law, therefore, hydrolysis of methyl acetate in aqueous solution is a pseudo first order reaction.

(b) Average rate

$$\frac{[\text{CH}_3 \quad \text{COOCH}_3]}{t}$$

$$\frac{0.10 \text{ mol } \text{L}^{-1} \quad 0.20 \text{ mol } \text{L}^{-1}}{40 \text{ s} \quad 20 \text{ s}}$$

$$\frac{0.1 \text{ mol } \text{L}^{-1}}{20 \text{ s}} \quad 0.005 \text{ mol } \text{L}^{-1} \text{ s}^{-1}$$

- (a) (i) The number of collisions per second per unit volume of the reaction mixture is known as collision frequency.
 - (ii) Refer to Ans. 7 Patna Region 2015

(b)
$$\log \frac{k_2}{k_1} - \frac{E_a}{2303 - R} - \frac{T_2 - T_1}{T_1 T_2}$$

Substituting k_1 4 10 2 , k_2 24 10 2 , T_1 300 K, T_2 350 K, R 8.314 J K 1 mol 1 in the above equation, we get

$$\log \frac{24 + 10^{-2}}{4 + 10^{-2}} = \frac{E_a}{2303 + 8.314} = \frac{350 + 300}{300 + 350}$$

$$\log 6 = \frac{E_a}{19.147} = \frac{50}{300 + 350}$$

$$0.7782 = \frac{E_a}{19.147 + 300 + 7}$$

$$E_a = 0.7782 + 19.147 + 300 + 7$$

$$E_a = 0.7782 + 19.147 + 300 + 7$$

$$E_a$$
 31290.41 J mol ¹ 31.29 kJ mol ¹


- **25.** (*a*) (*i*) Because 3 oxidation state of Bi is more stable than Sb due to inert pair effect.
 - (ii) Because chlorine is more electronegative than iodine.
 - (iii) Oxygen is more electronegative than sulphur due to this bond pair electrons of the O—H bond will be closer to oxygen than the bond pair electrons of S-H bond which will be little away from the sulphur atom. Consequently bond pair-bond pair repulsions would be stronger in O—H bond than in the S—H bond and hence bond angle in H₂O is greater than H₂S.


$$(iv) 4H_3 PO_4 3H_3 PO_4 + PH_3$$

$$(v) 2F_2 + 2H_2O$$

$$4HF + O_2$$

(*b*)





(i) Due to formation of fumes of HCl

POCl₃ 2HCl

- (ii) Rhombic sulphur or sulphur
- (iii) The yellow colour of chlorine water is due to dissolved Cl2. On standing, the Cl2 is assumed in reacting with water to form colourless products.

$$Cl_{2} + H_{2}O \qquad HOCl \quad HCl$$

$$2HOCl \qquad 2HCl + O_{2}$$


$$O \qquad \qquad 0$$

$$\parallel \qquad \qquad \qquad NH-C-CH_{3}$$

$$Acetanilide \qquad Br$$

$$p-bromo$$

$$acetanilide$$



OR

(a) (i)
$$+ KI \longrightarrow I$$
 $+ KCI + N_2$
Benzene diazonium Iodobenzene

chloride

(iii)
$$Cu/HCl$$
 Cl Cl Cu/HCl Cu/HCl $Chlorobenzene$

- $(b) \; ({\rm CH_3})_3 \, {\rm N} < {\rm CH_3NH_2} < ({\rm CH_3})_2 \, {\rm NH}$
- $-\mathrm{NH}_2$ on treatment with NaNO_2 and dil. HCl at 273-278 K followed by treatment with

an alkaline solution of 2-napthol gives an orange coloured azo dye.

$$NH_{2} \xrightarrow{+ \text{NaNO}_{2}/\text{HCl}} NH_{2} \xrightarrow{+ \text{NaNO}_{2}/\text{HCl}} VH_{2} \xrightarrow{+ \text{NaNO}_{2}/\text{HCl}} VH_{2} VH_{2$$

1-phenylazo-2-naphthol (orange dye)

 NH_2 under these conditions gives brisk evolution of N_2 gas On the other hand, CH₃

$$\begin{array}{c} & \text{HNO}_2 \text{ (NaNO}_2/\text{HCl)} \\ \text{CH}_3 \text{ NH}_2 & \text{CH}_3 \text{ OH} + \text{N}_2 & \text{H}_2 \text{O} \\ \text{Methyl} & \text{273-278 K} & \text{Methyl} \\ \text{amine} & \text{Methyl} & \text{alcohol} \end{array}$$

CBSE Examination Paper Chennai Region-2015 (Code No. 56/3/MT)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination paper, Delhi-2015.

1. How much charge in Faradays is required for the reduction of 1 mol of Al³⁺ to Al?

2. Which would undergo S_N2 reaction faster in the following pair:

$$CH_3$$
— CH_2 — Br and CH_3 — CH_3
 CH_3
 CH_3
 CH_3
 CH_3
 CH_3
 CH_3

3. Write the IUPAC name of compound.

- **4.** Write the dispersed phase and dispersion medium of paints.
- **5.** Copper atom has completely filled *d*-orbitals in its ground state but it is a transition element. Why?
- **6.** Arrange the following:
 - (i) in increasing order of basic strength:

(ii) in increasing order of boiling point:

- 7. Define rate constant (k). Write the unit of rate constant for the following:
 - (i) First order reaction
- (ii) Second order reaction
- **8.** Write the structures of the following:
 - (i) H₂S₂O₇

- (ii) XeO₃
- **9.** Derive the relationship between relative lowering of vapour pressure and molar mass of the solute.
- 10. Write down the IUPAC name of the complex $[Co(NH_3)_5(CO)_3]$ Cl. What type of isomerism is shown by this complex?

OR

Using IUPAC norms write the formulae for the following coordination compounds:

- (i) Tetrachloridocuprate (II)
- (ii) Potassium tetrahydroxozincate (II)
- 11. Give reasons for the following:
 - (a) p-nitrophenol is more acidic than o-nitrophenol.
 - (b) Bond angle C-O-C in ethers is slightly higher than the tetrahedral angle (109°28′).
 - (c) (CH₃)₃C–Br on reaction with NaOCH₃ gives an alkene instead of an ether.
- **12.** How do you convert the following:
 - (i) Aniline to benzene
- (ii) Ethanamide to methanamine
- (iii) Nitrobenzene to aniline

OR

Write the chemical equations involved when C₂H₅NH₂ is treated with the following reagents:

- (i) CH₃COCl/pyridine
- (ii) C₆H₅SO₂Cl

- (iii) CHCl₃ + KOH
- **13.** Define the following terms:
 - (i) F-centre

(ii) p-type semiconductor

- (iii) Ferrimagnetism
- 14. The rate constant of a first order reaction increases from 2×10^{-2} to 8×10^{-2} when the temperature changes from 300 K to 320 K. Calculate the energy of activation (E_a) .

$$(\log 2 = 0.301, \log 3 = 0.4771, \log 4 = 0.6021)$$

- **15.** Define the following terms:
 - (i) Homogeneous catalysis
- (ii) Coagulation
- (iii) Macromolecular colloids
- **16.** Write the structure of the major product in each of the following reactions:
 - (i) $CH_3-CH = CH_2 + H_2O$
 - ethanol/heat

- 17. (i) Mention the principle behind the zone refining of metals.
 - (ii) What is the role of dilute NaCN in the extraction of gold?
 - (iii) Which form of iron is the purest form of commercial iron?
- 18. When 1.5 g of a non-volatile solute was dissolved in 90 g of benzene, the boiling point of benzene raised from 353.23 K to 353.93 K. Calculate the molar mass of the solute.

 $(K_h \text{ for benzene} = 2.52 \text{ K kg mol}^{-1})$

- 19. (i) Write the product obtained when D-glucose reacts with Br_2 water.
 - (ii) What type of linkage is present in proteins?
 - (iii) Write one difference between DNA and RNA.
- **20.** (a) Write the hybridisation and shape of the following complexes?
 - (i) $[Co(NH_3)_6]^{3+}$
- $(ii) [NiCl_4]^{2-}$

(Atomic number: Co = 27, Ni = 28)

- (b) Out of NH₃ and 'en', which ligand forms more stable complex with metal and why?
- **21.** Write the names and structures of the monomers of the following polymers:
 - (i) Buna-N

(ii) Bakelite


- (iii) Teflon
- **22.** Give reasons for the following:
 - (i) Dinitrogen is a gas but phosphorus is a solid.
 - (ii) Bond angle decreases from H₂O to H₂Te.
 - (iii) Halogens have the maximum negative electron gain enthalpy.
- 23. Seeing the growing cases of diabetes and depression among young children, Mr. Chopra, the principal of one reputed school organised a seminar in which he invited parents and principals. They all resolved this issue by strictly banning junk food in schools and introducing healthy snacks and drinks like soup, *lassi*, milk, etc. in school canteens. They also decided to make compulsory half an hour of daily physical activities for the



students in the morning assembly. After six months, Mr. Chopra conducted the health survey in most of the schools and discovered a tremendous improvement in the health of the students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Chopra?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) Why should antidepressant drugs not be taken without consulting a doctor?
- (iv) Give two examples of artificial sweeteners.
- **24.** (a) Write the structures of A, B, C and D in the following reactions:



- (b) Distinguish between the following:
 - (i) $C_6H_5 COCH_3$ and $C_6H_5 COCH_2CH_3$
- (ii) Benzoic acid and Phenol
- (c) Write the structure of 2-hydroxybenzaldehyde.

- (a) Write the structures of the main products when ethanal (CH₃-CHO) reacts with the following reagents:
 - (i) HCN

$$(ii)$$
 H₂N—NH₂/H⁺

- (iii) LiAlH₄
- (b) Arrange the following in the increasing order of their reactivity towards nucleophilic addition reaction:

(c) Give a simple chemical test to distinguish between the following pair of compounds:

25. Calculate emf and G for the following cell:

$$\operatorname{Ni}(s)|\operatorname{Ni}^{2+}(0.01\ \mathrm{M}\parallel \mathrm{Ag}^+(0.001\ \mathrm{M})|\ \mathrm{Ag}(s)$$

Given:
$$E_{\text{(Ni}^2 / \text{Ni)}}^{\text{o}} = -0.25\text{V}, E_{\text{(Ag /Ag)}}^{\text{o}} = +0.80\text{V}$$

- (a) The conductivity of 0.1 mol L⁻¹ solution of NaCl is 1.06×10^{-2} S cm⁻¹. Calculate its molar conductivity and degree of dissociation (). Given ${}^{0}(Na^{+}) = 50.1 \text{ S cm}^{2} \text{ mol}^{-1}$ and ${}^{0}(C1^{-}) = 76.5 \text{ S cm}^{2} \text{ mol}^{-1}$.
- (b) What is the difference between primary battery and secondary battery? Give one example of each type.
- **26.** (a) Account for the following:
 - (i) Ce⁴⁺ is a strong oxidising agent in aqueous solution.
 - (ii) Transition metals have high enthalpy of atomisation.
 - (iii) Mn shows maximum number of oxidation states in 3d series.
 - (b) Complete the following equations:
 - (i) $2\text{MnO}_4^ 6\text{H}^+$ 5NO_2^- (ii) Cr_2O_7^2 14H 6Fe^2

- (a) Account for the following:
 - (i) Transition metals form coloured compounds.
 - (ii) Cr²⁺ is a strong reducing agent.
 - (iii) Actinoids show irregularities in their electronic configurations.
- (b) Define lanthanoid contraction. Write the common oxidation state of lanthanoids.





Solutions

1. Al³ 3eAl

Quantity of charge required for reduction of 1 mol of Al³

96500 C 289500 C

- 2. Refer to Ans. 3 Delhi Region 2015.
- 3. 1-methoxypropan-2-ol
- **4.** Dispersed phase : Solid; Dispersion medium : Liquid
- 5. Copper exhibits +2 oxidation state wherein it has incompletely filled d orbitals $(3d^9 4s^0)$ hence a transition element.
- (i) $C_6H_5NH_2$ $C_6H_5NH-CH_3$ $CH_3-CH_2-NH_2$
 - (ii) $CH_3 NH CH_3 CH_3 CH_2 NH_2 C_2H_5 OH$
- 7. The rate of reaction when molar concentration of each of the reactants is unity is called rate constant. Its value depends on the temperature but independent of concentration of reactants.

Order of reaction


Unit of k

(i) First


(ii) Second

L mol
1
 time 1

8. (i)



Pyrosulphuric acid (Oleum) $(H_2S_2O_7)$



9. From Raoult's we have, *P* P_R

If solute B is non-volatile, then

For a dilute solution n_B n_A , so n_B can be neglected in denominator in comparison to n_A .

10. Pentaammine carbonato cobalt (III) chloride

Ionisation isomerism and the ionisation isomer is [Co(NH₃)₅ Cl] CO₃.

- (i) $[CuCl_4]^2$
- (ii) $K_2[Zn(OH)_4]$



H-bonding in o-nitrophenol

Intramolecular H-bonding in o-nitrophenol makes loss of proton difficult. Therefore, p-nitrophenol is more acidic than o-nitrophenol

- bond angle in ether is slightly higher than 109°28' due to repulsive interaction between the two bulky alkyl groups.
- (c) It is because NaOCH3 is a strong nucleophile as well as a strong base. Thus, elimination reaction predominates over substitution reaction.

Ethyl amine Acetyl chloride N - Ethyl acetamide

(ii) C₂H₅NH₂ $C_6H_5SO_2CI$ C_2H_5NH O_2S C_6H_5 Ethyl amine Benzene N - Ethyl benzene sulphonyl sulphonamide chloride

(iii) C2H5NH2 CHCl₃ 3KOH(alc.) C2H5NC 3KCl 3H₂O Ethyl amine Ethyl isocyanide

- (i) The anionic sites occupied by unpaired electrons are called F-centres. They impart colour to alkali halide like NaCl, KCl, LiCl.
 - (ii) When Si or Ge is doped with a trivalent impurity like B or Al then electron vacancies are created called positive holes which impart conduction. Such type of semiconductors are called *p*-type semiconductors.
 - (iii) Ferrimagnetism is observed when the magnetic moments of the domains in the substance are aligned in parallel and antiparallel directions in unequal numbers leading to a small net magnetic moment e.g., magnetite (Fe $_3$ O $_4$).

Alignment of magnetic moment of domains in a ferrimagnetic substance



14. Substituting
$$k_1$$
 2 10 ², k_2 8 10 ², T_1 300 K, T_2 320 K, R 8.314 J K ¹mol ¹ in the expression $\log \frac{k_2}{k_1} = \frac{E_a}{2303 - R} = \frac{T_2 - T_1}{T_1 T_2}$, we get

$$\log \frac{8 \quad 10^{2}}{2 \quad 10^{2}} \quad \frac{E_{a}}{2303 \quad 8314} \quad \frac{320 \quad 300}{320 \quad 300}$$
$$\log 4 \quad \frac{E_{a}}{19147} \quad \frac{20}{320 \quad 300}$$

$$E_a$$
 0.6021 19.147 16 300 5533636 J mol ¹ E_a 55.34 kJ mol ¹

(i) In a catalysis process if the catalyst and the reactants are in the same phase (liquid or gas), the process is said to be homogeneous catalysis. For example, oxidation of SO₂ to SO₃ with O₂ in the presence of NO as a catalyst.

$$2SO_2(g)$$
 $O_2(g)$ $NO(g)$ $2SO_3(g)$

- (ii) The process of settling of colloidal particles forming a precipitate is called coagulation.
- (iii) Macromolecules in a suitable solvent form solutions in which the size of the macromolecules may be in colloidal range. Such colloids are called macromolecular colloids. These colloids are quite stable and resemble true solutions in many respect e. g., starch dispersed in water.


(ii)
$$\mathrm{CH_3}$$
 $\mathrm{CH_2}$ $\mathrm{CH_3}$ $\mathrm{CH_3}$ KOH $\mathrm{CH_3CH}$ $\mathrm{CH_3CH}$ $\mathrm{CHCH_3}$ $\mathrm{CH_3}$ $\mathrm{CH_2}$ $\mathrm{CH_2}$ $\mathrm{CH_2}$ $\mathrm{CH_2}$ $\mathrm{CH_2}$ $\mathrm{CH_3}$ $\mathrm{CH_2}$ $\mathrm{CH_3}$ $\mathrm{CH$

- **17.** (*i*) Refer to Ans. 15 (*i*) Patna Region 2015.
 - (ii) Refer to Ans. 14 (ii) Foreign 2015.
 - (iii) Wrought iron.
- T_h 353.93 K 353.23 K 0.7 K 18.


Substituting $W_B=1.5$ g, $W_A=90$ g, $T_b=0.7$ K, $K_b=2.52$ K kg mol 1 in the expression $M_B=\frac{K_b-W_B-1000}{T_b-W_A}$, we get

$$M_B = \frac{2.52 \text{ K kg mol}^{-1} - 1.5 \text{ g} - 1000 \text{ g kg}^{-1}}{0.7 \text{ K} - 90 \text{ g}}$$


$$M_B$$
 60 g mol ¹



- (ii) Peptide linkage (C NH)
- (iii) Refer to Answer of Q. 11 (ii) Bhubaneshwar Region 2015.
- **20.** (a) (i) $Co^3 : 3d^6 4s^0$



Hybridisation = d^2sp^3 ; Shape = Octahedral



(b) 'en', as it forms chelate.

21.

	Polymer	Name of Monomers	Structures of Monomers
(<i>i</i>)	Buna-N	Buta-1,3-diene,	$CH_2 = CH CH = CH_2$
		Acrylonitrile	CH ₂ CH CN
(ii)	Bakelite	Phenol,	ОН
		Formaldehyde	НСНО
(iii)	Teflon	Tetrafluoroethene	F ₂ C CF ₂

- 22. (i) Nitrogen due to small size and high electronegativity forms p p multiple bonds and exist as a diatomic molecule. These molecules are held together by weak vander Waal forces and hence N₂ exist as a gas at room temperature. In contrast due to larger size and lower electronegativity P does not form p bond with itself rather it prefer to form single P-P bonds and exist as tetrahedral P₄ molecules. Due to bigger size, the force of attraction holding P_4 molecules are quite strong and cannot broken by the collision of molecules at room temperature and hence solid at room temperature.
 - (ii) As we move down the group from O to Te, the size of central atom goes on increasing and its electronegativity goes on decreasing. Consequently, the bond pairs of electrons tend to lie away from the central atom as we move from H₂O to H₂Te. As a result of this the force of repulsion between the bond pairs decreases as we move from H₂O to H₂Te and hence the bond angle decreases in the same order.

Due to decrease in the tendency of the central atom to undergo sp^3 hybridisation from H_2O to H_2 Te.



- (iii) Halogens have one electron less than the nearest noble gas configuration. Therefore, they have a strong tendency to accept an additional electron to acquire nearest noble gas configuration and hence they have maximum negative electron gain enthalpy.
- 23. Refer to Ans. 22 Allahabad Region 2015.

24. (a)
$$A = \bigcup_{Benzaldehyde}$$
, $B \text{ or } C = \bigcup_{Sodium}$, $C \text{ or } B = \bigcup_{Benzyl}$, $D = \bigcup_{1-Phenyl}$ ethanol

(b) (i) C₆H₅COCH₃ being a methyl ketone gives iodoform test while C₆H₅COCH₂CH₃ does not. C₆H₅COCH₃ 3NaOI C₆H₅COONa CHI_3 2NaOH

Iodoform Acetophenone (Yellow ppt)

(ii) Benzoic acid gives brisk effervescence with NaHCO₃ solution whereas phenol does not.

Η

$$(c) \begin{picture}(200,0) \put(0,0){\line(0,0){100}} \put(0,0){\line(0,0$$

(*i*) CH₃ C H+ HCN CH_3 CN Hydrogen Ethanal cyanide

$$(ii) \begin{tabular}{lll} CH_3 & & & & & & & \\ & \operatorname{CH}_3 & & & & & & \\ & \operatorname{C} & \operatorname{O} + \operatorname{H}_2\operatorname{NNH}_2 & & & & & \\ & \operatorname{Hydrazine} & & & \operatorname{H} & & \\ & \operatorname{Ethanal} & & & & \operatorname{Ethanal oxime} \\ & \operatorname{O} & & & & \\ \end{tabular}$$

- (b) $C_6H_5COCH_3 < CH_3COCH_3 < CH_3CHO$
- (c) CH₃CHO gives yellow precipitate of iodoform when treated with NaOI (NaOH/I) while CH₃CH₂CHO does not.

$$\mathrm{CH_{3}CHO} + \mathrm{3NaOI}$$
 $\mathrm{CHI_{3}}$ + HCOO $\mathrm{Na}^{+} + \mathrm{2NaOH}$ $\mathrm{Iodoform}$ (Yellow ppt.)

25. At anode: Ni Ni²⁺ 2e

At cathode:
$$2Ag^+$$
 2e $2Ag$

$$Ni + 2Ag^+ Ni2+ + 2Ag; n 2$$



$$E_{cell}$$
 $E_{cathode}$ E_{anode} $E_{Ag /Ag}$ $E_{Ni^{2+}/Ni}$

$$E_{cell}$$
 0.80 V - (- 0.25 V) = 1.05 V

Substituting E_{cell} 1.05 V, n 2, $[Ni^2]$ 1 10 2 M, $[Ag^+]$ = 1 10 3 M in Nernst equation,

$$E_{cell}$$
 E_{cell} $\frac{0.059}{n} \log \frac{[\text{Ni}^2]}{[\text{Ag}]^2}$, we get

$$E_{cell} = 1.05 = \frac{0.059}{2} \log \frac{10^{-2}}{(10^{-3})^2}$$

$$E_{cell}$$
 1.05 0.0295 log $(10)^4 = 1.05 - 0.0295$ 4 log 10

$$E_{cell}$$
 1.05 - 0.1180 V = 0.9320 V

$$E_{cell}$$
 0.932 V

0.932 V, n=2, F=96500 C mol $^{-1}$ in the expression, $G=nFE_{cell}$, we get Substituting E_{cell}

$$G \qquad 2 \quad 96500 \; \mathrm{C} \; \, \mathrm{mol} ^{-1} \quad 0.932 \; \, \mathrm{V} \quad -179876 \; \mathrm{J} \; \, \mathrm{mol} ^{-1}$$

$$G - 179.8 \text{ kJ mol}^{-1}$$

(a) Substituting
$$K = 1.06 - 10^{-2} \text{ S cm}^{-1}$$
, $M = 0.1 \text{ mol L}^{-1}$ in the expression, $m = \frac{K - 1000}{M}$, we get
$$\frac{1.06 - 10^{-2} \text{ S cm}^{-1} - 1000 \text{ cm}^3 \text{ L}^{-1}}{0.1 \text{ mol L}^{-1}}$$

$$106 \text{ S cm}^2 \text{ mol}^{-1}$$

o o Na⁺ Na⁺ O = 50.1 S cm² mol
1
 + 76.5 S cm² mol 1 = 126.6 S cm² mol 1

Substituting
$$_m$$
 106 S cm² mol⁻¹, $_m^{o}$ 126.6 S cm² mol⁻¹ in the expression, $\frac{m}{o}$, we get

$$\frac{106 \text{ S cm}^2 \text{ mol}^{-1}}{126.6 \text{ S cm}^2 \text{ mol}^{-1}} \quad 0.8373$$

(b) A primary battery is one in which the redox reaction occurs only once and after use over a period of time battery becomes dead and hence cannot be used again, e.g., dry cell, mercury cell.

A secondary battery after use can be recharged by passing electric current through it and hence can be used over and again, e.g., lead storage battery.

- **26.** (a) (i) In aqueous solutions Ce^{4+} tends to change to Ce^{3+} as +3 oxidation state is more stable and hence Ce⁴⁺ acts as an oxidising agent.
 - (ii) Because of large number of unpaired electrons in their atoms, transition metals have strong interatomic interactions (metallic bonds) resulting in high enthalpy of atomisation.
 - (iii) Refer to Ans. 10 Foreign 2015.
 - (b) (i) Refer to Ans. 14 (b) Delhi 2015.

(ii)
$$\operatorname{Cr_2O_7^2} + 14\operatorname{H}^+ + 6e \qquad 2\operatorname{Cr}^{3+} + 7\operatorname{H_2O}$$

$$\operatorname{Fe}^{2+} \qquad \operatorname{Fe}^{3+} \quad e \quad] \quad 6$$

$$\operatorname{Cr_2O_7^2} + 6\operatorname{Fe}^{2+} + 14\operatorname{H}^+ \qquad 2\operatorname{Cr}^{3+} \quad 6\operatorname{Fe}^{3+} + 7\operatorname{H_2O}$$



OR

- (a) (i) This is due to d-d transition. Under the influence of the ligand attached, the d-orbitals in a transition metal complex split in two sets of orbitals having slightly different energies. When white light falls on these complexes they absorb a particular colour from the white light for the promotion of electrons and remaining colours are emitted. The colour of the complex is due to this emitted radiations.
 - (ii) Cr^{2+} is a strong reducing agent because after the loss of one electron Cr^{2+} becomes Cr^{3+} which has more stable t_{2g}^3 (half-filled) configuration in a medium like water.
 - (iii) The irregularities in the electronic configuration of the actinoids is due to stability of the f^0 , f^7 and f^{14} occupancies of the 5f-orbitals.
- (b) The overall decrease in atomic and ionic radii from lanthanum to lutetium, due to poor shielding effect of 4f electrons is called lanthanoid contraction. The common oxidation state of lanthanoid is +3.





CBSE Examination Paper Guwahati Region-2015 (Code No. 56/3/G)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination paper, Delhi-2015.

1. Write the IUPAC name of the given compound:

- 2. Why is zinc not regarded as a transition element?
- 3. How much charge in Faraday is required for the reduction of 1 mol of Ag⁺ to Ag?
- **4.** Which would undergo S_N1 reaction faster in the following pair:

C₆H₅ CH₂ Br and C₆H₅ CH Br

- **5.** Write the dispersed phase and dispersion medium of smoke.
- **6.** Write two differences between order of a reaction and molecularity of a reaction.
- **7.** Write the structures of the following:
 - $(i) N_2 O_5$
 - (ii) BrF3
- **8.** Write down the IUPAC name of the complex [Co(en)₂ Cl₂] . What type of isomerism is shown by this complex?

OR

Using IUPAC norms write the formulae for the following coordination compounds:

- (i) Tetracarbonylnickel(0)
- (ii) Potassium tetracyanidoferrate(II)
- **9.** What do you understand by depression of freezing point? Derive the relationship between depression of freezing point and molar mass of the solute.
- **10.** Arrange the following:
 - (i) In increasing order of their basic strength

(ii) In increasing order of solubility in water

$$CH_3$$
 NH_2 $(CH_3)_3$ N , CH_3 NH CH_3

- 11. Write the names and structures of the monomers of the following polymers:
 - (i) polystyrene
 - (ii) Nylon-6,6
 - (iii) Terylene

Describe the classification of polymers on the basis of structure.


- **12.** (i) Write the product obtained when D-glucose reacts with HCN.
 - (ii) What type of bonding stabilises the -helix structure of proteins.
 - (iii) Write the name of the disease caused by the deficiency of vitamin B_{12} .
- **13.** Define the following terms:
 - (i) Sorption
 - (ii) Lyophilic colloids
 - (iii) Associated colloids
- **14.** (i) Indicate the principle behind the method used for the refining of zirconium.
 - (ii) What is the role of CO in the extraction of iron?
 - (iii) What is 'copper matte'?
- **15.** Give reasons for the following:
 - (i) NH₃ has a higher boiling point than PH₃.
 - (ii) H_2 Te is more acidic than H_2 S.
 - (iii) Chlorine water on standing loses its yellow colour.
- **16.** Give reasons for the following:
 - (i) Aniline does not undergo Friedel Crafts reaction.
 - (ii) p-methylaniline is more basic than p-nitroaniline.
 - (iii) Acetylation of NH₂ group is done in aniline before preparing its ortho and para compounds.
- 17. (a) Write the hybridisation and shape of the following complexes:
 - (i) $[FeF_6]^3$
 - (ii) [Ni(CO)₄]

(Atomic number : Fe = 26, Ni = 28)

- (b) Out of CN and CO, which ligand forms more stable complex with metal and why?
- 18. Write the structures of the major product in each of the following reactions:

$$(ii) H_2O_2 / OH$$

(ii)
$$CH_3 - CH_2 - CH_3 - CH_3 - KOH(aq.)$$
Br



- **19.** Define the following terms:
 - (i) Crystalline solids
 - (ii) Frenkel defect
 - (iii) n-type semiconductor
- **20.** A first order reaction takes 10 minutes for 25% decomposition. Calculate $t_{1/2}$ for the reaction.

(Given: $\log 2 = 0.3010$, $\log 3 = 0.4771$, $\log 4 = 0.6021$)

21. Vapour pressure of water at 20°C is 17.5 mm Hg. Calculate the vapour pressure of water at 20°C when 15 g of glucose (molar mass = 180 g mol^{-1}) is dissolved in 150 g of water.



- **22.** How do you convert the following:
 - (i) Phenol to 2-hydroxyacetophenone
 - (ii) Ethyl chloride to methoxy ethane
 - (iii) Acetone to 2-methylpropan-2-ol
- Mr. Roy, the principal of one reputed school organized a seminar in which he invited parents and principals to discuss the serious issue of diabetes and depression in students. They all resolved this issue by strictly banning junk food in schools and introducing healthy snacks and drinks like soup, lassi, milk, etc. in school canteens. They also decided to make compulsory half an hour of daily physical activities for the students in the morning assembly. After six months, Mr. Roy conducted the health survey in most of the schools and discovered a tremendous improvement in the health of the students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Roy?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) What are tranquilizers? Give an example.
- (iv) Why is the use of aspartame limited to cold foods and drinks?
- **24.** Calculate e.m.f and G for the following cell at 298 K:

Mg (s) | Mg² (0.01 M) || Ag⁺(0.0001 M) | Ag (s)
Given:
$$E^0_{(Mg^{2+}/Mg)}$$
 2.37V, $E^0_{(Ag^{-}/Ag)}$ 0.80V


(i) The conductivity of 0.001 mol L ¹ solution of CH₃COOH is 4.95 10 ⁵ S cm ¹. Calculate its molar conductivity and degree of dissociation ().

Given:
0
(H) 349.6 S cm² mol 1 and 0 (CH₃COO) 40.9 S cm² mol 1

- (ii) What is a fuel cell? Write its one advantage over other ordinary cells.
- 25. (i) Account for the following:
 - (a) Eu² is a strong reducing agent.
 - (b) Transition metals form coloured compounds.
 - (c) Zn has lowest enthalpy of atomization in 3d series.
 - (ii) Complete the following equations:

(a)
$$\operatorname{KMnO}_4$$
 (b) $\operatorname{Cr}_2\operatorname{O}_7^2$ 14H 6Fe²

- (i) Account for the following:
 - (a) Transition elements form interstitial compounds.
 - (b) Mn^3 (3d⁴) is strongly oxidizing whereas Cr^{2+} (3d⁴) is strongly reducing.
 - (c) Transition metals have high melting points.
- (ii) What is 'misch metal'? Write its one use.
- **26.** (*i*) Write the structures of A, B, C, and D in the following reactions:





- (ii) Distinguish between the following:
 - (a) CH_3 CO $\operatorname{CH}_2\operatorname{CH}_3$ and CH_3 CH_2 CH_2 CHO
 - (b) Ethanal and ethanoic acid
- (iii) Write the structure of 4-chloropentan-2-one.

OR

- (i) Write the structures of the main products when propanal (CH₃ CH₂ CHO) reacts with the following reagents:
 - (a) Zn Hg / conc. HCL
 - (b) H_2N OH/H
 - (c) HCN
- (ii) Arrange the following in the decreasing order of their reactivity towards nucleophilic addition reaction:

HCHO, CH₃ CO CH₃, CH₃ CHO

(iii) Give a simple chemical test to distinguish between the following pairs of compounds:

C₆H₅CHO and C₆H₅ CO CH₃








- 1. 2, 4-dimethylphenol
- 2. As zinc atom has completely filled d-orbitals $(3d^{10})$ in its ground state as well as oxidised state therefore, it is not regarded as transition element.
- 3. 1 F

CH₂

- Br, Because of greater stability of secondary carbocation than primary.
- **5.** Dispersed phase = Solid; Dispersion medium = Gas.
- 6. Difference between Order and Molecularity of a Reaction

S. No.	Order	Molecularity
(<i>i</i>)	Sum of the powers of the concentration of the reactants in a rate law expression.	Number of reacting species taking part in a elementary reaction which must colloid simultaneously in order to bring about a chemical reaction.
(ii)	Determined experimentally.	A theoretical concept.
(iii)	Can be zero or a fraction.	Cannot be zero or a fraction.
(iv)	Applicable to elementary as well as complex reactions.	Applicable only for elementary reactions. For a complex reaction it has no meaning.

7.



Slightly bent "T"

8. Dichlorido bis (ethane-1, 2-diamine) cobalt (III) ion.

Geometrical isomerism. The cis-geometrical isomer will also show optical isomerism.

OR

- (i) [Ni(CO)₄]
- (ii) $K_2[Fe(CN)_4]$
- 9. The decrease in the freezing point of a solvent due to lowering of vapour pressure on addition of a non-volatile solute is called depression in freezing point.

Depression in freezing point (T_f) Freezing point of solvent (T_f) Freezing point of solution (T_f)

It has been found that for a dilute solution $T_f = m \pmod{\text{molality}}$

$$T_f K_f m$$
 ...(i)

where K_f is a proportionality constant known as molal depression constant.

$$m = \frac{W_B}{M_B} = \frac{1000}{W_A}$$

Substituting the value of molality in equation (i), we get

$$T_f = \frac{K_f \quad W_B \quad 1000}{M_B \quad W_A} \quad \text{or} \quad M_B = \frac{K_f \quad W_B \quad 1000}{T_f \quad W_A}$$



10. (*i*)
$$C_6H_5NH_2 < CH_3$$
 CH_2 $NH_2 < CH_3NHCH_3$ (*ii*) $(CH_3)_3N < (CH_3)_2NH < CH_3NH_2$


11.

	Polymer	Name of Monomers	Structures of Monomers
(i)	Polystyrene	Styrene	\sim CH = CH ₂
(ii)	Nylon-6,6	Adipic acid Hexamethylene-diamine	$HOOC$ $(CH_2)_4$ $COOH$ H_2N $(CH_2)_6$ NH_2
(iii)	Terylene	Ethylene glycol Terephthalic acid	HOOC—COOH

OR

Classification based on structure of polymer:

- (i) Linear polymers: These polymers consist of long and straight chains, e.g., high density polythene, polyvinyl chloride, nylon, etc.
- (ii) Branched chain polymers: These polymers contain linear chains having some branches, e.g., low density polythene, glycogen, etc.
- (iii) Cross-linked or network polymers: In this type of polymers, the initially formed linear polymer chains are joined together to form three dimensional network structure. Due to presence of cross links these polymers are also called cross-linked polymers, e.g., bakelite, melamine, etc.



CHO
$$CH_{OH}$$

12. (i) (CHOH)₄ $CH_{2}OH$

CH₂OH

Glucose Glucose cyanohydrin

- (ii) Intermolecular hydrogen bonding
- (iii) Pernicious anaemia.
- (i) The process in which adsorption and absorption take place simultaneously is called sorption. Certain dyes get adsorbed as well as absorbed on the fibers resulting in sorption.
 - (ii) The colloidal solutions in which the particles of the dispersed phase have a great affinity for the dispersion medium are known as lyophilic colloids, e.g., starch, gum, gelatin, etc.
 - (iii) There are certain substances which at low concentrations behave as normal strong electrolytes, but at higher concentrations exhibit colloidal behaviour due to formation of aggregates of small particles called micelles and such type of colloids are known as associated colloids, e.g., soap, synthetic detergent.



- 14. (a) Impure Zr reacts with I₂ to form volatile ZrI₄ which when heated at higher temperature decomposes to give pure Zr.
 - (b) CO acts as a reducing agent.
 - (c) It is a mixture of Cu₂S and FeS.
- 15. (i) Due to small size and high electronegativity of N, NH₃ undergoes extensive intermolecular H-bonding and hence exist as an associated molecule. In contrast due to low electronegativity of P, PH3 does not undergo H-bonding and thus exist as a discrete molecule. That is, why boiling point of NH3 is higher than PH₃.
 - (ii) Because bond dissociation enthalpy of H-Te bond is less than H—S bond as size of Te is larger than that of S.
 - (iii) Cl₂ + H₂O HCl + HOCl

Due to formation of hydrochloric acid and hypochlorous acid chlorine water loses yellow colour on standing.

16. (i) Aniline being a Lewis base reacts with Lewis acid AlCl₃ (catalyst) to form a salt.

Due to presence of positive charge on N-atom in the salt, the group NH₂AlCl₃ reduces electron density in benzene ring and acts as a strong deactivating group and hence aniline does not undergo Friedal craft reaction.

- CH_3 being electron donating group released electrons, stabilizes the conjugate acid of p-methyl (ii) NO₂ being electron withdrawing, withdraws electrons, destabilizes conjugate acid of p-nitroaniline that is why p-methyl aniline is stronger base than p-nitroaniline.
- (iii) Acetylation of NH₂ group reduces the activating effect of it and thus prevail the formation of di and tri-substitution products.
- **17.** (a) (i) Refer to Ans. 17 (a) (i) Patna Region 2015.
 - (ii) Ni : $3d^8 4s^2$

(b) CO, because of synergic bonding.

18. (i)
$$CH_3$$
 CH_2 CH_2 CH_3 CH_3 CH_4 CH_5 CH_5 CH_5 CH_6 CH_7 CH_8 CH_8



- 19. (i) Crystalline solids are those solids in which the constituent particles (atoms, molecules or ions) are arranged in a definite geometrical pattern in three dimensional space so that there is short range as well as long range order of constituent particles, e.g., Ag, C (diamond), NaCl etc.
 - (ii) When some ions, usually cations are missing from the lattice sites and occupy the interstitial sites so that electrical neutrality as well as stoichiometry is maintained, it is called Frenkel defect. Silver halides AgCl, AgBr, AgI exhibit Frenkel defect.
 - (iii) Si or Ge when doped with atoms of pentavalent element like P, As, free electrons become available which increase the conductivity. Such type of semiconductors are called *n*-type semiconductors.
- **20.** Refer to Ans. 24 (b) Foreign 2015.
- **21.** Substituting P_A 17.5 mm Hg, W_b 15 g, W_A 150 g, M_B 180 g mol ¹, M_A 18 g mol ¹ in the expression, $\frac{P_A}{P_A} = \frac{P}{M_B} = \frac{M_A}{M_B}$, we get

$$\frac{17.5}{17.5} \frac{P}{180} \frac{15}{180} \frac{18}{150} \quad 0.01$$

$$17.5 - P \quad 0.175$$

$$P \quad 17.5 - 0.175$$

$$P \quad 17.325 \text{ mm Hg}$$

22.

or

$$(ii) \ \, \text{CH}_{3} \ \, \text{CH}_{2} \ \, \text{CI} \\ \text{Ethyl chloride} \ \, \text{CH}_{3} \ \, \text{O Na} \\ -\text{NaCl} \ \, \text{CH}_{3} \ \, \text{CH}_{2} \ \, \text{O} \ \, \text{CH}_{3} \\ \text{Methoxy ethane} \ \, \text{OH} \\ \\ (iii) \ \, \text{CH}_{3} \ \, \text{C} \ \, \text{CH}_{3} \$$

- 23. Refer to Ans. 23 Delhi region 2015.
- **24.** At anode: At cathode : $Ag^{+} e Ag = 2$ $Mg + 2Ag^{+} Mg^{2+} + 2Ag = 0.80 \text{ V} - (-2.37 \text{ V}) = 3.17 \text{ V}$ $E_{cell} E_{cathode} E_{Ag /Ag} E_{Mg^{2+}/Mg} = 0.80 \text{ V} - (-2.37 \text{ V}) = 3.17 \text{ V}$



Substituting
$$E_{cell}$$
 3.17 V, n 2, [Mg 2] 1 10 2 M, [Ag $^+$] = 1 10 4 M in Nernst equation for above cell reaction, E_{cell} E_{cell} $\frac{0.059}{n}$ log $\frac{[\text{Mg}^2]}{[\text{Ag}]^2}$, we get

$$E_{cell}$$
 3.17 $\frac{0.059}{2} \log \frac{10^{-2}}{(10^{-4})^2}$

$$E_{cell}$$
 3.17 0.0295 log 10⁶

$$E_{cell}$$
 3.17 - 0.177 V = 2.993 V

$$E_{cell}$$
 2.993 V

96500 C mol 1 , E_{cell} 2.993 V in the expression, G nFE_{cell} , we get 2, *F* Substituting *n*

$$G = nFE_{cell}$$
 2 96500 C mol ¹ 2.993 V

$$G$$
 -577649 J mol 1 -577.649 kJ mol 1

(a) Substituting
$$K = 4.95 - 10^{-5}$$
 S cm⁻¹, $M = 0.001$ mol L⁻¹ in the expression, $_{m} = \frac{K - 1000}{M}$, we get

$$^{m} \quad \frac{4.95 \quad 10^{-5} \text{ S cm}^{-1} \quad 1000 \text{ cm}^{3} \text{ L}^{-1}}{0.001 \text{ mol L}^{-1}}$$

$$m \text{ (CH}_3\text{COOH)}$$
 H^+ $CH_3\text{COO} = 349.6 \text{ S cm}^2 \text{ mol}^{-1} + 40.9 \text{ S cm}^2 \text{ mol}^{-1}$
= 390.5 S cm² mol⁻¹

Substituting $_m$ 49.5 S cm² mol¹, $_m$ 390.5 S cm² mol¹ in the expression, $\frac{m}{m}$, we get

$$\frac{49.5 \text{ S cm}^2 \text{ mol}^{-1}}{390.5 \text{ S cm}^2 \text{ mol}^{-1}} \quad 0.1267$$

(b) Galvanic cells which are designed to convert the energy of combustion of fuels like hydrogen, methane, methyl alcohol etc. directly into electrical energy are called fuel cells.

Advantages of fuel cell:

- (i) It is pollution free.
- (ii) It has high efficiency of 60-70%.
- **25.** (a) (i) Eu^{2+} is a strong reducing agent as it changes to more stable Eu^3 state by losing an electron.
 - (ii) Refer to Ans. 26 (OR) (a) (i) Kerala region 2015.
 - (iii) Because in the formation of metallic bonds no electron from 3d orbitals are involved in case of zinc as it has completely filled d-orbitals $(3d^{10} 4s^2)$, while in all other metals of 3d series electrons from the *d*-orbitals are always involved in the formation of metallic bonds.

(ii)
$$\operatorname{Cr_2O_7^2} + 14\operatorname{H}^+ + 6e \qquad 2\operatorname{Cr}^{3+} + 7\operatorname{H_2O}$$

$$\operatorname{Fe}^{2+} \qquad \operatorname{Fe}^{3+} \qquad e \quad] \quad 6$$

$$\operatorname{Cr_2O_7^2} + 6\operatorname{Fe}^{2+} + 14\operatorname{H}^+ \qquad 2\operatorname{Cr}^{3+} \quad 6\operatorname{Fe}^{3+} + 7\operatorname{H_2O}$$



- (a) (i) Transition elements form interstitial compounds because small size atoms like H, B, C, N etc. can easily occupy interstitial sites in the crystal lattices transition elements, e.g., TiC, Fe₃H, Mn₄N etc.
 - (ii) Mn³⁺ (3 d^4) is strongly oxidising as a change from 3 d^4 to 3 d^5 configuration results a more stable half filled d^5 configuration.

While $\operatorname{Cr}^{2+}(3d^4)$ is strongly reducing as Cr^{2+} to Cr^{3+} change result a more stable half filled t_{2g}^3 configuration.

- (iii) The high melting points of transition metals attributed to the involvement of greater number of 1) d orbitals in addition to ns electrons in the metallic bonding.
- (b) A misch metal is an alloy which contain lanthanoid metals (95%), iron (5%) and traces of S, C, Ca and Al. Misch metal is used in magnesium based alloy to produce bullets, shell and lighter flint.

- Cl; B CH_3 C H; C CH_3 CH CH_3 ; D CH_3 CH CH_2 Ethanol **26.** (*a*) A CH₃ C
 - (b) (i) Being an aldehyde CH₃CH₂CH₂CHO reduces Fehling solution to red ppt. of Cu₂O while CH₃COCH₂CH₃ being a ketone does not.

 CH_2 oxide (red ppt.)

(ii) Ethanoic acid being an acid gives brisk effervescence with NaHCO3 solution whereas ethanal does

 $CH_3COO Na^+ + CO_2 + H_2O$ CH₃COOH + NaHCO₃ Ethanoic acid Sodium bicarbonate

O Cl

(c) CH₃ CH CH₂ C CH₃

(a) (i)
$$CH_3$$
 CH_2 $CHO + 4[H]$ CH_3 CH_3 CH_3 CH_4 $CH_3 + H_2 CH_3$ $CH_3 + H_4 CH_5$ CH_5 CH_5 CH_6 CH_7 CH_8 CH_8 CH_8 CH_8 CH_8 CH_9 CH_9

Η Η

(iii)
$$CH_3$$
 CH_2 C $O+HCN$ CH_3 CH_2 C OH

CN

Propanal cyanohydrin

- (b) $HCHO > CH_3$ $CHO > CH_3COCH_3$
- (c) C₆H₅CHO being an aldehyde gives silver mirror when treated with Tollen's reagent while C₆H₅COCH₃ being a ketone does not.

$$\begin{array}{cccc} C_6H_5\text{CHO} + 2[\text{Ag}(\text{NH}_3)_2]^+ + 3\text{OH} & C_6H_5\text{COO} & +2\text{Ag} & +2\text{H}_2\text{O} + \text{NH}_3 \\ & & \text{Silver} & & \text{mirror} \end{array}$$



CBSE Examination Paper Panchkula Region -2015 (Code No. 56/1/C)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper, Delhi-2015.

1. Write a method by which lyophobic colloids can be coagulated.

- 2. What is the formula of a compound in which the element Y forms hcp lattice and atoms of X occupy $2/3^{rd}$ of tetrahedral voids?
- 3. Out of white phosphorus and red phosphorus, which one is more reactive and why?
- **4.** Write the IUPAC name of the given compound:

$$\begin{array}{c} \operatorname{HO}-\operatorname{CH}_2-\operatorname{CH}-\operatorname{CH}_2-\operatorname{OH} \\ \mid \\ \operatorname{CH}_3 \end{array}$$

5. Which would undergo S_N1 reaction faster in the following pair and why?

$$CH_3$$
 CH_3 CH_3 CH_3 CH_3 CH_3 CH_3 CH_3

6. State Henry's law. Why do gases always tend to be less soluble in liquids as the temperature is raised?

OR

State Raoult's law for the solution containing volatile components. Write two differences between an ideal solution and a non-ideal solution.

7. (a) Following reactions occur at cathode during the electrolysis of aqueous sodium chloride solution:

Na⁺
$$(aq) + e^{-}$$
 Na (s) $E^{0} = -2.71 \text{ V}$
H⁺ $(aq) + e^{-}$ $\frac{1}{2} \text{H}_{2} (g)$ $E^{0} = 0.00 \text{ V}$

On the basis of their standard reduction electrode potential (E^0) values, which reaction is feasible at the cathode and why?

- (b) Why does the cell potential of mercury cell remain constant throughout its life?
- **8.** Why do the transition elements have higher enthalpies of atomisation? In 3d series (Sc to Zn), which element has the lowest enthalpy of atomisation and why?
- 9. (i) Write down the IUPAC name of the following complex: $[Co(NH_3)_5(NO_2)] (NO_3)_2$
 - (*ii*) Write the formula for the following complex: Potassium tetracyanidonickelate(II)
- 10. Name the reagents used in the following reactions:

(i)
$$CH_3 - CO - CH_3 \xrightarrow{?} CH_3 - C - CH_3$$

OH

$$(ii)$$
 CH₃ – COOH $^{?}$ CICH₂ – COOH

11. Calculate the amount of CaCl₂ (molar mass = 111 g mol⁻¹) which must be added to 500 g of water to lower its freezing point by 2 K, assuming CaCl2 is completely dissociated.

 $(K_f \text{ for water} = 1.86 \text{ K kg mol}^{-1}).$

- 12. An element with density 10 g cm^{-3} forms a cubic unit cell with edge length of 3×10^{-8} cm. What is the nature of the cubic unit cell if the atomic mass of the element is 81 g mol^{-1} ?
- 13. Calculate emf of the following cell at 25°C:

Sn | Sn²⁺ (0.001 M) || H⁺ (0.01 M) | H₂ (g) (1 bar) | Pt (s) $E_{(Sn^2/Sn)}^0$ 0.14 V, $E_{(H/H_2)}^0$ 0.00 V

- **14.** Give reasons for the following observations:
 - (i) Physisorption decreases with increase in temperature
 - (ii) Addition of alum purifies the water.
 - (iii) Brownian movement provides stability to the colloidal solution.
- **15.** (i) Name the method used for the refining of zirconium.
 - (ii) What is the role of CO in the extraction of iron?
 - (iii) Reduction of metal oxide to metal becomes easier if the metal obtained is in liquid state. Why?
- **16.** (a) How would you account for the following:
 - (i) The chemistry of actinoids is more complicated as compared to lanthanoids.
 - (ii) Transition metals form complex compounds.
 - (b) Complete the following equation:

 $2 \text{ MnO}_4^- + 6 \text{ H}^+ + 5 \text{ SO}_3^2$

- 17. (i) Draw the geometrical isomers of complex $[Co(en)_2Cl_2]^+$.
 - (ii) On the basis of crystal field theory, write the electronic configuration for d^4 ion if $_0 > P$.
 - (iii) $[NiCl_4]^{2-}$ is paramagnetic while $[Ni(CO)_4]$ is diamagnetic, though both are tetrahedral. Why? (Atomic number of Ni = 28)
- **18.** How do you convert the following:
 - (i) Prop-1-ene to 1-fluoropropane
 - (ii) Chlorobenzene to 2-chlorotoluene
 - (iii) Ethanol to propanenitrile

OR

Write the main products when

- (i) n-butyl chloride is treated with alcoholic KOH.
- (ii) 2, 4, 6-trinitrochlorobenzene is subjected to hydrolysis.
- (iii) methyl chloride is treated with AgCN.
- **19.** Give reason for the following:
 - (i) o-nitrophenol is more acidic than o-methoxyphenol.
 - (ii) Butan-1-ol has a higher boiling point than diethylether.
 - (iii) (CH₃)₃ C O CH₃ on reaction with HI gives (CH₃)₃ C I and CH₃ OH as the main products and not $(CH_3)_3 C - OH$ and $CH_3 - I$.
- **20.** Predict the products of the following reactions:

(i)
$$CH_3 - C = O \xrightarrow{HCN} ?$$

 CH_3
(ii) $C_6H_5 - CH_2 - CH_3$ (a) $KMnO_4 / KOH$?

(b) H

NH₃/ (iii) CH₃COOH

- 21. Write the names and structures of the monomers of the following polymers:
 - (i) Nylon-6

(ii) Novolac

- (iii) Buna-N
- **22.** (*i*) Which one of the following is a polysaccharide:

Starch, Maltose, Fructose, Glucose

- (ii) What is the difference between native protein and denatured protein?
- (iii) Write the name of the vitamin responsible for the coagulation of blood.
- 23. Seeing the growing cases of diabetes and depression among young children, Mr. Lugani, the principal of one reputed school organized a seminar in which he invited parents and principals. They all resolved this issue by strictly banning junk food in schools and introducing healthy snacks and drinks like soup, lassi, milk, etc., in school canteens. They also decided to make compulsory half an hour of daily physical activities for the students in the morning assembly. After six months, Mr. Lugani conducted the health survey in most of the schools and discovered a tremendous improvement in the health of students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Lugani?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) What are antidepressant drugs? Give an example.
- (iv) Name the sweetening agent used in the preparation of sweets for a diabetic patient.
- **24.** (a) Illustrate the following reactions giving suitable example in each case:
 - (i) Hoffmann bromamide degradation reaction
 - (ii) Diazotisation
 - (iii) Gabriel phthalimide synthesis
 - (b) Distinguish between the following pairs of compounds:
 - (i) Aniline and N-methylaniline
- (ii) (CH₃)₂NH and (CH₃)₃N

OR

- (a) Write the structures of main products when benzene diazonium chloride $(C_6H_5N_2^+Cl^-)$ reacts with the following reagents:
 - (i) CuCN/KCN

(ii) H₂O

- (iii) CH₃CH₂OH
- (b) Arrange the following:
 - (i) $C_2H_5NH_2$, C_2H_5OH , $(CH_3)_3N$ in the increasing order of their boiling point.
 - (ii) Aniline, p-nitroaniline, p-methylaniline in the increasing order of their basic strength.
- **25.** (a) What is rate of reaction? Write two factors that affect the rate of reaction.
 - (b) The rate constant of a first order reaction increases from 4×10^{-2} to 8×10^{-2} when the temperature changes from 27°C to 37°C. Calculate the energy of activation (E_a) .

$$(\log 2 = 0.301, \log 3 = 0.4771, \log 4 = 0.6021).$$

OR

(a) For a reaction A + B

P, the rate is given by

Rate = $k[A] [B]^2$

- (i) How is the rate of reaction affected if the concentration of B is doubled?
- (ii) What is the overall order of reaction if A is present in large excess?
- (b) A first order reaction takes 23.1 minutes for 50% completion. Calculate the time required for 75% completion of this reaction.

$$(\log 2 = 0.301, \log 3 = 0.4771, \log 4 = 0.6021)$$

- **26.** (a) Account for the following:
 - (i) Interhalogens are more reactive than pure halogens.



- (ii) N_2 is less reactive at room temperature.
- (iii) Reducing character increases from NH₃ to BiH₃.
- (b) Draw the structures of the following:
 - (i) H₄P₂O₇ (Pyrophosphoric acid) (ii) XeF₄

OR

- (a) Which poisonous gas is evolved when white phosphorus is heated with conc. NaOH solution? Write the chemical equation involved.
- (b) Which noble gas has the lowest boiling point?
- (c) Fluorine is a stronger oxidising agent than chlorine. Why?
- (d) What happens when H_3PO_3 is heated?
- (e) Complete the equation:

 $PbS + O_3$







- 1. The coagulation of lyophobic sol can be carried out-in the following ways:
 - (i) By adding excess of electrolyte
 - (ii) By boiling
 - (iii) By persistent dialysis
 - (iv) By mixing two oppositely charged sols.
 - (v) By electrophoresis.
- 2. Let the number of atoms of element Y in hcp lattice N

Number of tetrahedral voids

Number of atoms of element $X = \frac{2}{3} = 2N = \frac{4}{3}N$

$$X: Y = \frac{4}{3} N: N = 4:3$$

Formula of the compound X_4Y_3

- 3. White phosphorus, because of angular strain in the P_4 molecule where the bond angles are only 60° .
- 2-methylpropane-1, 3-diol.

 CH_3

5. CH₂ C CH₃, tertiary halide reacts faster than primary halide because of greater stability of Br

3°-carbocation.

6. It states that the mole fraction of a gas in the solution is proportional to the partial pressure of the gas over the solution.

As dissolution is an exothermic process, so according to Le Chatelier's principle solubility will decrease with increase of temperature

OR

Raoult's law states that for a solution of volatile liquids the partial vapour pressure of each component is directly proportional to its mole fraction.

Differences between Ideal and non-Ideal solutions

S.No.	Ideal solution	Non-Ideal solution		
(<i>i</i>)	Obeys Raoult's law over entire range of concentration	Does not obey Raoult's law over entire range of		
	$i.e., P_A \qquad P_A x_A; P_B \qquad P_B x_B$	concentration, i.e., $P_A P_A x_A$; $P_B P_B x_B$		
(ii)	$_{\text{mix}} H$ 0; $_{\text{mix}} V$ 0	$_{ ext{mix}} H$ 0; V 0		
(iii)	Does not form azeotrope.	Forms azeotrope.		

- $\frac{1}{2}$ H₂(g) is feasible at cathode as it has higher standard reduction (a) The reaction H (aq) potential than other reaction.
 - (b) As the overall reaction does not involve any ion in the solution whose concentration changes during its life period.
- **8.** Refer to Ans. 26 (*a*) (*ii*) Chennai Region 2015.

 $\text{Zn}(3d^{10}4s^2)$, as in the formation of metallic bond, no electrons from 3d orbitals are involved in case of zinc therefore weak metallic bonds exist in zinc.

- **9.** (i) Pentaamminenitrito-N-cobalt (III) nitrate
 - (ii) $K_2[Ni(CN)_4]$



- **10.** (*i*) CH₃MgBr; H₂O
 - (ii) Cl₂/red P; H₂O
- 11. $CaCl_2$ Ca^2 2Cl , i 3

Substituting W_A 500 g, M_B 111 g mol 1 , K_f 1.86 K kg mol $^{-1}$, i 3, in the expression,

$$W_B = \frac{T_f \quad W_A \quad M_B}{i \quad K_f \quad 1000}$$
, we get

$$W_B = \frac{2 \text{ K} \quad 500 \text{ g} \quad 111 \text{ g mol}^{-1}}{3 \quad 1.86 \text{ K kg mol}^{-1} \quad 1000 \text{ g kg}^{-1}} = 19.89 \text{ g}$$

12. Substituting $d = 10 \text{ g cm}^{-3}$, $a = 3 = 10^{-8} \text{ cm}$, $M = 81 \text{ g mol}^{-1}$, $N_A = 6.022 = 10^{23} \text{ mol}^{-1}$ in the expression, $z = \frac{d - a^3 - N_A}{M}$, we get

$$z = \frac{10 \text{ g cm}^3 \quad (3 \quad 10^{-8} \text{ cm})^3 \quad 6.022 \quad 10^{23} \text{ mol}^{-1}}{81 \text{ g mol}^{-1}} = 2.007$$

As thus are 2 atoms of the element per unit cell. Hence, the unit cell must be body centred cubic (bcc).

 Sn^2 2e Sn **13.** At anode :

At cathode :
$$\frac{2H}{Sn} = \frac{2e}{Sn} = \frac{H_2}{Sn^2 + H_2}$$
; $n = 2$

$$E_{cell}^0 = E_{cathode}^0 = E_{anode}^0 = E_{H^+/H_2} = E_{Sn^{2+}/Sn}^0 = 0 - (-0.14 \text{ V}) = 0.14 \text{ V}$$

$$E_{\text{cell}}^{\text{o}}$$
 $E_{\text{cathode}}^{\text{o}}$ $E_{\text{anode}}^{\text{o}}$ $E_{\text{H}^{+}/\text{H}_{2}}^{\text{o}}$ $E_{\text{Sn}^{2+}/\text{Sn}}^{\text{o}} = 0 - (-0.14 \text{ V}) = 0.14 \text{ V}$

Substituting
$$E_{\text{cell}}^{\text{o}}$$
 0.14 V, n 2, $[\text{Sn}^2]$ 1 10 3 M, $[\text{H}]$ 1 10 2 M

In the equation
$$E_{\text{cell}} = E_{\text{cell}}^{\text{o}} = \frac{0.059}{n} \log \frac{[\text{Sn}^2]}{[\text{H}]^2}$$
, we get

$$E_{\text{cell}} = 0.14 = \frac{0.059}{2} \log \frac{10^{-3}}{(10^{-2})^2}$$

$$E_{\text{cell}}$$
 0.14 0.0295 log 10 = 0.1105 V

$$E_{\rm cell}$$
 0.11 V

- 14. (i) As physisorption is an exothermic process so according to Le Chatelier's principle increase in temperature will decrease the physisorption.
 - (ii) The Al³ ions furnished by alum help in the coagulation of negatively charged colloidal particles present in water as impurities.
 - (iii) The Brownian movement is due to the unbalanced bombardment of the colloidal particles of the molecules of dispersion medium. It has a stirring effect which does not permit the particles to settle and thus responsible for stability of sols.
- **15.** (*i*) van Arkel method.
 - (ii) CO acts as a reducing agent.
 - (iii) Entropy is higher when a metal is in the liquid state than when it is in the solid state. The value of entropy change (S) of reduction process is more on +ve side when metal formed is in liquid state than the metal oxide being reduced in solid state. Thus, the value of G is more on – ve side and the reduction becomes easier.
- **16.** (a) (a)

$E^{0}_{(M^{2+}/M)}$	Cr	Mn	Fe	Co	Ni	Cu
-(M ²⁺ /M)	-0.91	-1.18	-0.44	-0.28	-0.25	+0.34




From the given data E^0 values, answer the following questions:


- (i) Why is $E_{(Cu^{2+}/Cu)}^{o}$ value exceptionally positive?
- (ii) Why is $E_{(Mn^{2+}/Mn)}^{0}$ value highly negative as compared to other elements?
- (iii) Which is a stronger reducing agent Cr²⁺ or Fe²⁺? Give reason.
- (ii) Refer to Ans. 21(a) (ii) Allahabad Region 2015.

(b)
$$MnO_4$$
 8H 5e Mn^2 4H₂O] 2
 SO_3^2 H₂O SO_4^2 2H 2e] 5
 $2MnO_4$ 5SO $_3^2$ 6H $2Mn^2$ 5SO $_4^2$ 3H₂O





- (ii) $t_{2g}^4 e_g^0$.
- (iii) Because in $[NiCl_4]^2$, Cl being a weak field ligand unable to pair up the d electrons whereas in [Ni(CO)₄], CO being a strong field ligand pair up the electrons.



- Alc. KCN CH₃ CH_2 CH₂ CH_2 (iii) CH₃ Ethanol Propanenitrile
- OR CH_2 CH_2 Cl + KOH (alc.) $CH_2 + KCl + H_2O$ (i) CH₃ CH_2 CH_3 CH₂ CH *n* - Butyl chloride But -1-ene

(ii)
$$O_2N$$
 O_2N $O_$

(iii) CH_3 Cl + AgCN (alc.)N C + AgCl CH_3 Methyl isocyanide Methylchloride (Major product)



- **19.** (i) R and I effect of NO₂ group decreases the electron density in O—H bond and make loss of proton easy in o-nitrophenol whereas R effect of —OCH₃ group increases the electron density in O—H bond and makes release of proton difficult in o-methoxyphenol. That is, why o-nitrophenol is stronger acid than o-methoxyphenol.
 - (ii) Butan-1-ol undergoes intermolecular hydrogen bonding due to presence of a hydrogen atached to the electronegative oxygen atom and hence exists as associated molecule. On the other hand, diethyl ether does not form H-bonds. Due to this reason butan-1-ol has higher boiling point than diethyl ether.
 - (iii) Refer to Ans. 19 (iii) Allahabad Region-2015.

CN

20. (*i*) CH_3 C OH (ii) C₆H₅COOH Benzoic acid

(iii) CH₃CONH₂ Ethanamide

 CH_3 Propanone cyanohydrin

21.

	Polymer	Name of Monomers	Structure of Monomers
(i)	Nylon-6	Caprolactum	$H_{2}C^{N}C=0$ $H_{2}C$ CH_{2} $H_{2}C-CH_{2}$
(ii)	Novolac	Phenol,	OH
(iii)	Buna-N	Formaldehyde Butan-1, 3-diene Acrylonitrile	H C H CH ₂ CH CH CH ₂ CH ₂ CH CN

- 22. (i) Starch
 - (ii) Protein found in a biological system with unique three-dimensional structure and biological activity is called native protein.

When a protein in its native form is subjected to change such as change in temperature, change in pH, its 2° and 3° structures are destroyed and it loses its biological activity. The protein thus formed is called denatured protein.

- (iii) Vitamin K.
- 23. Refer to Ans. 23 Patna Region 2015.
- (i) Hoffmann Bromamide degradation reaction:

(ii) Diazotisation:

$$NH_2 + NaNO_2 + 2HCl \xrightarrow{273-278 \text{ K}} N_2Cl + NaCl + 2H_2O$$



(iii) Gabriel Phthalimide synthesis:

(b) (i) Aniline being a primary amine gives carbylamine test, i.e., when heated with an alcoholic solution of KOH and CHCl₃, it gives four smell of phenyl carbylamine whereas N-methylaniline being a secondary amine does not give this test.

Aniline
$$NH_2 + CHCl_3 + 3KOH(alc.)$$
 \longrightarrow $NC + 3KCl + 3H_2O$ Phenyl carbylamine (foul smell)

(ii) (CH₃)₂ NH and (CH₃)₃ N can be distinguished using Hinsberg's reagent, i.e., benzene sulphonyl chloride. When treated with Hinsberg's reagent, (CH₃)₂ NH gives N, N-dimethyl benzene sulphonamide which is insoluble in aqueous KOH solution.

(CH₃)₃ N, on the other hand does not react with Hinsberg's. reagent.

OR

(a) (i)
$$N_2Cl$$
 $CuCN/KCN$ $CN + N_2$

Benzene diazonium chloride

(ii) $N_2Cl + H_2O$ 283 K $OH + N_2 + Cl$

Phenol

(iii) $N_2Cl + CH_3 - CH_2 - OH$ $OH + CH_3CHO + N_2 + HO$

Benzene

- (b) (i) $(CH_3)_3 N < C_2H_5NH_2 < C_2H_5OH$
 - (ii) p-Nitroaniline < aniline < p-methylaniline
- **25.** (*a*) Refer to Ans. 7 Patna Region 2015.
 - (b) Substituting k_1 4 10 ², k_2 8 10 ², T_1 (27 273)K 300K, T_2 (37 310 K, $R = 8.314 \text{ J K}^{-1} \text{mol}^{-1}$, in the expression $\log \frac{k_2}{k_1} = \frac{E_a}{2.303 R} = \frac{T_2 - T_1}{T_1 T_2}$, we get



$$\log \frac{8 \quad 10^{-2}}{4 \quad 10^{-2}} \quad \frac{E_a}{2303 \quad 8.314} \quad \frac{310 \quad 300}{300 \quad 310}$$

$$\log 2 \quad \frac{E_a}{19.147} \quad \frac{10}{300 \quad 310}$$

$$E_a \quad \log 2 \quad 19.147 \quad 300 \quad 31 \quad 0.3010 \quad 19.147 \quad 300 \quad 31 \quad 53598.19 \text{ J mol}^{-1}$$

$$E_a \quad 53.6 \text{ kJ mol}^{-1}$$


OR

(a) Refer to Ans. 26 (OR) (a) Delhi Region 2015.

(b) 75% of
$$[R]_0$$
 $\frac{75 \quad [R]_0}{100}$ 0.75 $[R]_0$ $[R] \quad [R]_0$ 0.75 $[R]_0$ 0.25 $[R]_0$ $k \quad \frac{0.693}{t_{1/2}} \quad \frac{0.693}{23.1 \text{ min}} \quad 0.03 \text{ min}^{-1}$

Substituting $k = 0.03 \text{ min}^{-1}$, $[R] = 0.25 [R]_0$ in the expression $t = \frac{2.303}{k} \log \frac{[R]_0}{[R]}$, we get $t = \frac{2.303}{0.03} \log \frac{[R]_0}{0.25 [R]_0} = \frac{2.303}{0.03} \log 4 = \frac{2.303}{0.03} = 0.602$

- (i) This is because X X bond in interhalogens is weaker than X—X bond in halogens (except F—F bond).
 - (ii) Due to presence of triple bond between two N-atoms the bond enthalpy (941.4 kJ mol 1) is very high. Hence, nitrogen is less reactive at room temperature.
 - (iii) Because E—H bond dissociation ethalpy decreases from NH₃ to BiH₃ as the size of central atom E increases from NH₃ to BiH₃.



(a) Phosphine (PH₃).

$$\begin{array}{ccc} P_4 & + & 3 \text{NaOH} + 3 \text{H}_2 \text{O} & & P \text{H}_3 & + & 3 \text{NaH}_2 \text{PO}_2 \\ \text{White} & & & \text{Phosphine} \end{array}$$

- (b) Helium (He).
- (c) It is due to the following reasons:
 - (i) Low enthalpy of dissociation of F—F bond.
 - (ii) high hydration enthalpy of F.
- (d) H₃PO₃ disproportionates on heating to give H₃PO₄ and PH₃.

$$4H_3PO_3$$
 $3H_3PO_4 + PH_3$





CBSE Examination Paper Patna Region-2015 (Code No. 56/1/P)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination paper, Delhi-2015.

1. How much charge is required for the reduction of 1 mol of Zn^{2+} to Zn?

- 2. Write the dispersed phase and dispersion medium of butter.
- 3. Zn^{2+} salts are white while Cu^{2+} salts are coloured. Why?
- **4.** Write the IUPAC name of the given compound:

$$CH_2 = C - CH_2 - OF$$

$$CH_3$$

5. Which would undergo S_N1 reaction faster in the following pair:

- **6.** Why does a solution containing non-volatile solute have higher boiling point than the pure solvent? Why is elevation of boiling point a colligative property?
- 7. Define rate of reaction. Write two factors that affect the rate of reaction.
- **8.** Write the structures of the following molecules:
 - (i) H₂SO₃
 - (ii) XeOF₄
- **9.** Write down the IUPAC name of the complex $[Pt(en)_2CI_2]^{2+}$. What type of isomerism is shown by this complex?

OR

Using IUPAC norms write the formulae for the following coordination compounds:

- (i) Hexaamminecobalt(III) chloride
- (ii) Potassium tetrachloridonickelate(II)
- **10.** Arrange the following in increasing order of their basic strength:

(i)
$$C_6H_5 - NH_2, C_6H_5 - CH_2 - NH_2, C_6H_5 - NH - CH_3$$

(ii)
$$NH_2$$
 NH_2 NH_2 NH_2 NH_2 NH_2 NH_2 NH_2 NO_2 NO

- 11. Calculate the freezing point of the solution when 31 g of ethylene glycol ($C_2H_6O_2$) is dissolved in 500 g of water. (K_f for water = 1.86 K kg mol⁻¹)
- **12.** Define the following terms:
 - (i) Primitive unit cells
- (ii) Schottky defect
- (iii) Ferromagnetism

$$(\log 2 = 0.301, \log 3 = 0.4771, \log 4 = 0.6021)$$

- **14.** Define the following terms:
 - (i) Brownian movement
- (ii) Peptization
- (iii) Multimolecular colloids
- **15.** (*i*) What is the principle behind the zone refining of metals?
 - (ii) What is the role of silica in the extraction of copper?
 - (iii) How is 'cast iron' different from 'pig iron'?
- **16.** Give reasons for the following:
 - (i) N₂ is less reactive at room temperature.
 - (ii) H₂Te is the strongest reducing agent amongst all the hydrides of Group 16 elements.
 - (iii) Helium is used in diving apparatus as a diluent for oxygen.
- 17. (a) Write the hybridization and shape of the following complexes:
 - (*i*) $[CoF_6]^{3-}$

(ii) $[Ni(CN)_4]^{2-}$

(Atomic number: Co = 27, Ni = 28)

- (b) Out of NH₃ and CO, which ligand forms a more stable complex with a transition metal and why?
- **18.** Write the structure of the major product in each of the following reactions:

(ii)
$$CH_3 - CH_2 - CH_2 - CH_3 + KOH$$
 ethanol/heat

(iii)
$$\rightarrow$$
 + CH₃Cl \rightarrow Anhyd. AlCl₃

- **19.** Give reasons for the following:
 - (i) Phenol is more acidic than ethanol.
 - (ii) Boiling point of ethanol is higher in comparison to methoxymethane.
 - (iii) $(CH_3)_3C O CH_3$ on reaction with HI gives CH_3OH and $(CH_3)_3C I$ as the main products and not $(CH_3)_3C OH$ and CH_3I .
- **20.** How do you convert the following:
 - (i) $C_6H_5CONH_2$ to $C_6H_5NH_2$
- (ii) Aniline to phenol
- (iii) Ethanenitrile to ethanamine

OR

Write the chemical equations involved when aniline is treated with the following reagents:

(i) Br₂ water

(ii) CHCI₃ + KOH

- (iii) HCl
- 21. Write the names and structures of the monomers of the following polymers:
 - (i) Buna-S

(ii) Glyptal

- (iii) Polyvinyl chloride
- 22. (i) Write the product obtained when D-glucose reacts with H_2N —OH.
 - (ii) Amino acids show amphoteric behaviour. Why?
 - (iii) Why cannot vitamin C be stored in our body?



23. Seeing the growing cases of diabetes and depression among young children, Mr. Lugani, the principal of one reputed school organised a seminar in which he invited parents and principals. They all resolved this issue by strictly banning junk food in school and introducing healthy snacks and drinks like soap, lassi, milk, etc. in school canteens. They also decided to make compulsory half an hour of daily physical activities for the students in the morning assembly. After six months, Mr. Lugani conducted the health survey in most of the schools and discovered a tremendous improvement in the health of the students.

After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Lugani?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) What are antidepressant drugs? Give an example.
- (iv) Name the sweetening agent used in the preparation of sweets for a diabetic patient.
- 24. (a) A compound 'A' of molecular formula C₂H₃OCl undergoes a series of reactions as shown below. Write the structures of A, B, C and D in the following reactions:

(C₂H₃OCl) A

H₂/Pd-BaSO₄

dil. NaOH

Heat

D

- (b) Distinguish between the following:
 - (i) C₆H₅—COCH₃ and C₆H₅—CHO
 - (ii) Benzoic acid and methyl benzoate
- (c) Write the structure of 2-methylbutanal.

OR

(a) Write the structures of the main products when acetone (CH₃—CO—CH₃) reacts with the following reagents:

(i) Zn—Hg/conc. HCI

(ii) H₂N—NHCONH₂/H⁺

- (iii) CH₃MgBr and then H₃O⁺
- (b) Arrange the following in the increasing order of their boiling points:

C₂H₅OH, CH₃—CHO, CH₃—COOH

(c) Give a simple chemical test to distinguish between the following pair of compounds:

CH₃CH₂CHO and CH₃CH₂COCH₃

25. Calculate e.m.f. and *G* for the following cell:

 $Mg(s) | Mg^{2+}(0.001 M) | Cu^{2+}(0.0001 M) | Cu(s)$

Given:
$$E_{\text{(Mg}^{2+}/\text{Mg)}}^{\text{o}} = -2.37 \text{ V}, E_{\text{(Cu}^{2+}/\text{Cu)}}^{\text{o}} = +0.34 \text{ V}.$$

- (a) The conductivity of 0.20 mol L⁻¹ solution of KCI is 2.48×10^{-2} S cm⁻¹. Calculate its molar conductivity and degree of dissociation (). Given ${}^{0}(K) = 73.5 \text{ S cm}^{2} \text{ mol}^{-1}$ and ${}^{0}(Cl^{-}) = 76.5 \text{ S cm}^{2} \text{mol}^{-1}$.
- (b) What type of battery is mercury cell? Why is it more advantageous than dry cell?
- **26.** (a) Account for the following:
 - (i) Zr and Hf have almost similar atomic radii.
 - (ii) Transition metals show variable oxidation states.
 - (iii) Cu⁺ion is unstable in aqueous solution.
 - (b) Complete the following equations:
 - (i) $2MnO_2 + 4KOH + O_2$
 - (ii) $2Na_2CrO_4 + 2H^+$



(*a*)

E _(M²⁺/M)	Cr	Mn	Fe	Co	Ni	Cu
M^{2+}/M	-0.91	-1.18	-0.44	-0.28	-0.25	+0.34

From the given data E⁰ values, answer the following questions:

- (i) Why is $E_{(Cu^{2+}/Cu)}^{o}$ value exceptionally positive?
- (ii) Why is $E_{(Mn^{2+}/Mn)}^{o}$ value highly negative as compared to other elements?
- (iii) Which is a stronger reducing agent Cr^{2+} or Fe^{2+} ? Give reason.
- (b) Why do actinoids show a wide range of oxidation states? Write one similarity between the chemistry of lanthanoids and actinoids.



Solutions •

1. $Zn^{2+} + 2e$ Zn

Quantity of charge required for the reduction of one mole of Zn^{2+} 2F = 2 96500 C 193000 C

2. Dispersed phase: Liquid

Dispersion medium: Solid

- 3. $Cu^{2+}(3d^9 4s^0)$ has one unpaired electron in *d*-subshell which absorbs radiation in visible region resulting in d-d transition and hence Cu²⁺ salts are coloured. Zn²⁺ (3 d^{10} 4 s^0) has completely filled d-orbitals. No radiation is absorbed for d-d transition and hence Zn^{2+} salts are colourless.
- **4.** 2-methylprop-2-en-1-ol.

 CH_3

CH₃, as the tertiary carbonium ion formed in the slowest step is more stable than primary Br

carbonium ion.

6. The addition of a non-volatile solute to a volatile solvent lowers its vapour pressure. In order to boil the solution, i.e., to make its vapour pressure equal to atmospheric pressure, the solution has to be heated at a higher temperature. In other words, the boiling point of solution becomes higher than solvent.

As elevation in boiling point depends on the number of moles of solute particles and independent of their nature therefore it is a colligative property.

7. Rate of a reaction may be defined as the change in concentration of reactant or product in a unit time.

For a hypothetical reaction, R


Rate
$$\frac{[R]}{t}$$
 $\frac{[P]}{t}$

Factors that affect rate of a reaction:


- (i) Concentration of reactants
- (ii) Nature of the reactants
- (iii) Temperature
- (iv) Catalyst
- (v) Surface area of the reactants

(any two)

8. (*i*)



Sulphurous acid (H₂SO₃)



Square pyramidal

9. Dichloridobis-(ethane-1, 2-diamine) platinum (IV) ion

Geometrical isomerism. The cis-geometrical isomer will also show optical isomerism.

OR

- (i) [Co(NH₃)₆]Cl₃ (ii) K₂[NiCl₄]



10. (*i*)
$$C_6H_5NH_2 < C_6H_5NHCH_3 < C_6H_5$$
 CH_2 NH_2

$$(ii) \bigvee_{\mathrm{NO}_2}^{\mathrm{NH}_2} < \bigvee_{\mathrm{CH}_3}^{\mathrm{NH}_2} < \bigvee_{\mathrm{CH}_3}^{\mathrm{NH}_2}$$

11. Substituting W_B 31 g, W_A 500 g, M_B 62 g mol ¹, K_f 1.86 K kg mol ¹ in the expression,

$$T_f = \frac{K_f - W_B - 1000}{M_B - W_A}$$
, we get
$$T_f = \frac{1.86 \text{ K kg mol}^{-1} - 31 \text{ g} - 1000 \text{ g kg}^{-1}}{62 \text{ g mol}^{-1} - 500 \text{ g}}$$

Freezing point of solution, T_f T_f T_f 273.15 K – 1.86 K *T_f* 271.29 K

- (i) When a unit cell contains constituent particles only at the corner positions, it is called as primitive unit **12.** cell.
 - (ii) If in an ionic solid of the type AB, equal number of cations and anions are missing from their usual positions so that the electrical neutrality is maintained, it is called Schottky defect.
 - (iii) The substances which show magnetism even in the absence of magnetic field are called ferromagnetic substances and this property is known as ferromagnetism, e.g., Fe, Co, CrO₂, etc. Ferromagnetism is due to permanent alignment of domains of metal ions in the same direction.

Ferromagnetism

13. Substituting k_1 2 10 ², k_2 4 10 ², T_1 300 K, T_2 310 K, R 8.314 J K ¹ mol ¹ in the expression

$$\log \frac{k_2}{k_1} = \frac{E_a}{2303} = \frac{T_2 - T_1}{T_1 T_2} \text{, we get}$$

$$\log \frac{4 - 10^{-2}}{2 - 10^{-2}} = \frac{E_a}{2303} = \frac{310 - 300}{300 - 310}$$

$$\log 2 = \frac{E_a}{19147} = \frac{10}{300 - 310}$$

$$E_a = 0.3010 - 19.147 - 300 - 31 - 53598 \text{ J mol}^{-1}$$

$$E_a = 53.598 \text{ kJ mol}^{-1}$$

- 14. (i) The motion of the colloidal particles in a zig-zag path due to unbalanced bombardment by the particles of dispersion medium is called Brownian movement.
 - (ii) Peptization: The process of converting a precipitate into colloidal sol by shaking it with dispersion medium in the presence of a small amount of suitable electrolyte is called peptization. During peptization, the precipitate absorbs one of the ions of the electrolyte on its surface. This causes development of positive or negative charge on precipitates, which ultimately break up into particles of colloidal dimension.
 - (iii) Multimolecular colloids: A large number of atoms or smaller molecules (diameter < 1 nm) of a substance on dissolution aggregate together to form species having size in the colloidal range. Such species are called multimolecular colloids. Examples: a sulphur sol consist of particles containing thousands of S₈ sulphur molecules, a platinum or gold sol may have particles of various sizes having many atoms.



- (i) Zone refining is based on the principle that the impurities are more soluble in melt than in the solid state 15. of the metal.
 - (ii) Refer to Ans. 12 (ii) Allahabad Region-2015.
 - (iii) The iron obtained from blast furnace is called pig iron. It contains about 4% carbon and other impurities such as S, P, Si, Mn in smaller amount. Cast iron is made by melting pig iron and coke using hot blast of air. It has slightly lower carbon content (about 3%) and is extremely hard brittle.
- (i) Due to presence of a triple bond between the two N atoms the bond enthalpy (941.4 kJ mol ¹) of N N is very high and hence N₂ is less reactive at room temperature.
 - (ii) Because of lowest bond dissociation enthalpy of Te-H bond due to larger size of Te.
 - (iii) Because of its low solubility in blood as it has large value of K_H .
- (i) Refer to Ans. 14 (iii) Allahabad Region 2015.
 - (ii) $Ni^{2+} = 3d^84s^0$ $[Ni(CN)_4]^{2-}$ XX = Electron pair from ligand CNT Low spin complex

Hybridisation = dsp^2 ; shape = square planar

(b) CO forms more stable complex due to synergic bonding.

Br

18. (i)
$$CH_3$$
 CH C $CH_3 + H$ Br CH_3 CH_2 C CH_3 CH_3

(iii) CH₃ CH₂ CH₂ CH CH₃ + KOH

Br

$$CH_3 CH_2 CH CH CH_3$$

$$EH_3 CH_2 CH_2 CH CH_3$$

$$CH_3 CH_2 CH_2 CH CH_2$$

$$EH_3 CH_3 CH_3 CH_3$$

$$EH_4 CH_3 CH_3$$

$$EH_4 CH_3 CH_3$$

$$EH_5 CH_3$$

$$EH_6 CH_6$$

$$EH_6$$

$$EH_6 CH_6$$

$$EH_6$$

Ethanol / Heat

- 19. (i) Phenol is stronger acid than ethanol, because the phenoxide ion left after the removal of a proton is stabilized by resonance whereas the ethoxide ion left after the removal of proton from ethanol is not. Moreover, ethyl group due to its I effect intensify -ve charge on oxygen atom in ethoxide ion and destabilize it.
 - (ii) Ethanol undergoes intermolecular hydrogen bonding due to presence of a hydrogen attached to electronegative oxygen atom and hence exists as associated molecules. On the other hand, methoxy methane does not form hydrogen bonds.
 - (iii) Refer to Ans. 19 (iii) Allahabad Region 2015.
- $C_6H_5NH_2$ $K_2CO_3 + 2KBr + 2H_2O$ Aniline **20.** (*i*) $C_6H_5CONH_2 + Br_2 + 4KOH$ Benzamide



$$(ii) \begin{picture}(100,0) \put(0.5,0){\line(1,0){100}} \put(0.5,0){\line(1,$$

$$\begin{array}{ccc} (\it{iii}) & \rm{CH_3C} & \rm{N+4[H]} & \begin{array}{c} \rm{Na-Hg\,/C_2H_5OH} \\ & \rm{or\,\,LiAlH_4} \end{array} & \begin{array}{c} \rm{CH_3\,CH_2\,NH_2} \\ \rm{Ethanamine} \end{array}$$

(i)
$$Aniline$$
 $Aniline$ OR $Aniline$ $Aniline$ $Aniline$ OR $Aniline$ $Anil$

(ii)
$$NH_2$$
 2.4,6-Tribromo aniline NC NC $Heat$ $+ 3KCl + 3H_2O$ $Phenyl$ isocyanide

$$(iii) \qquad \begin{array}{c} NH_2 \\ NH_3Cl \\ \end{array}$$

$$Aniline \qquad Anilinium chloride$$

21.

	Polymer	Name of Monomers	Structure of Monomers
(<i>i</i>)	Buna-S	1, 3-butadiene,	CH ₂ CH CH CH ₂
		Styrene	CH=CH ₂
(ii)	Glyptal	Ethylene glycol, Phthalic acid	H O CH ₂ CH ₂ OH COOH COOH
(iii)	Polyvinyl chloride	Vinyl chloride	CH ₂ CH Cl

CHOCH NOH

22. (i)
$$(CHOH)_4 + NH_2OH$$
 $(CHOH)_4$

$$CH_2OH$$

$$Glucose$$

$$CH_2OH$$

$$Glucose oxime$$



In acidic medium COO ion of the zwitter ion accepts a proton to form the cation (I) while in basic medium N H₃ loses a proton to form the cation (II).

- (iii) As vitamin C is water soluble, therefore, it is readily excreted in urine and hence cannot be stored in the body.
- **23.** (*i*) Values displayed by Mr. Lugani are :
 - Caring nature, empathy
 - Concern for health of students
 - Social awareness
 - Application of knowledge of chemistry in daily life.
 - (ii) One can spread awareness about this issue by:
 - performing play in the assembly.
 - performing *nukkad natak* in community.
 - organising competitions like poster making, essay writing, slogan making.
 - conducting seminars.
 - (iii) The drugs which are used for the treatment of depression are called antidepressant. These drugs inhibit the enzymes which catalyse the degradation of noradrenaline, *e.g.*, iproniazid, phenelzine.
 - (iv) Saccharin.

(b) (i) C₆H₅CHO being an aldehyde gives silver mirror with Tollens' reagent while C₆H₅COCH₃ being a ketone does not give silver mirror with Tollens' reagent.

(ii) Benzoic acid gives brisk effervescence with sodium bicarbonate solution while methyl benzoate does not.



OR

(a) (i)
$$CH_3$$
 C $CH_3 + 4[H]$ CH_3 CH_3 CH_4 CH_5 CH

Propanone Propanone semicarbazone

OMgBr OH 0 CH_3 C CH_3 CH_3 C CH_3 (iii) CH₃ C CH₃ + CH₃ MgBr CH_3 CH_3 tert.-Butyl alcohol

- (b) CH_3 $CHO < CH_3CH_2OH < CH_3COOH$
- (c) On adding Tollens' reagent CH₃CH₂CHO forms silver mirror while CH₃CH₂COCH₃ does not.

 Mg^{2+} 2e Mg **25.** At anode :

At cathode : $\frac{\text{Cu}^{2+}}{\text{Mg} + \text{Cu}^{2+}} \frac{2e}{\text{Mg}^{2+} + \text{Cu}}; n = 2$ $E_{\text{cell}}^{0} = \frac{E_{\text{Cu}^{2}}^{0}}{\text{Cu}^{2}} \frac{E_{\text{Mg}^{2}}^{0}}{\text{Mg}} \frac{0.34 \text{ V} - (-2.37 \text{ V})}{\text{Mg}^{2+} + \text{Cu}^{2}}$

2.71 V

• Substituting, E_{cell}^0 2.71 V, n 2, $[\text{Mg}^{2+}]$ 1 10 3 M, $[\text{Cu}^2]$ 1 10 4 M in Nernst equation

$$\begin{split} E_{\rm cell} & E_{\rm cell}^{\rm o} & \frac{0.059}{n} \log \frac{\rm [Mg^2]}{\rm [Cu^2]} \,, \, \text{we get} \\ E_{\rm cell} & 2.71 & \frac{0.059}{2} \log \frac{10^{-3}}{10^{-4}} \end{split}$$

 $E_{\rm cell}$ $2.71 \quad 0.0295 \log 10 \quad 2.71 - 0.0295 = 2.6805 \text{ V}$ 2.68 V $E_{\rm cell}$

Substituting n=2, F=96500 C mol $^{-1}$, $E_{cell}=2.68$ V in the expression $G=nFE_{cell}$, we get 2 96500 C mol ¹ 2.68 V -517240 J mol⁻¹ G -517.24 kJ mol ¹

OR

(a) Substituting $K = 2.48 - 10^{-2} \text{ S cm}^{-1}$, $M = 0.20 \text{ mol L}^{-1}$ in the expression $_{m} = \frac{K - 1000}{M}$, we get $_{m}$ $\frac{2.48 \cdot 10^{-2} \text{ S cm}^{-1} \cdot 1000}{0.20 \text{ mol L}^{-1}}$ $124 \text{ S cm}^{2} \text{ mol}^{-1}$



•
$$m \text{ (KCl)}$$
 $K \text{ Cl}^-$ 73.5 S cm² mol ¹ 76.5 S cm² mol ¹
 $m \text{ (KCl)}$ 150 S cm² mol ¹

Substituting
$$_m$$
 124 S cm² mol¹, $_m^{\circ}$ 150 S cm² mol¹ in the expression $\frac{m}{\circ}$, we get

Degree of dissociation,
$$\frac{124 \text{ S cm}^2 \text{ mol}}{150 \text{ S cm}^2 \text{ mol}} = 0.82666$$

$$82.67\%$$

- (b) Primary cell. Mercury cell is more advantageous than dry cell because its cell potential remains constant during its life as the overall reaction does not involve any ion in solution whose concentration can change during its life period.
- **26.** (a) (i) This is due to filling of 4f orbitals which have poor shielding effect (Lanthanoid contraction).
 - (ii) Transition elements show variable oxidation states as (n 1) d and ns orbital electrons have comparable energies so that both can enter into chemical bond formation.
 - (iii) Refer to Ans. 21 (i) CBSE Allahabad Region 2015.

(ii)
$$2\text{Na}_2\text{CrO}_4 + 2\text{H}^+$$
 $\text{Na}_2\text{Cr}_2\text{O}_7 + 2\text{Na}^+ + \text{H}_2\text{O}$
Sodium Sodium dichromate

OR

- (a) (i) Copper has high enthalpy of atomisation and low enthalpy of hydration. Since the high energy to transform Cu (s) to Cu²⁺ (aq) is not balanced by hydration enthalpy, therefore, $E_{\text{Cu}^2}^{\text{O}}$ value is exceptionally positive.
 - (ii) This is due to extra stability of half-filled 3d-orbitals of $Mn^{2+}(3d^5)$.
 - (iii) Cr^{2+} , d^4 d^3 occurs in case of Cr^{2+} to Cr^{3+} while d^6 d^5 occurs in case of Fe^{2+} to Fe^{3+} . As in a medium like water d^3 is more stable as compared to d^5 , therefore, Cr^{2+} is stronger reducing agent
- (b) Due to comparable energies of 5f, 6d and 7s orbitals actinoids show a wide range of oxidation states. For the answer of remaining part see Ans. 6 Allahabad Region 2015.





CBSE Examination Paper Foreign-2015 (Code No. 56/2/1/F)

Time allowed: 3 hours Maximum marks: 70

General Instructions: Same as CBSE Examination Paper Delhi-2015.

1. Which would undergo S_N 2 reaction faster in the following pair and why?

$$CH_3 - CH_2 - Br$$
 and $CH_3 - CH_2 - I$

- 2. Which allotrope of sulphur is thermally stable at room temperature?
- 3. Write the IUPAC name of the given compound:

$$\begin{array}{ccc} \operatorname{HO} - \operatorname{CH}_2 - \operatorname{CH} & \operatorname{C} - \operatorname{CH}_3 \\ & | & \\ & \operatorname{CH}_2 \end{array}$$

- **4.** What is the formula of a compound in which the element Y forms *ccp* lattice and atoms of X occupy 2/3rd of octahedral voids?
- **5.** Physisorption is reversible while chemisorption is irreversible. Why?
- **6.** (*i*) Write down the IUPAC name of the following complex:

$$[Cr (en)_3]Cl_3$$

(ii) Write the formula for the following complex:

Potassium tri oxalato chromate (III)

7. What is meant by negative deviation from Raoult's law? Give an example. What is the sign of mix H for negative deviation?

OR

Define azeotropes. What type of azeotrope is formed by negative deviation from Raoult's law? Give an example.

8. Name the reagents used in the following reactions:

(i)
$$CH_3 - CHO$$
 ? $CH_3 - CH - CH_3$ | OH

9. (a) Following reactions occur at cathode during the electrolysis of aqueous copper(II) chloride solution:

$$Cu^{2+}(aq) + 2e^{-}$$
 $Cu(s)$ $E^{0} = +0.34 \text{ V}$
 $H^{+}(aq) + e^{-}$ $\frac{1}{2} H_{2}(g)$ $E^{0} = 0.00 \text{ V}$

On the basis of their standard reduction electrode potential (E⁰) values, which reaction is feasible at the cathode and why?

- (b) State Kohlrausch law of independent migration of ions. Write its one application.
- **10.** Why do transition elements show variable oxidation states? In 3d series (Sc to Zn), which element shows the maximum number of oxidation states and why?

- 11. Calculate the mass of NaCl (molar mass = 58.5 g mol^{-1}) to be dissolved in 37.2 g of water to lower the freezing point by 2°C, assuming that NaCl undergoes complete dissociation. (K_f for water = 1.86 K kg mol⁻¹)
- 12. Write the names and structures of the monomers of the following polymers:
 - (i) Terylene
- (ii) Bakelite
- (iii) Buna-S
- **13.** (*i*) Which one of the following is a monosaccharide:

starch, maltose, fructose, cellulose

- (ii) What is the difference between acidic amino acids and basic amino acids?
- (iii) Write the name of the vitamin whose deficiency causes bleeding of gums.
- **14.** (*i*) Indicate the principle behind the method used for the refining of Nickel.
 - (ii) What is the role of dilute NaCN in the extraction of gold?
 - (iii) What is 'copper matte'?
- **15.** Calculate the emf of the following cell at 25°C:

 $Zn | Zn^{2+} (0.001M) | H^{+} (0.01M) | H_{2}(g) (1 bar) | Pt(s)$

$$E^{0}(Zn^{2}/Zn) - 0.76 \text{ V}, E^{0}(H/H_{2}) = 0.00 \text{ V}$$

- **16.** Give reasons for the following observations:
 - (i) A delta is formed at the meeting point of sea water and river water.
 - (ii) NH_3 gas adsorbs more readily than N_2 gas on the surface of charcoal.
 - (iii) Powdered substances are more effective absorbents.
- 17. (i) Draw the geometrical isomers of complex $[Pt(en)_2Cl_2]^{2+}$
 - (ii) On the basis of crystal field theory, write the electronic configuration for d^4 ion, if
 - (iii) Write the hybridization type and magnetic behaviour of the complex $[Ni\ (CN)_4]^{2-}$. (Atomic number of Ni = 28).
- **18.** (a) Why are alkyl halides insoluble in water?
 - (b) Why is Butan-1-ol optically inactive but Butan-2-ol is optically active?
 - (c) Although chlorine is an electron withdrawing group, yet it is ortho-, para-, directing in electrophilic aromatic substitution reactions. Why?
- **19.** How do you convert the following:
 - (a) Benzoic acid to Benzaldehyde
 - (b) Ethyne to Ethanal
 - (c) Acetic acid to Methane

OR

Write the equations involved in the following reactions:

- (i) Stephen reaction
- (ii) Wolff-Kishner reduction
- (iii) Etard reaction
- **20.** (a) How would you account for the following:
 - (i) Highest fluoride of Mn is MnF₄ whereas the highest oxide is Mn₂O₇.
 - (ii) Transition metals and their compounds show catalytic properties.
 - (b) Complete the following equation:

 $3MnO_4^2$ $4H^{+}$



(i)
$$CH_3 - CH = CH_2$$
 (i) B_2H_6 ?

(*ii*)
$$C_6H_5 - OH$$
 $Br_2 (aq)$?

- 22. An element X (molar mass = 60 g mol^{-1}) has a density of 6.23 g cm^{-3} . Identify the type of cubic unit cell, if the edge length of the unit cell is 4×10^{-8} cm.
- 23. Seeing the growing cases of diabetes and depression among children, Mr. Chopra, the principal of one reputed school organised a seminar in which he invited parents and principals. They all resolved this issue by strictly banning the junk food in schools and by introducing healthy snacks and drinks like soup, lassi, milk, etc. in school canteens. They also decided to make compulsory half an hour physical activities for the students in the morning assembly daily. After six months, Mr. Chopra conducted the health survey in most of the schools and discovered a tremendous improvement in the health of students.


After reading the above passage, answer the following questions:

- (i) What are the values (at least two) displayed by Mr. Chopra?
- (ii) As a student, how can you spread awareness about this issue?
- (iii) Why should antidepressant drugs not be taken without consulting a doctor?
- (iv) Give two examples of artificial sweeteners.
- **24.** (*a*) Define the following terms:
 - (i) Activation energy
- (ii) Rate constant
- (b) A first order reaction takes 10 minutes for 25% decomposition. Calculate $t_{1/2}$ for the reaction.

(Given:
$$\log 2 = 0.3010$$
, $\log 3 = 0.4771$, $\log 4 = 0.6021$)

OR

(a) For a chemical reaction R P, the variation in the concentration, ln [R] vs. time (s) plot is given as



- (i) Predict the order of the reaction.
- (ii) What is the slope of the curve?
- (iii) Write the unit of rate constant for this reaction.
- (b) Show that the time required for 99% completion is double of the time required for the completion of 90% reaction.
- **25.** (a) Account for the following:
 - (i) Bond angle in NH₄ is higher than NH₃.
 - (ii) H₂S has lower boiling point than H₂O.
 - (iii) Reducing character decreases from SO_2 to TeO_2 .
 - (b) Draw the structures of the following:
 - (i) H₄P₂O₇ (Pyrophosphoric acid)
 - (ii) XeF₂



OR

- (a) Draw the structures of the following:
 - (i) XeF₄
 - (ii) H₂S₂O₇
- (b) Account for the following:
 - (i) Iron on reaction with HCl forms FeCl₂ and not FeCl₃.
 - (ii) HClO₄ is a stronger acid than HClO.
 - (iii) BiH₃ is the strongest reducing agent amongst all the hydrides of group 15.
- **26.** (a) Illustrate the following reactions giving suitable example in each case:
 - (i) Ammonolysis
 - (ii) Coupling reaction
 - (iii) Acetylation of amines
 - (b) Describe Hinsberg method for the identification of primary, secondary and tertiary amines. Also write the chemical equations of the reactions involved.

OR

- (a) Write the structures of main products when benzene diazonium chloride ($C_6H_5N_2Cl^-$) reacts with the following reagents:
 - (*i*) HBF₄/
 - (ii) Cu/HBr
- (b) Write the structures of A, B and C in the following reactions:
 - A NaNO₂ HCl (i) $C_6H_5NO_2$ 273 K
 - $LiAlH_4$ HNO_2 (ii) CH₃Cl





Solutions -

- 1. CH₃—CH₂—I, as iodine is a better leaving group due to its larger size, it will release at a faster rate in the presence of incoming nucleophile.
- 2. Rhombic sulphur
- **3.** 3-Methylbut-2-en-1-ol.
- **4.** Let the number of atoms of element *Y* in ccp lattice N

Number of octahedral voids

Number of atoms of element $X = \frac{2}{3}$ N

$$X: Y = \frac{2}{3} N: N$$

$$X:Y$$
 2:3

Formula of compound X_2Y_3

- 5. Because physisorption involves weak vander Waal forces whereas chemisorption involves chemical bond formation.
- (i) Tris-(ethane-1,2-diamine) chromium (III) chloride
 - (*ii*) $K_3[Cr(C_2O_4)_3]$
- 7. The deviation is called negative deviation if the partial pressure of each component (A and B) and vapour pressure of solution are less than the pressure expected on the basis of Raoult's law. This type of deviation is shown by the solutions in which F_{A} $_{B} > F_{A}$ $_{A}$ and F_{B} $_{B}$.

A mixture of chloroform and acetone forms a solution with negative deviation from Raoult's law.

Sign of $_{mix}H$ is negative.

OR

Azeotropes are binary liquid mixtures having the same composition in liquid and vapour phase and boil at a constant temperature.

Maximum boiling azeotrope.

A mixture of 68% nitric acid and 32% water by mass is an example of maximum boiling azeotrope.

- **8.** (*i*) CH₃MgBr/H₂O
 - (ii) SOCl₂ or PCl₅
- **9.** (a) The reaction, Cu^2 (aq) Cu(s) is feasible at cathode as it has higher reduction potential.
 - (b) Kohlrausch law states that the limiting molar conductivity of an electrolyte can be represented as the sum of the individual contribution of anion and cation of the electrolyte.

Applications of Kohlrausch law:

(i) In calculation of m for weak electrolyte, e.g.,

$$m(CH_3COOH)$$
 (CH_3COO) H $m(CH_3COONa)$ $m(HCl)$ m (NaCl

- (ii) In calculation of degree of dissociation () and dissociation constant (K) of weak electrolytes.
- **10.** Refer to Ans. 26 (OR) (*a*) (*ii*) Patna Region 2015.

Manganese $(25 \text{Mn} - 3d^5 4s^2)$ from + 2 to + 7, as its atom has maximum number of unpaired electrons.



11. NaCl Na Cl ,
$$i = \frac{2}{1}$$
 2

Substituting K_f 1.86 K kg mol 1 , W_A 37.2 g, M_B 58.5 g mol 1 , i 2, T_f 2 C or 2 K in the equation, W_B $\frac{T_f \quad M_B \quad W_A}{i \quad K_f \quad 1000}$, we get

$$W_B = \frac{2 \text{K}}{12} \frac{58.5 \text{ g mol}}{1.86 \text{ K kg mol}} \frac{1}{1000 \text{ g kg}} \frac{37.2 \text{ g}}{1} = 1.17 \text{ g}$$

12.

	Name of polymer	Name of Monomer	Structure of Monomer
(<i>i</i>)	Terylene	Ethylene glycol	HO—H ₂ C—CH ₂ —OH
		Terephthalic acid	ноос-
(ii)	Bakelite	Phenol	ОН
		Formaldehyde	НСНО
(iii)	Buna-S	1,3-Butadiene	CH ₂ CH—CH CH ₂
		Styrene	CH=CH ₂

13. (*i*) Fructose

- (ii) An acidic amino acid has more number of carboxylic groups than amino groups, e.g., aspartic acid, glutamic acid whereas a basic amino acid has more number of amino groups than carboxyl groups, e.g., lysine, arginine.
- (iii) Vitamin C.
- **14.** (*i*) Impure nickel is heated with CO to form volatile complex Ni(CO)₄ and collected elsewhere. It is then decomposed to give pure nickel.

(ii) NaCN acts as a leaching agent to form soluble complex with gold while the impurities remain unaffected which are then filtered off.

$$4Au(s)$$
 8CN (aq) $2H_2O(l)$ $O_2(g)$ $4[Au(CN)_2]$ (aq) $4OH(aq)$

(iii) Copper matte is a mixture of Cu₂S and FeS.

$$E_{\text{cell}}$$
 E_{H/H_2} $E_{\text{Zn}^2/\text{Zn}}$ 0 (0.76 V) 0.76 V

Substituting $E_{\text{cell}} = 0.76 \text{ V}$, n = 2, $[\text{Zn}^2] = 1 = 10^{-3} \text{ M}$, $[\text{H}] = 1 = 10^{-2} \text{ M}$ in the Nernst equation

$$E_{\text{cell}} - E_{\text{cell}} - \frac{0.059}{n} \log \frac{[\text{Zn}^2]}{[\text{H}]^2}$$
, we get




$$E_{\text{cell}} = 0.76 = \frac{0.059}{2} \log \frac{10^{-3}}{(10^{-2})^2} = 0.76 = 0.0295 \log 10$$
 $E_{\text{cell}} = 0.76 = 0.0295 = 0.7305 \text{ V}$
 $E_{\text{cell}} = 0.73 \text{ V}$

- **16.** (i) Refer to Ans. 3 Allahabad Region 2015.
 - (ii) NH_3 as it has higher critical temperature than N_2 .

Higher the critical temperature more easily the gas can be liquified, i. e., greater is the intermolecular forces of attraction and hence greater will be adsorption.

(iii) Because powdered substances have larger surface area.





$$cis-[Pt (en)_2Cl_2]^{2+}$$

trans- $[Pt (en)_2Cl_2]^{2+}$

(ii)
$$t_{2g}^{4} e_{g}^{0}$$

(*iii*) Ni² :
$$3d^8 4s^0$$

Hybridisation = dsp^2

Magnetic behaviour = Diamagnetic


18. (a) This is due to the inability of alkyl halide molecule to form intermolecular hydrogen bonds with water molecules.

OH

(b) CH₃—CH₂—*C—CH₃, due to presence of a chiral carbon butan-2-ol is an optically active Н

compound.

(c) As the weaker resonance (+R) effect of Cl which stabilise the carbocation formed tends to oppose the stronger inductive (-I) effect of Cl which destabilise the carbocation at ortho and para positions and makes deactivation less for ortho and para position.





O OH H_2O $CH_3 - C - H$ (b) CH CH CH_2 CH $HgSO_4/H_2SO_4$ Ethyne Ethanal

NaOH NaOH / CaO CH₃COONa (c) CH₃COOH CH_4 Acetic acid Methane

OR

 H_3O (i) Stephen reaction: R—C **HCI** SnCl₂ RCH NH **RCHO** Aldehyde Cyanide

(ii) Wolff-kishner reduction:

(iii) Etard reaction:

$$\begin{array}{c|c} CH_3 & CH(OCrOHCl_2)_2 & CHO \\ \hline \\ + CrO_2Cl_2 & H_3O^+ & \\ \hline \\ Toluene & Chromium & Benzaldehyde \\ \hline \\ complex & \\ \end{array}$$

- **20.** (a) (i) As oxygen stabilizes manganese more than fluorine by forming multiple bonds.
 - (ii) The catalytic activity of transition metals and their compounds is attributed to the following reasons:
 - Due to their tendency to show variable oxidation states transition metals form instable intermediate compounds and provide a new path for the reaction with lower activation energy.
 - In some cases, the transition metals provide a suitable large surface area with free valencies on which reactants are adsorbed.
 - (b) 3MnO₄² 4H $2MnO_4^ MnO_2$ $2H_2O$ Manganate ion Permanganate ion

2,4,6-Tribromophenol

$$\begin{array}{cccc} \textit{(iii)} \ \text{CH}_{3}\text{CH}_{2}\text{OH} & & \text{Cu} & & \text{CH}_{3}\text{CHO} & \text{H}_{2} \\ & \text{Ethanol} & & 573 \text{ K} & & \text{Ethanal} \end{array}$$



22. Substituting
$$M = 60 \text{ g mol}^{-1}$$
, $d = 6.23 \text{ g cm}^{-3}$, $a = 4 = 10^{-8} \text{ cm}^{-3}$

$$N_A = 6.022 - 10^{23} \text{ mol}^{-1} \text{ in the expression } z = \frac{d - N_A - a^3}{M}, \text{ we get}$$

$$z = \frac{6.23 \text{ g cm}^{-3} - 6.022 - 10^{23} \text{ mol}^{-1}}{60 \text{ g mol}^{-1}} = \frac{(4 - 10^{-8} \text{ cm})^3}{60} = \frac{240.1}{60}$$

As there are 4 atoms of the element *X* per unit cell. Hence, the cubic unit cell must be face centered cubic (fcc).

- 23. Refer to Ans. 27 Allahabad Region 2015
- **24.** (*a*) (*i*) The energy required to form the intermediate, activated complex is called activation energy. Activation energy = Threshold energy Average energy of the reactants
 - (ii) The rate of reaction when the molar concentration of each of the reactants is taken as unity is called rate constant. Its value depends on the temperature but independent of the concentration of reactants.

(b)
$$[R]$$
 $[R]_0$ 25% of $[R]_0$ $[R]_0$ $\frac{25}{100}$ $\frac{[R]_0}{100}$ $\frac{75[R]_0}{100}$ $\frac{3[R]_0}{4}$

Substituting [R] $\frac{3[R]_0}{4}$, t 10 min, in the expression k $\frac{2.303}{10}$ log $\frac{[R]_0}{[R]}$, we get

$$k = \frac{2.303}{10} \log \frac{[R]_0}{[R]} = \frac{2.303}{10} \log \frac{4}{3}$$

$$k = \frac{2.303}{10} (\log 4 - \log 3) = \frac{2.303}{10} (0.6021 - 0.4771)$$

$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{0.02879 \text{ min}} = 24.07 \text{ min}$$

OR

- (a) (i) First-order
 - (ii) Slope of the curve k (rate constant)

(iii) time
$$^{1}(s^{-1})$$

(b)
$$t = \frac{2303}{k} \log \frac{[R]_0}{[R]}$$

$$\frac{t_{99\%}}{t_{90\%}} = \frac{\frac{2.303}{k} \log \frac{100}{100 - 99}}{\frac{2.303}{k} \log \frac{100}{100 - 90}} = \frac{\log 100}{\log 10} = 2 \frac{\log 10}{\log 10}$$


Hence, $t_{99\%} = 2t_{90\%}$

- 25. (a) (i) N atom both in NH₃ and NH₄ is sp^3 hybridised. In NH₄ all the four orbitals are bonded whereas in NH₃ there is a lone pair on N, which is responsible for lone pair-bond pair repulsion in NH₃ reducing the bond angle from $109^{\circ}28'$ to 107° .
 - (ii) Due to high electronegativity and same size of oxygen, H₂O undergoes extensive intermolecular H-bonding and exist as an associated molecule. On the other hand, due to low electronegativity of S, H₂S is unable to form effective H-bonds and exists as a discrete molecule which are held by weak



vander Waal forces. To overcome these forces only small amount of energy is required. Therefore H_2 S has lower boiling point than H_2 O.

(iii) Because the stability of +4 oxidation state increases from S to Te on moving down the group in group 16 due to inert pair effect.



(i) HCl on reaction with iron produces H₂

2HCl FeCl₂

Liberation of hydrogen prevent the formation of ferric chloride.

HClO₄ H_2O H_3O ClO_4

- (ii) Greater the number of oxygen atom in conjugate base greater will be the dispersal of negative charge through p d back bonding and greater will be its stability. Thus, ClO₄ is more stable than $ClO\,\,$. As the conjugate base ClO_4 is more stable than $ClO\,\,$. Therefore, $HClO_4$ is stronger acid than HClO.
- (iii) As we move down the group, the size of the atom of the element increases and therefore E—H bond strength decreases. Thus, Bi—H bond is the weakest amongst the hydrides of elements of group 15 and hence BiH₃ is the strongest reducing agent.

(i) Ammonolysis:

(ii) Coupling Reaction:



(iii) Acetylation:

$$C_{2}H_{5}-N-H$$

$$C_{-}Cl$$

$$Ethyl amine$$

$$Acetyl chloride$$

$$Acetyl chloride$$

$$Acetic anhydride$$

$$Acetic anhydride$$

$$Acetic anhydride$$

$$Acetic anhydride$$

$$All CH_{3}$$

$$All CH_{3}$$

$$All CH_{3}$$

$$Acetic anhydride$$

- (b) Hinsberg's test: In this test, the amine is first treated with Hinsberg's reagent (benzene sulphonyl chloride) and then shaken with aqueous KOH solution when the three amines behave in different ways.
 - (i) A 1° amine gives a clear solution which on acidification gives an insoluble N-alkyl benzene sulphonamide.

(ii) A 2° amine gives an insoluble N, N-dialkyl benzene sulphonamide which remains unaffected on addition of acid.

$$\begin{array}{c|c} O & R & O & R \\ \hline O & S - Cl & + & H - N - R \\ \hline O & 2^{\circ} \text{ Amine} & O & R \\ \hline O & S - N - R & + & HCl \\ \hline O & N, N-Dialkylbenzene \\ sulphonyl chloride & sulphonamide \\ (insoluble in KOH) \\ \hline \end{array}$$

(iii) A 3° amine does not react at all.

$$\begin{array}{cccc} C_6H_5SO_2Cl & + & R_3N & & \text{KOH} \\ \text{Benzene-sulphonyl} & & 3 & \text{amine} \\ & & \text{chloride} & & & \end{array}$$

(a) (i)
$$N_2Cl + HBF_4 \longrightarrow N_2BF_4 \longrightarrow N_2BF_4 \longrightarrow F + BF_3 + N_2$$

Fluorobenzene

(ii) $N_2Cl \longrightarrow N_2Cl \longrightarrow N_$



(b) (i)
$$A = \bigcup_{Aniline}^{NH_2}$$
; $B = \bigcup_{Benzene diazonium chloride}^{+}$; $C = \bigcup_{Phenol chloride}^{+}$

